(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau





(43) International Publication Date 30 October 2003 (30.10.2003)

(51) International Patent Classification7:

PCT

(10) International Publication Number WO 03/089583 A2

CI2N

(21) International Application Number: PCT/US03/11497

(22) International Filing Date: 15 April 2003 (15.04.2003)

(25) Filing Language: English

(26) Publication Language: English


(30) Priority Data: 60/372,669 16 April 2002 (16.04,2002) US 60/374.823 24 April 2002 (24.04.2002) US 60/376,558 1 May 2002 (01.05.2002) US 60/381.366 20 May 2002 (20.05.2002) US 60/403,648 16 August 2002 (16.08.2002) US 60/411.882 20 September 2002 (20.09,2002) US 60/424.336 7 November 2002 (07.11.2002) US

(71) Applicant (for all designated States except US): ORI-GENE TECHNOLOGIES, INC. [US/US]: Suite 100, 6 Taft Court. Rockville, MD 20850 (US).

- (72) Inventors; and
- (75) Inventors/Applicants (for US only): JAY, Gilbert [US/US]; 5801 Nicholson Lane, North Bethesda, MD 20852 (US). LEBOVITZ, Richard, M. [US/US]; 3800 North Fairfax Drive, Arlington, VA 22203 (US), LIU, Nuan [US/US]; 14213 Day Road, Rockville, MD 20850 (US). SHU, Voumin [US/US]; 2508 Chilham Place. Potomac, MD 20854 (US), SUN, Zairen [CN/US]; 1083 Copperstone Court, Rockville, MD 20852 (US). WU, Meng [PG/US]; 18016 Rockingham Place, Germantown, MD 20874 (US).
- (74) Agent: LEBOVITZ, Richard, M.; Origene Technologies, Inc., Suite 100, 6 Taft Court, Rockville, MD 20850 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU. AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ. DE. DK. DM. DZ, EC. EE, ES, FI, GB, GD, GE, GH. GM. HR. HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ. TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

[Continued on next page]

(54) Title: TISSUE SPECIFIC GENES AND GENE CLUSTERS



(57) Abstract: The present invention relates to genes and genes clusters which are expressed in a tissue specific manner. For example, the invention relates to a group of genes encoding GPCR-like receptors that are involved in the function and activity of the immune system. These genes are organized into a discrete cluster at chromosomal location 1q22 (the "immune gene complex") and span about 700 kb of DNA. The region closest to the centromere comprises genes that are expressed predominantly in the thymus, while the distal region comprises genes which are expressed predominantly in the bone marrow and other hematopoietic cells. Another cluster of GPCR genes is located at chromosomal band 11q24. These genes are expressed predominantly in pancreatic tissue, establishing this region of chromosome 11 as a unique gene complex involved in

pancreatic function. A cluster of transmembrane and GPCR-type receptor genes is also located at chromosomal band 11q12.2. These genes are expressed predominantly in the spleen (hence, "spleen gene" cluster), as well as other tissues of the immune and reticuloendothelial system (RES), indicating that establishing this region of the chromosome is involved is spleen, lymphoid, and/or reticuloendothelial function. Finally, genes coding for membrane proteins have been identified which are expressed selectively in bone marrow, kidney, pancreas, and retina.

BNSDOCID: <WO _ __ 03089583A2_I >



(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE. AG. AL. AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

- as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM). European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

Published:

 without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

10

15

20

25

30

TISSUE SPECIFIC GENES AND GENE CLUSTERS

This application claims the benefit of U.S. Application Serial Nos. 60/372,669 April 16, 2002, 60/374,823 filed April 24, 2002, 60/376,558 filed May 1, 2002, 60/381,366 filed May 20, 2002, 60/403,648 filed August 16, 2002, 60/411,882 filed September 20, 2002, and 60/424,336 filed November 7, 2002, which are hereby incorporated by reference in their entirety.

DESCRIPTION OF THE DRAWINGS

Figs. 1 and 2 show a physical map of the immune system gene complex. Sequence-tagged site ("STS") markers are used to characterize the chromosomal regions. An STS is defined by two short synthetic sequences (typically 20 to 25 bases each) that have been designed from a region of sequence that appears as a single-copy in the human genome (the reference numbers, and the sequences which they represent, are hereby incorporated by reference in their entirety). These sequences can be used as primers in a polymerase chain reaction (PCR) assay to determine whether the site is present or absent from a DNA sample.

Fig. 3 shows the expression pattern of transmembrane proteins homologous to the olfactory G-protein-coupled receptor ("GPCR") family in human tissues. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 5 indicates the SEQ ID NO for each primer ("FOR" is the forward primer and "REV" is the reverse primer).

Fig. 4 shows the expression pattern of two olfactory G-protein-coupled receptor ("GPCR") family members in human tissues. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 6 indicates the SEQ ID NO for each primer ("FOR" is the forward primer and "REV" is the reverse primer).

Figs. 5 (a and b) and 6 show the expression pattern in human tissues of genes selectively expressed in kidney tissue. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 11 indicates the SEQ ID NO for each primer ("FOR" is the forward primer and "REV" is the reverse primer).

Fig. 7 (a-b) show organization of pancreatic gene complex on chromosome 11q24.

10

15

20

25

30

Fig. 8 is a schematic drawing of five of the pancreatic olfactory G-protein-coupled receptor ("GPCR") family members located in the gene complex showing regions of overlap. The numbering underneath the lines indicates amino acid position.

Fig. 9 (a and b) show the expression pattern of TMD0986, XM_061780 (TMD0987), XM_061781 (TMD0353), XM_061784 (TMD0989), and XM_061785 (TMD058) in human tissues. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 12 indicates the SEQ ID NO for each primer ("FOR" is the forward primer and "REV" is the reverse primer).

Fig. 10 shows the expression pattern of TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), and TMD0621 (XM_166205) in human tissues. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 17 indicates the SEQ ID NO for each primer ("F-oligo" is the forward primer and "R-oligo" is the reverse primer).

Fig. 11 shows the organization of the spleen gene complex on chromosome 11q12.2.

Fig. 12 (a-c) shows the expression of the pancreas genes in human tissues. To detect gene expression, PCR was carried out on aliquots of the normalized tissue samples using a forward and reverse gene-specific primers. Table 23 indicates the SEQ ID NO for each primer ("FOR" is the forward primer and "REV" is the reverse primer).

Expression patterns were analyzed as described below. A twenty-four tissue panel was used (lanes from left to right): 1, adrenal gland; 2, bone marrow; 3, brain; 4, colon; 5, heart; 6, intestine; 7, pancreas; 8, liver; 9, lung; 10, lymph node; 11, lymphocytes; 12, mammary gland; 13, muscle; 14, ovary; 15, pancreas; 16, pituitary; 17, prostate; 18, skin; 19, spleen; 20, stomach; 21, testis; 22, thymus; 23, thyroid; 24, uterus. The lane at the far left of each panel contains molecular weight standards. Polyadenylated mRNA was isolated from tissue samples, and used as a template for first-strand cDNA synthesis. The resulting cDNA samples were normalized using beta-actin as a standard. For the normalization procedure, PCR was performed on aliquots of the first-strand cDNA using beta-actin specific primers. The PCR products were visualized on an ethidium bromide stained agarose gel to estimate the quantity of beta-actin cDNA present in each sample. Based on these estimates, each sample was diluted with buffer until each contained the same quantity of beta-actin cDNA per unit volume. PCR was carried out using the primers described above, and reaction

-3-

products were loaded on to an agarose (e.g., 1.5-2%) gel and separated electrophoretically.

DESCRIPTION OF THE INVENTION

The present invention relates to tissue-selective genes and tissue-selective gene clusters. The polynucleotides and polypeptides are useful in variety of ways, including, but not limited to, as molecular markers, as drug targets, and for detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, determining predisposition to, etc., diseases and conditions, associated with genes of the present invention. The identification of specific genes, and groups of genes, expressed in pathways physiologically relevant to particular tissues, permits the definition of functional and disease pathways, and the delineation of targets in these pathways which are useful in diagnostic, therapeutic, and clinical applications. The present invention also relates to methods of using the polynucleotides and related products (proteins, antibodies, etc.) in business and computer-related methods, e.g., advertising, displaying, offering, selling, etc., such products for sale, commercial use, licensing, etc.

Immune Gene Complex

5

10

15

20

25

30

The present invention relates to a group of genes involved in the function and activity of the immune system. These genes are organized into a discrete cluster at chromosomal location 1q22 (the "immune gene complex") and span hundreds of kb of DNA, e.g., about 700 kb of DNA. See, Figs. 1 and 2. The region closest to the centromere comprises genes that are expressed predominantly in the thymus, while the distal region comprises genes which are expressed predominantly in the bone marrow and other hematopoietic cells.

The present invention relates to a composition consisting essentially of the 1q22 immune gene complex, comprising TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) genes, or a fragment thereof comprising at least two said genes. As discussed in more detail, the composition can comprise or consist essentially of the chromosome region between STS markers that define the genomic DNA, e.g., between SHGC-81033 and SHGC-145403, or a fragment thereof comprising at least two said genes.

10

15

20

25

30

The CD1 family, a cluster of genes previously identified as coding for proteins involved in antigen presentation (Sugita and Brenner, Seminars in Immunology, 12:511-516, 2000), are located at the proximal boundary of the immune gene complex. The expression of CD1a, b, and c genes are restricted to professional antigen-presenting cells, including dendritic cells and some B-cell subsets (Sugita and Brenner, ibid). CD1d is present on other cell types, in addition to hematopoietic cells, such as intestinal cells (Sugita and Brenner, ibid).

Adjacent to the CD1 family, is a cluster of genes coding for transmembrane proteins homologous to the olfactory G-protein-coupled receptor ("GPCR") family. These genes include XM_060945 (TMD0024), XM_060346 (TMD1779), XM_060947 (TMD0884), and XM_060948 (TMD0025), and are expressed predominantly in thymus tissues (e.g., thymocytes). XM_089421 (TMD1781) is also expressed in thymus, but it is present in much higher amounts in lymphocytes ("PBL"). This chromosomal region can be defined by STS markers, e.g., between SHGC-81033 and D1S3249, G15944, GDB:191077, GDB:196442, RH68459, RH102597, RH69635, or RH65132, or fragments thereof, such as fragments which comprise two or more genes.

The gene for human erythroid alpha spectrin (SPTA1) is distal to the GPCR thymus-restricted family. It is expressed in bone marrow cells, and is localized to the red cell membrane (Wilmotte et al., Blood, 90(10):4188-96, 1997). Next to it, is another cluster of genes coding for proteins that resemble the olfactory GPCR family. These include XM_060956 (TMD0304), XM_060957 (TMD0888), and XM_060959 (TMD089), and are expressed predominantly in the bone marrow, although other sites of expression are observed as well. See, e.g., Table 1. This chromosomal region can be defined by STS markers, e.g., between GDB:181583 or RH118729, and D1S2577 or SHGC-145403.

The gene for myeloid cell nuclear differentiation antigen ("MNDA") is next. MNDA is also expressed in bone marrow cells, particularly in normal and neoplastic myelomonocytic cells and a subset of normal and neoplastic B lymphocytes (Miranda et al., Hum. Pathol., 30(9):1040-9, 1999).

The phrase "immune system" indicates any processes and cells which are involved in generating and carrying out an immune response. Immune system cells includes, but are not limited to, e.g., stem cells, pluripotent stem cell, myeloid progenitor, lymphoid progenitor,

WO 03/089583 PCT/US03/11497
-5-

lymphocytes, B-lymphocytes, T-lymphocytes (e.g., naive, effector, memory, cytotoxic, etc.), thymocytes, natural killer, erythroid, megakaryocyte, basophil, eosinophil, granulocytemonocyte, accessory cells (e.g., cells that participate in initiating lymphocyte responses to antigens), antigen-presenting cells ("APC"), mononuclear phagocytes, dendritic cells, macrophages, alveolar macrophages, etc., and any precursors, progenitors, or mature stages thereof.

Table I is a summary of the genes and their expression patterns in accordance with the present invention. The genes and the polypeptides they encode can be used as diagnostic, prognostic, therapeutic, and research tools for any conditions, diseases, disorders, or applications associated with the tissues and cells in which they are expressed.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

In view of their selectivity and display on the cell surface, the olfactory GPCR family members of the present invention are a useful target for histological, diagnostic, and therapeutic applications relating to the cells in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to breast cancer. They can also be used to detect metastatic cells, in biopsies to identify bone marrow and thymus tissue, etc. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc.

Useful antibodies or other binding partners include those that are specific for parts of the

5

10

15

20

25

polypeptide which are exposed extracellularly as indicated in Table 2. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo (e.g., bone marrow cells or peripheral blood lymphocytes can be treated ex vivo and then returned to the body).

The expression patterns of the selectively expressed polynucleotides disclosed herein can be described as a "fingerprint" in that they are a distinctive pattern displayed by a tissue. Just as with a fingerprint, an expression pattern can be used as a unique identifier to characterize the status of a tissue sample. The list of expressed sequences disclosed herein provides an example of such a tissue expression profile. It can be used as a point of reference to compare and characterize samples. Tissue fingerprints can be used in many ways, e.g., to classify an unknown tissue, to determine the origin of metastatic cells, to assess the physiological status of a tissue, to determine the effect of a particular treatment regime on a tissue, to evaluate the toxicity of a compound on a tissue of interest, etc.

For example, the tissue-selective polynucleotides disclosed herein represent the configuration of genes expressed by a normal tissue. To determine the effect of a toxin on a tissue, a sample of tissue can be obtained prior to toxin exposure ("control") and then at one or more time points after toxin exposure ("experimental"). An array of tissue-selective probes can be used to assess the expression patterns for both the control and experimental samples. As discussed in more detail below, any suitable method can be used. For instance, a DNA microarray can be prepared having a set of tissue-selective genes arranged on to a small surface area in fixed and addressable positions. RNA isolated from samples can be labeled using reverse transcriptase and radioactive nucleotides, hybridized to the array, and then expression levels determined using a detection system. Several kinds of information can be extracted: presence or absence of expression, and the corresponding expression levels. The normal tissue would be expected to express substantially all the genes represented by the tissue-selective probes. The various experimental conditions can be compared to it to

While the expression profile of the complete gene set represented by the sequences disclosed here may be most informative, a fingerprint containing expression information from less than the full collection can be useful, as well. In the same way that an incomplete fingerprint may contain enough of the pattern of whorls, arches, loops, and ridges, to identify

determine whether a gene is expressed, and how its levels match up to the normal control.

5

10

15

20

25

10

15

20

25

30

the individual, a cell expression fingerprint containing less than the full complement may be adequate to provide useful and unique identifying and other information about the sample. Moreover, because of heterogeneity of the population, as well differences in the particular physiological state of the tissue, a tissue's "normal" expression profile is expected to differ between samples, albeit in ways that do not change the overall expression pattern. As a result of these individual differences, each gene although expressed selectively in spleen, may not on its own 100% of the time be adequately enough expressed to distinguish said tissue. Thus, the genes can be used in any of the methods and processes mentioned above and below as a group, or one at a time.

Binding partners can also be used as to specifically deliver therapeutic agents to a tissue of interest. For example, a gene to be delivered to a tissue can be conjugated to a binding partner (directly or through a polymer, etc.), in liposomes comprising cell surface, and then administered as appropriate to the subject who is to be treated. Additionally, cytotoxic, cytostatic, and other therapeutic agents can be delivered specifically to the tissue to treat and/or prevent any of the conditions associated with the tissue of interest.

The present invention relates to methods of detecting immune system cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for a gene selected from Table 1, or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 3, 4, 8, 9, 14, 15, 22, 23, 27, 28, 35, 36, 42, 43, 49, 50, 57, and 58 (see, Table 5), and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting an immune system cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by gene selected from Table 1, or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be

10

15

20

25

30

accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface as indicated in Table 2.

As indicated above, binding partners can be used to deliver agents specifically to the immune system, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to an immune cell can comprise, e.g., contacting an immune cell with an agent coupled to binding partner specific for a gene selected from Table 1 (i.e., TMD0024 (XM 060945), TMD1779 (XM 060946), TMD0884 (XM 060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM 060956), TMD0888 (XM 060957), and TMD0890 (XM 060959)), whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the immune system can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose.

The maturation of the immune system can also be modulated in accordance with the

10

15

20

25

30

present invention, e.g., by methods of modulating the maturation of an immune system cell, comprising, e.g., contacting said cell with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from Table 1, or a mammalian homolog thereof, whereby the maturation of an immune cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

The phrase "immune system cell maturation" includes indirect or direct effects on immune system cell maturation, i.e., where modulating the gene directly effects the maturational process by modulating a gene in a immune system cell, or less directly, e.g., where the gene is expressed in a cell-type that delivers a maturational signal to the immune system cell. Immune system maturation includes B-cell maturation, T-cell maturation, such as positive selection, negative selection, apoptosis, recombination, expression of T-cell receptor genes, CD4 and CD8 receptors, antigen recognition, MHC recognition, tolerization, RAG expression, differentiation, TCR expression, antigen expression, etc. See also below and, e.g., Abbas et al., *Cellular and Molecular Immunology*, 4th Edition, W.B. Saunders Company, 2000, e.g., Pages 149-160. Process include reception of a signal, such as cytokinin or other GPCR ligand. Any suitable agent can be used, e.g., agents that block the maturation, such as an antibody to a GPCR of Table 1, or other GPCR antagonist.

The interactions between lymphoid and non-lymphoid immune system cells can also be modulated comprising, e.g., contacting said cells with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from Table 1, or a mammalian homolog thereof, whereby the interaction is modulated. Lymphoid cells, includes, e.g., lymphocytes (T- and B-), natural killer cells, and other progeny of a lymphoid progenitor cell. Non-lymphoid cells include accessory cells, such as antigen presenting cells, macrophages, mononuclear phagocytes dendritic cells, non-lymphoid thymocytes, and other cell types which do not normally arise from lymphoid progenitors. Interactions that can be modulated included, e.g., antigen presentation, positive selection, negative selection, progenitor cell differentiation, antigen expression, tolerization, TCR expression, apoptosis. See, also above and below, for other immune system processes.

Promoter sequences obtained from GPCR genes of the present invention can be utilized to selectively express heterologous genes in immune system cells. Methods of

-10-

expressing a heterologous polynucleotide in immune system cells can comprise, e.g., expressing a nucleic acid construct in immune system cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected from Table 5. In addition to the cell lines mentioned below, the construct can be expressed in primary cells, such as thymocytes, bone marrow cells, stem cells, lymphoid progenitor cells, myeloid progenitor cells, monocytes, antigen presenting cells, macrophages, and cell lines derived therefom, cell lines such as JHK3 (CRL-10991), KG-1 (CCL-246), KG-1a (CCL-246.1), U-937 (CRL-1593.2), VA-ES-BJ (CRL-2138), TUR (CRL-2367), ELI (CRL-9854), 28SC (CRL-9855), KMA (CRL-9856), THP-1 (TIB-2002), WEHI-274.1 (CRL-1679), M-NFS-60 (CRL-1838), MH-S (CRL-2019), SR-4987 (CRL-2028), NCTC 3749 (CCL-461), AMJ2-C8 (CRL 2455), AMJ2-C11 (CRL2456), PMJ2-PC (CRL-2457), EOC2 (CRL-2467), as well as any primary and established immune system cell lines.

15 Thymus

5

10

20

25

30

The thymus is the site of T-cell lymphocyte maturation. Immature lymphocytes migrate into the thymus from the bone marrow and other organs in which they are generated. The selection process that shape the antigen repertoire of T-cells takes place in the thymus organ. Both positive and negative selection processes take place. For a review, see, e.g., Abbas et al., Cellular and Molecular Immunology, 4th Edition, W.B. Saunders Company, 2000, e.g., Pages 126-130 and 149-160.

There are various diseases and disorders related to thymus tissue, including, but not limited to, thymic carcinoma, thymoma, Omenn syndrome, autoimmune diseases, allergy, Graves disease, Myasthenia gravis, thymic hyperplasia, DiGeorge syndrome, Good syndrome, promoting immune system regeneration after bone marrow transplantation, immuno-responsiveness, etc. The thymic selective genes and polypeptides encoded thereby can be use to treat or diagnose any thymic condition. For instance, chemotherapeutic and cytotoxic agents can be conjugated to thymic selective antibodies and used to ablate a thymoma or carcinoma. They can be used alone or in combination with other treatments. See, e.g., Graeber and Tamin, Semin. Thorac. Cardiovasc. Surg., 12:268-277, 2000; Loehrer, Ann. Med., 31 Suppl. 2:73-79, 1999.

Bone marrow

5

10

15

20

25

30

All circulating blood cells in the adult, including all immature lymphocytes, are produced in the bone marrow. In addition, the bone marrow is also the site of B-cell maturation. The marrow consists of a spongelike reticular framework located between long trabeculae. It is filled with fat cells, stromal cells, and precursor hematopoietic cells. The precursors mature and exit through the vascular sinuses

All the blood cells are believed to arise from a common stem cell. Lineages that develop from this common stem cell include, e.g., myeloid and lymphoid progenitor cells. The myeloid progenitor develops into, erythrocytes (erythroid), platelets (megokaryocytic), basophils, eosinophils, granulocytes, neutrophils, and monocytes. The lymphoid progenitor is the precursor to B-lymphocytes, T-lymphocytes, and natural killer cells.

There are various diseases and disorders related to bone marrow, including, not limited to, e.g., red cell diseases, aplastic anemia (e.g., where there is a defect in the myeloid stem cell), pure red cell aplasia, white cell diseases, leukopenia, neutropenia, reactive (inflammatory) proliferation of white cells and nodes such as leukocytosis and lymphadenitis, neoplastic proliferation of white cells, malignant lymphoma, Non-Hodgkin's Lymphomas, Hodgkins disease, acute leukemias (e.g., acute lymphoblastic leukemia, acute myeloblastic leukemia, myelodysplatic snydrome), chromic myeloid leukemia, chronic leukemia. hairy cell leukemia, myeloproliferative disorders, plasma cell disorders, multiple myeloma, histiocytoses, etc.

Immune System Selective Genes

The present invention relates to genes involved in the function and activity of the immune system. XM_062147 (TMD0088) and XM_061676 (TMD0045) code for seven membrane spanning polypeptides which are homologous to members of the olfactory G-protein-coupled receptor ("GPCR") family. XM_062147 is expressed predominantly in bone marrow tissue, with no detectable expression in other tissues. XM_061676 is also expressed predominantly in bone marrow tissue, but it is detected in peripheral blood lymphocytes, as well. As discussed in more detail below, XM_062147 (TMD0088), XM_061676 (TMD0045), and the polypeptides they encode, can be used as diagnostic, prognostic, therapeutic, and research tools for any conditions, diseases, disorders, or applications

PCT/US03/11497

WO 03/089583

-12-

associated with the immune system and the cells in which they are expressed.

In view of their selectivity and display on the cell surface, the GPCR family members of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to the cells (e.g., B-cells and B-cell progenitors) in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to breast cancer. They can also be used to detect metastatic cells, in biopsies to identify bone marrow, lymphocytes, etc. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly as indicated in Table 2. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo (e.g., bone marrow cells or peripheral blood lymphocytes can be treated ex vivo and then returned to the body). Ex vivo methods can be used to eliminate cancerous cells from the bone marrow, to modulate bone marrow cells, to prime bone marrow cells for an immune response, to expand a particular class of cells expressing XM_062147 (TMD0088) or XM_061676 (TMD0045), to transfer genes into said cells (e.g., Banerjee and Bertino, Lancet Oncol., 3:154-158, 2002), etc.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

The phrase "immune system" indicates any processes and cells which are involved in

5

10

15

20

25

generating and carrying out an immune response. Immune system cells includes, but are not limited to, e.g., stem cells, pluripotent stem cell, myeloid progenitor, lymphoid progenitor, lymphocytes, B-lymphocytes, T-lymphocytes (e.g., naive, effector, memory, cytotoxic, etc.), thymocytes, natural killer, erythroid, megakaryocyte, basophil, eosinophil, granulocytemonocyte, accessory cells (e.g., cells that participate in initiating lymphocyte responses to antigens), antigen-presenting cells ("APC"), mononuclear phagocytes, dendritic cells, macrophages, etc., and any precursors, progenitors, or mature stages thereof.

XM_062147 contains seven transmembrane segments. It is located on chromosomal band 11q12 within proximity to the locus for an inherited form of atopic hypersenstivity (OMIM 147050, e.g., associated with asthma, hay fever, and eczema). It has been suggested that the condition is a result of defect in the regulation of immunoglobulin E. XM_061676 also is seven membrane spanning polypeptide. The chromosomal locus, 11p15, to which it maps is rich in genes associated with immune disorders, including Fanconi anemia, nucleoporin, myeloid leukemia, and T-cell lymphoblastic leukemia. Arthrogryposis multiplex congenita (distal type IIB) also maps closely to this chromosomal location.

The present invention relates to methods of detecting immune system cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for a gene selected from Table 6, or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 67, 68, 76, and 77 (see, Table 6), and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting an immune system cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by gene selected from Table 6, or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be

5

10

15

20

25

accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc.

Useful epitopes include those exposed to the surface as indicated in Table 7.

As indicated above, binding partners can be used to deliver agents specifically to the immune system, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to an immune cell can comprise, e.g., contacting an immune cell with an agent coupled to binding partner specific for a gene selected from Table 6, whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the immune system can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos. 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose.

The maturation of the immune system can also be modulated in accordance with the present invention, e.g., by methods of modulating the maturation of an immune system cell, comprising, e.g., contacting said cell with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from Table 6, or a mammalian homolog thereof,

5

10

15

20

25

WO 03/089583 PCT/US03/11497 -15-

whereby the maturation of an immune cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

The phrase "immune system cell maturation" includes indirect or direct effects on immune system cell maturation, i.e., where modulating the gene directly effects the maturational process by modulating a gene in a immune system cell, or less directly, e.g., where the gene is expressed in a cell-type that delivers a maturational signal to the immune system cell. Immune system maturation includes B-cell maturation, T-cell maturation, such as positive selection, negative selection, apoptosis, recombination, expression of T-cell receptor genes, CD4 and CD8 receptors, antigen recognition, MHC recognition, tolerization, RAG expression, differentiation, TCR expression, antigen expression, etc. See also below and, e.g., Abbas et al., *Cellular and Molecular Immunology*, 4th Edition, W.B. Saunders Company, 2000, e.g., Pages 149-160. Processes include reception of a signal, such as cytokinin or other GPCR ligand. Any suitable agent can be used, e.g., agents that block the maturation, such as an antibody to a GPCR of Table 6, or other GPCR antagonist.

The interactions between lymphoid and non-lymphoid immune system cells can also be modulated comprising, e.g., contacting said cells with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from Table 6, or a mammalian homolog thereof, whereby the interaction is modulated. Lymphoid cells, includes, e.g., lymphocytes (T- and B-), natural killer cells, and other progeny of a lymphoid progenitor cell. Non-lymphoid cells include accessory cells, such as antigen presenting cells, macrophages, mononuclear phagocytes dendritic cells, non-lymphoid thymocytes, and other cell types which do not normally arise from lymphoid progenitors. Interactions that can be modulated included, e.g., antigen presentation, positive selection, negative selection, progenitor cell differentiation, antigen expression, tolerization, TCR expression, apoptosis. See, also above and below, for other immune system processes.

Promoter sequences obtained from GPCR genes of the present invention can be utilized to selectively express heterologous genes in immune system cells. Methods of expressing a heterologous polynucleotide in immune system cells can comprise, e.g., expressing a nucleic acid construct in immune system cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said

5

10

15

20

25

promoter sequence is selected from Table 6. In addition to the cell lines mentioned below, the construct can be expressed in primary cells, such as thymocytes, bone marrow cells, stem cells, lymphoid progenitor cells, myeloid progenitor cells, monocytes, B-cells, antigen presenting cells, macrophages, and cell lines derived therefrom.

5

10

15

20

Kidney Selective Genes

The present invention relates to genes and polypeptides which are selectively expressed in kidney tissues: TMD0049 (XM 057351), TMD0190 (XM 087157), TMD0242 (XM 088369), TMD0335 (XM 089960), TMD0371, TMD0374, TMD0469 (XM 038736), TMD0719 (XM 059548), TMD0731 (XM 059703), TMD0785 (XM 060310), TMD0841 (XM_060623), TMD1114 (NM 019841), and/or TMD 1148 (XM 087108). These genes and polypeptides are expressed predominantly in kidney tissues, making them, and the polypeptides they encode, useful as selective markers for kidney tissue and function, as well as diagnostic, prognostic, therapeutic, and research tools for any conditions, diseases, disorders, or applications associated with the kidney and the cells in which they are expressed. TMD0049 (XM 057351), TMD0190 (XM 087157), TMD0242 (XM 088369), TMD0335 (XM 089960), TMD0371, TMD0374, TMD0469 (XM 038736), TMD0719 (XM 059548), TMD0731 (XM 059703), TMD0785 (XM 060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108) includes both human and mammalian homologs of it. SEQ ID NOS 78-103 represent particular alleles, but the present invention relates to other alleles, including naturally-occurring polymorphisms (i.e., a polymorphism in the nucleotide sequence which is identified in populations of mammals) and homologs thereof. More information on these genes is summarized in Tables 8-11.

25

30

In view of their selectivity and display on the cell surface, the polypeptides and polynucleotides of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to the cells (e.g., juxtaglomerular cells which secrete renin, peritubular cells, endothelial cells, e.g., of the cortex and outer medulla, mesangial cells which secrete inflammatory mediators including NO and products of cyclooxygenase, visceral epithelial cells, parietal epithelial cells, podocytes, early proximal tubule cells which secrete, e.g., angiotensin converting enzyme and neutral endopeptidase, late distal tubule

-17-

cells that produce, e.g., prolyl endopeptidase, serine endopeptidase, carboxypeptidase, and neutral endopeptidase, renomedullary interstitial cells, etc) in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to breast cancer. They can also be used to detect metastatic cells, in biopsies, to identify kidney, etc. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly as indicated in Table 9. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

The present invention relates to methods of detecting kidney cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below,

5

10

15

20

25

such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 104, 105, 107, 108, 111, 112, 115, 116, 119, 120, 122, 123, 126, 127, 131, 132, 135, 136, 138, 139, 142, 143, 145, 146, 149, 150, and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting a kidney cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface as indicated in Table 9.

As indicated above, binding partners can be used to deliver agents specifically to the kidney, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to a kidney cell can comprise, e.g., contacting a kidney cell with an agent coupled to binding partner specific for TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the kidney can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target),

5

10

15

20

25

-19-

present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers. Any cell expressing a polypeptide coded for by TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108) can be targeted, including, e.g., juxtaglomerular, peritubular, endothelial, mesangial, visceral epithelial, parietal epithelial, podocytes, early proximal tubule, late distal tubule, renomedullary interstitial, etc.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose.

A kidney cell (see above for examples of kidney cell types) can also be modulated in accordance with the present invention, e.g., by methods of modulating a kidney cell, comprising, e.g., contacting said cell with an agent effective to modulate TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), or the biological activity of a polypeptide encoded thereby, or a mammalian homolog thereof, whereby said kidney cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

5

10

15

20

25

-20-

An activity or function of the kidney cell can be modulated, including, e.g., glomerular filtration rate, filtration pressure, renal autoregulation (including via myogenic mechanism and tubuloglomerular feedback mechanism), tubular reabsorption, tubular secretion, and renal clearance. In addition, the transcription, translation, synthesis, degradation, expression, etc., of any secretory or polypeptide produced by a kidney cell can be modulated, including, but not limited to, renin-angiotensin activity, production and secretion of prostaglandins, nitric oxide, kallikrein, adenosine, endothelin, erythropoietin, and other hormones, enzymes, and other secretory and intracellular factors. The response of a kidney cell to stimuli can also be modulated, including, but not limited to, ligands to TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), oxygen levels, blood pressure, etc.

The present invention also relates to polypeptide detection methods for assessing kidney function, e.g., methods of assessing kidney function, comprising, detecting a polypeptide coded for by TMD0049 (XM 057351), TMD0190 (XM_087157), TMD0242 (XM 088369), TMD0335 (XM 089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM 059548), TMD0731 (XM 059703), TMD0785 (XM 060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108), fragments thereof, polymorphisms thereof, in a body fluid, whereby the level of said polypeptide in said fluid is a measure of kidney function. Kidney function tests are usually performed to determine whether the kidney is functioning normally as a way of diagnosing kidney disease. Various tests are commonly used, including, e.g., BUN (blood urea nitrogen), serum creatinine, estimated GFR, ability to concentrate urine, BUN/creatine ratio, urine sodium and other electrolytes, urine NAG (N-acetyl-beta-glucosaminidase, adenosine deaminase, urinary alkaline phosphatase, serum and urine beta-2-microglobulin, serum uric acid, isotope scans, Doppler sonogram, positron emission tomography, specific gravity of urine, microalbumin, total protein, etc. Detection of TMD0049 (XM 057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM 038736), TMD0719 (XM 059548), TMD0731 (XM_059703), TMD0785 (XM 060310), TMD0841 (XM 060623), TMD1114 (NM 019841), and/or TMD 1148

5

10

15

20

25

-21-

(XM_087108) provides an additional assessment tool, especially in diseases such as chromic renal failure, urinary tract infections, kidney stones, nephrotic syndrome, nephritic syndrome, kidney disease due to diabetes or high blood pressure, etc., As with the other tests, elevated levels of said polypeptide in blood, or other fluids, can indicate impaired kidney function. Values can be determined routinely, as they are for other kidney function markers, such as those mentioned above. Detecting can be performed routinely (see below), e.g., using an antibody which is specific for said polypeptide, by RIA, ELISA, or Western blot, etc.

Promoter sequences obtained from genes of the present invention can be utilized to selectively express heterologous genes in kidney cells. Methods of expressing a heterologous polynucleotide in kidney cells can comprise, e.g., expressing a nucleic acid construct in kidney cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected SEQ ID NOS 106, 109, 110, 113, 114, 117, 118, 121, 124, 125, 128-130, 133, 134, 137, 140, 141, 144, 147, 148, and 151. In addition to the cell lines mentioned below, the construct can be expressed in primary cells or in established cell lines.

Kidney

5

10

15

20

25

30

The kidney maintains the constancy of fluids in an organism's internal environment, and is therefore of great importance in maintaining health and vitality. Each day, the kidney filters the blood, removing and concentrating toxins, metabolic wastes, and excess ions, allowing them to be excreted by the body in the form of urine. The excretory function of the kidney is performed by over one million blood units called nephrons, each a miniature blood filtering and processing unit. A nephron consists of a glomerulus, a tuft of capillaries, and a renal tubule. In addition to their excretory function, kidneys produce a number of different hormones, enzymes, and other secreted molecules, including the enzyme renin and the hormone erythropoietin. The kidney also is responsible for metabolizing vitamin D into its active form, calcitriol. For a full description of the kidney's function and structure, see, e.g., *Human Anatomy and Physiology*, Marieb, E.N., 3rd Edition, Benjamin/Cummings Publishing Company, Inc., 1995, pp 896-923.

The glomerulus is a high pressure capillary bed which filters out most substances smaller than large plasma proteins across the fenestrated glomerular epithelium, the

10

15

20

25

30

intervening basement membrane, and the podocyte-containing visceral membrane of the glomerulus capsule. The external layer of the glomerulus is called the parietal layer, consisting predominally of a squamous epithelium. This layer is structural. Underneath it, is the visceral layer which consists of the modified branching epithelial cells called podocytes. These sit on top of the fenestratrated glomerular endothelium. The glomerulus is connected to the renal tubule, a highly differentiated and long tube, having three major elements: the proximal convoluted tubule, the loop of Henel, and the distal convoluted tubule. Different regions of the tubule have different functions in absorption and secretion.

Renal cells produce a variety of different hormones and chemicals, including, prostaglandins, nitric oxide, kallikrein family, adenosine, endothelin family, renin, erythropoietin, aldosterone, antidiuretic hormone (vasopressin), natriuretic hormones, etc. Renin is involved in modulating blood pressure. It cleaves angiotensinogen, a plasma peptide, splitting off a fragment containing 10 amino acids called angiotensin I. Angiotensin I is cleaved by a peptidase secreted by blood vessels called angiotensin converting enzyme (ACE), producing angiotensin II, which contains 8 amino acids. Angiotensin II has many direct effects on blood pressure. Erythropoietin stimulates red blood cell production in the bone marrow.

TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108) can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of the kidney. These include, but are not limited to, diseases that affect the four basic morphologic components, glomeruli, tubules, interstitium, and blood vessels. Diseases include, e.g., acute nephritic syndrome, nephritic syndrome, renal failure, urinary tract infections, renal stones, cystic diseases of the kidney, e.g., cystic renal dysplasia, polycystic disease (autosomal dominant and recessive types), medullary cystic disease, acquired cystic disease, renal cysts, parenchymal cysts, perihilar renal cysts (pyelocalyceal cysts, hilar lymphangitic cysts), glomerular diseases, diseases of tubules, tubulointerstitial diseases, tumors of the kidney, such as benign tumors (cortical adenoma, renal fibroma, renomedullary interstitial cell tumor), malignant tumors (renal cell carcinoma, hypernephroma,

WO 03/089583

-23-

PCT/US03/11497

adenocarcinoma of kidney, Wilms' tumor, nephroblastoma, urothelial carcinoma), renal coloboma, nephorblastoma, clear cell sarcoma of kidney (CCSK), rhabdoid tumor of kidney (RTK), von Hippel-Lindau disease, oncocytoid renal cell carcinoma (RCC), renal leiomyoblastoma, etc. TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108) can also be used for staging and classifying conditions and diseases of the present invention, alone, or in combination with conventional staging and classification schemes.

10

15

5

Pancreatic Gene Complex

The present invention relates to a cluster of olfactory GPCR (G-protein coupled) receptor genes located at chromosomal band 11q24. These genes are expressed predominantly in pancreatic tissue, establishing this region of chromosome 11 as a unique gene complex involved in pancreatic function. See, Table 12. Because of their exquisite selectivity for pancreatic tissues, the pancreatic gene complex ("PGC"), and the genes which comprise it, are useful to assess pancreas tissue and function for diagnostic, prognostic, therapeutic, and research purposes.

The spatial organization of the pancreatic gene complex ("PGC") is illustrated in Fig.

7. It spans several hundred kilobases of chromosome 11, e.g., from about LOC160205 to LOC119954, from about LOC119944-LOC119954, and any part thereof. Within this region, is a cluster of genes coding for polypeptides which share sequence identity with the olfactory GPCR family. These include, but are not limited to, TMD0986, XM_061780 (TMD0987), XM_061781 (TMD0353), XM_061784 (TMD0989), XM_061785 (TMD058). Fig. 8

25 illustrates the relationship between the lengths of the different coding sequences. As shown in the figure, XM_061784 is shorter at its C-terminus than the other family members.

As members of the GPCR family, the PGC genes all share a degree of amino acid sequence identity and similarity. See, Table 14 for values (% sequence identity is the first place; % sequence similarity is in parenthesis in the second place; calculations were performed using the publicly-available BLASTP pair-wise alignment program). TMD0986, XM_061780, XM_061781, and XM_061785 each share about 40% sequence identity.

-24-

BLAST searching of publicly available sequences indicates that these polypeptides share less amino acid sequence identity with each other than they do with other olfactory GPCR homologs located elsewhere in the genome. Significantly higher amino acid sequence identity – 81% – is observed between the adjacent genes XM_061784 and XM_061785. These genes appear to be part of a sub-cluster within PGC that share high polypeptide similarity between them.

The phrase "a gene of Table 12" which is used throughout the description include the specific sequences for the listed XM numbers as well as other human alleles, and mammalian homologs, such as murine homologs. For example, Table 14 lists several of the mouse homologs that are included in the present invention. While SEQ ID NOS. 152, 153, 162, 163, 167, 168, 171, 172, 175, and 176 may represent particular alleles, the present invention relates to other alleles, as well, including naturally-occurring polymorphisms (i.e., a polymorphism in a nucleotide sequence which is identified in populations of mammals).

TMD0986 (SEQ ID NO 152 and 153) is a full-length sequence of the previously identified XM_061779. It contains an additional 117 amino acids not present in XM_061779. The present invention relates to nucleic acids comprising or consisting essentially of this sequence in its entirety (e.g., amino acids 1-314), comprising or consisting essentially of nucleic acids coding for amino acids 1-117, and comprising or consisting essentially of fragments of nucleic acids coding for amino acids 1-117. Polypeptides encoded by these nucleic acids are also claimed, including polypeptide fragments of 1-117, such as 1-23, 79-97, 164-198, 261-274, and other extracellularly exposed peptides. In addition, the present invention relates to binding partners, such as antibodies, that bind to epitopes within amino acids 1-117 (e.g., SEQ ID NO 153).

25 Pancreas

5

10

15

20

30

Diabetes and other pancreatic disorders are a major health concern. Worldwide, it is estimated that 5-10% of the population suffers from some form of diabetes. Pancreatic cancer is the fifth leading cause of cancer-related mortality. In 2002, it was estimated that about 30,000 Americans would be diagnosed with pancreatic cancer, and 90% would die within 12 months. Despite the prevalence of pancreatic disease, the genetics and physiology of normal pancreatic function and pancreatic disease is still poorly understood.

-25-

The pancreas is a mixed gland comprised of exocrine and endocrine tissues. The exocrine portion comprises about 80-85% of the organ. It is divided into lobes by connective tissue septa, and each lobe is divided into several lobules. These lobules are composed of grape-like clusters of secretory cells that form sacs known as acini. An acinus is a functional unit of the pancreatic exocrine gland. All acini drain into interlobular ducts which merge to form the main pancreatic duct. It, in turn, joins together with the bile duct from the liver to form the common bile duct that empties into the duodenum. Pancreatic acinar cells make up more than 80% of the total volume of the pancreas and function in the secretion of the various enzymes that assist digestion in the gastrointestinal tract. Scattered among the acinar cells are approximately a million pancreatic islets ("islets of Langerhans") that secrete the pancreatic endocrine hormones. These dispersed islets comprise approximately 2% of the total volume of the pancreas.

The basic function of the pancreatic endocrine cells is to secrete certain hormones that participate in the metabolism of proteins, carbohydrates, and fats. The hormones secreted by the islets include, e.g., insulin, glucagon, somatostatin, pancreatic polypeptide, amylin, adrenomedullin, gastrin, secretin, and peptide-YY. See, also, Shimizu et al., *Endocrin.*, 139:389-396, 1998. The islets contain about four major and two minor cell types. The major cell types are alpha (glucagon producing), beta (insulin and amylin producing), delta (somatostatin producing which suppresses both insulin and glucagon release), and F (pancreatic polypeptide and adrenomedullin producing) cells. The minor cell types are D1 (produce vasoactive intestinal peptide or VIP) and enterochromaffin (produce serotonin) cells. The cells can be distinguished, e.g., by their morphology, hormonal content, and polynucleotide expression patterns.

The ability of the pancreas to respond to a wide variety of metabolic signals is conferred by an expression profile comprising a rich assortment of receptor proteins. G-protein coupled receptors have been previously identified in the pancreas, including, e.g., receptors for glucagon, secretin, CCK (e.g., Roettger et al., *J. Cell Biol.*, 130:579-590, 1995), purines (e.g., P2 purinoreceptors), gastrin, KiSS-1 peptides (e.g., Kotani et al., *J. Biol. Chem.*, 276:34631-6, 2001), adrenomedullin (Martinez et al., *Endocrin.*, 141:406, 2000), and interleukins. G-protein subunits have also been localized to the pancreas, including G-proteins which were previously associated with the olfactory epithelium. See, e.g., Zigman et

5

10

15

20

25

-26-

al., *Endocrin.*, 133:2508-2514, 1993. In addition, pancreatic cells express neurotropin, neurotensin, and interleukin receptors.

As mentioned, the pancreas is sensitive to a variety of metabolic, soluble and hormonal signals involved in regulating blood sugar, modulating synthesis and release of pancreatic digestive enzymes, and other physiologically important processes involved in pancreas function. In analogy to the ability of olfactory receptors to detect odors and pheromones in the environment, the pancreatic GPCRs of the present invention can be used to "sniff" out and respond to various ligands in the blood which pass through the pancreas, including peptides, metabolites, and other biologically-active molecules. Biological activities include, but are not limited to, e.g., regulation of blood sugar, modulation of all aspects of the various secreted polypeptides (hormones, enzymes, etc.) produced by the pancreas, ligand-binding, exocytosis, amylase (and any of the other 20 or so digestive enzymes produced by the pancreas) secretion, autocrine responses, apoptosis (e.g., in the survival of beta-islet cells), zymogen granule processing, G-protein coupling activity, etc.

The polynucleotides, polypeptides, and ligands thereto, of the present invention can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of pancreas. These include, but are not limited to, e.g., disorders associated with loss or mutation to 11q24, such as Jacobsen syndrome (OMIM #147791), cystic fibrosis, acute and chronic pancreatitis, pancreatic abscess, pancreatic pseudocyst, nonalcoholic pancreatitis, alcoholic pancreatitis, classic acute hemorrhagic pancreatitis, chronic calcifying pancreatitis, familial hereditary pancreatitis, carcinomas of the pancreas, primary (idiopathic) diabetes (e.g., Type I (insulin dependent diabetes mellitus, IDDM) [insulin deficiency, beta cell depletion], Type II (non-insulin dependent diabetes mellitus, NIDDM) [insulin resistance, relative insulin deficiency, mild beta cell depletion]), nonobese NIDDM, obese NIDDM, maturity-onset diabetes of the young (MODY), islet cell tumors, diffuse hyperplasia of the islets of Langerhans, benign adenomas, malignant islet tumors, hyperfunction of the islets of Langerhans, hyperinsulinism and hypoglycemia, Zollinger-Ellison syndrome, beta cell tumors (insulinoma), alpha cell tumors (glucagonoma), delta cell tumors (somatostatinoma), vipoma (diarrheogenic islet cell tumor), pancreatic cancers, pancreatic carcinoid tumors, multihormonal tumors, multiple endocrine neoplasia (MEN), MEN I (Wermer syndrome), MEN II (Sipple syndrome), MEN III or IIb, pancreatic endocrine

5

10

15

20

25

tumors, etc.

5

10

15

20

25

30

In view of its selectivity and display on the cell surface, the olfactory GPCR family members of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to the cells (e.g., pancreatic progenitor, exocrine, endocrine, acinar, islet, alpha, beta, delta, F, D1, enterochromaffin, etc.) in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to breast cancer. They can also be used to detect metastatic cells, in biopsies to identify bone marrow, lymphocytes, etc. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly as indicated in Table 14. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

The present invention relates to methods of detecting pancreas cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for a gene of Table 12, or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and

-28-

technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 154, 155, 164, 165, 169, 170, 173, 174, 177, and 178, and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting a pancreas cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by a polypeptide of Table 12, or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface as indicated in Table 14.

As indicated above, binding partners can be used to deliver agents specifically to the pancreas, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to a pancreas cell can comprise, e.g., contacting a pancreas cell with an agent coupled to a binding partner specific for a polypeptide coding for a gene of Table 12, whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the pancreas can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers. Any cell expressing a polypeptide coded for by a gene of Table 12 can be targeted, including, e.g., pancreatic progenitor, exocrine, endocrine, secretory, acinar, islet, alpha, beta, delta, F, D1, enterochromaffin, etc.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body.

5

10

15

20

25

-29-

Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose. See, Bruehlmeier et al., *Nucl. Med. Biol.*, 29:321-327, 2002, for imaging pancreas using labeled receptor ligands. Antibodies and other ligands to receptors of the present invention can be used analogously.

A pancreas cell (see above for examples of pancreas cell types) can also be modulated in accordance with the present invention, e.g., by methods of modulating a pancreas cell, comprising, e.g., contacting said cell with an agent effective to modulate a gene of Table 12, or the biological activity of a polypeptide encoded thereby (e.g., SEQ ID NO 153, 163, 168, 172, or 176), or a mammalian homolog thereof, whereby said pancreas cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

An activity or function of the pancreas cell can be modulated, including, e.g., regulation of blood sugar, modulation of all aspects of the various secreted polypeptides (hormones, enzymes, etc.) produced by the pancreas, ligand-binding, exocytosis, amylase (and any of the other 20 or so digestive enzymes produced by the pancreas) secretion, autocrine responses, apoptosis (e.g., in the survival of beta-islet cells), etc.

The present invention also relates to polypeptide detection methods for assessing pancreas function, e.g., methods of assessing pancreas function, comprising, detecting a polypeptide coded for by a gene of Table 12, fragments thereof, polymorphisms thereof, in a body fluid, whereby the level of said polypeptide in said fluid is a measure of pancreas function. Pancreas function tests are usually performed to determine whether the pancreas is functioning normally as a way of diagnosing pancreas disease. Various tests are commonly used, including, e.g., assays for the presence of pancreatic enzymes in body fluids (e.g.,

5

10

15

20

25

10

15

20

25

30

amylase, serum lipase, serum trypsin-like immuoreactivity), studies of pancreatic structure (e.g., using x-ray, sonography, CT-scan, angiography, endoscopic retrograde cholangiopancreatography), and tests for pancreatic function (e.g., secretin-pancreozymin (CCK) tst, Lundh meal test, Bz-Ty-PABA test, chymotrypsin in feces, etc). Detection of a polypeptide coded for by a gene of Table 12 provides an additional assessment tool, especially in diseases such as pancreatitis and pancreatic cancer where pancreatic markers can appear in the blood, stool, urine, and other body fluids. As with the other tests, elevated levels of said polypeptide in blood, or other fluids, can indicate impaired pancreas function. Values can be determined routinely, as they are for other markers, such as those mentioned above. Detecting can be performed routinely (see below), e.g., using an antibody which is specific for said polypeptide, by RIA, ELISA, or Western blot, etc., in analogy to the tests for pancreatic enzymes in body fluids.

Promoter sequences obtained from GPCR genes of the present invention can be utilized to selectively express heterologous genes in pancreas cells. Methods of expressing a heterologous polynucleotide in pancreas cells can comprise, e.g., expressing a nucleic acid construct in pancreas cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected SEQ ID NOS 156-161, 166, 179, or 180. In addition to the cell lines mentioned below, the construct can be expressed in primary cells or in established cell lines.

The genes and polypeptides of Table 12 can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of the pancreas as mentioned above. The present invention relates to methods of identifying a pancreatic disease or pancreatic disease-susceptibility, comprising, e.g., determining the association of a pancreatic disease or pancreatic disease-susceptibility with a nucleotide sequence present within the pancreatic gene complex. An association between a pancreas disease or disease-susceptibility and nucleotide sequence includes, e.g., establishing (or finding) a correlation (or relationship) between a DNA marker (e.g., gene, VNTR, polymorphism, EST, etc.) and a particular disease state. Once a relationship is identified, the DNA marker can be utilized in diagnostic tests and as a drug target.

Any region of the pancreatic gene complex can be used as a source of the DNA marker (e.g., a nucleotide sequence present with PGC), including, e.g., TMD0986,

XM_061780 (TMD0987), XM_061781 (TMD0353), XM_061784 (TMD0989), XM_061785 (TMD058), and any part thereof, introns, intergenic regions, any DNA from about 29160-29310 kb of 11q24, NT 009215, etc.

-31-

Human linkage maps can be constructed to establish a relationship between a region within 11q24 and a pancreatic disease or condition. Typically, polymorphic molecular markers (e.g., STRP's, SNP's, RFLP's, VNTR's) are identified within the region, linkage and map distance between the markers is then established, and then linkage is established between phenotype and the various individual molecular markers. Maps can be produced individual family, selected populations, patient populations, etc. In general, these methods involve identifying a marker associated with the disease (e.g., identifying a polymorphism in a family which is linked to the disease) and then analyzing the surrounding DNA to identity the gene responsible for the phenotype.

Retina Selective Gene

5

10

15

20

25

30

The present invention relates to NM 013941 (GPCR181 or OR10C1), a multiple transmembrane spanning polypeptide which shares sequence identity with the olfactory Gprotein coupled receptor (GPCR) family. Like other GPCR, NM 013941 has seven transmembrane domains, at about amino acid positions 20-42, 54-76, 91-113, 134-156, 190-212, 233-255, and 265-287, of SEQ ID NO 182. It is located at about chromosomal band 6p21.31-22.2. There are several other GPCRs located nearby (e.g., OR2B3, AL022727; OR2J3, AL022727). NM 013941 is highly expressed in brain tissue, at lower levels in heart, pituitary, and skin, and at minimally detectable levels in colon, small intestine, kidney, lymphocytes, and mammary gland. In the neuronal tissue, it was selectively expressed in the retina, but was not detected in any other brain tissue regions. The selective expression of NM 013941 in the retina makes it useful as a marker for retinal tissue, e.g., in stem cell cultures and biopsy samples, as well as a diagnostic, prognostic, therapeutic, and research tool for any conditions, diseases, disorders, or applications associated with the retina and the cells in which it is expressed. NM 013941 includes both human and mammalian homologs of it (e.g., mouse XM 111729 which is similar to olfactory receptor MOR263-6). SEQ ID NOS. 181 and 182 represent a particular allele of NM 013941; the present invention relates to other alleles, as well, including naturally-occurring polymorphisms (i.e., a polymorphism in the nucleotide sequence which is identified in populations of mammals).

The chromosomal region within which NM_013941 is located comprises a number of genes involved in retinal function. These include, e.g., retinal cone dystrophy (OMIM 602093) which appears to be a result of mutation in guanylate cyclase activator-1A (e.g., Payne et al., *Human Molec. Genet.*, 7:273-277, 1998), retinal degeneration slow (OMIM 179605) which appears to be a defect in specific retinal protein homologous to rod outer segment protein-1, retinitis pigmentosa-7, retinitis pigmentosa-14 (OMIM 600132) which is associated with a mutation in the tubby-like protein TULP1 (e.g., Banerjee et al., *Nature Genet.*, 18:177-179, 1998; Hagstrom et al., *Nature Genet.*, 18:174-176, 1998), and others. Thus, this region appears to be important in eye function.

In view of its selectivity and display on the cell surface, the olfactory GPCR family members of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to retinal cells. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat retinal carcinomas (e.g., retinoblastoma) in analogy to how c-erbB-2 antibodies are used to breast cancer. See, e.g., Hayashi et al., *Invest. Ophthalmol. Vis. Sci.*, 40:265-72, 1999 for an example treating retinoblastoma using HSV-TK. Transfer of the gene into the retinal cells can be achieved by incorporating the gene into liposomes which have been made cell-selective by incorporating a NM_013941 specific antibody into its bilayer. See, also, Wu and Wu, *J. Biol. Chem.*, 262: 4429-4432, 1987.

The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule

5

10

15

20

25

PCT/US03/11497

comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

-33-

The present invention relates to methods of detecting retinal cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for NM_013941 (e.g., SEQ ID NOS 181), or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 183 and 184, and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting a retinal cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by NM_013941 (e.g., SEQ ID NO 182), or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface.

As indicated above, binding partners can be used to deliver agents specifically to the retina, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to a retinal cell can comprise, e.g., contacting a retinal cell with an agent coupled to binding partner specific for NM_013941 (SEQ ID NO 182), whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the retinal can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent

5

10

15

20

25

-34-

is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers. Any cell expressing a polypeptide coded for by NM_013941 can be targeted, including, e.g., pigmented epithelial cells, photoreceptor cells, cones, rods, bipolar cells, ganglion cells, etc.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose.

A retinal cell (see above for examples of retinal cell types) can also be modulated in accordance with the present invention, e.g., by methods of modulating a retinal cell, comprising, e.g., contacting said cell with an agent effective to modulate NM_013941, or the biological activity of a polypeptide encoded thereby (e.g., SEQ ID NO 182), or a mammalian homolog thereof, whereby said retinal cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

Any activity or function of the retinal cell can be modulated, including, e.g., light reception, phototransduction, excitation of rods, excitation of cones, metabolism of vitamin A, retinal, rhodopsin, and other functional molecules, cGMP binding and hydrolysis, sodium channel flux, membrane potential, phosphodiesterase activity, G-protein activity and coupling, vitamin A processing, sodium pump activity, calcium flux, etc. The response of a retinal cell to stimuli can also be modulated, including, but not limited to, ligands to

5

10

15

20

25

-35-

NM_013941, light, ion levels, second messenger levels, etc.

Promoter sequences can be utilized to selectively express heterologous genes in retinal cells. Methods of expressing a heterologous polynucleotide in retinal cells can comprise, e.g., expressing a nucleic acid construct in retinal cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is obtained from NM_01394, e.g., on genomic NT_007592. In addition to the cell lines mentioned below, the construct can be expressed in primary cells or in established cell lines.

10 Retina

5

15

20

25

30

The retina is a two-layered structure located on the back of the eye. It is the primary organ responsible for vision. The outer pigmented layer is comprised of pigmented epithelial cells that absorb light, preventing it from scattering in the eye, and store vitamin A needed by the photoreceptor cells. The inner neural layer is comprised of three main cell types: photoreceptor cells, bipolar cells, and ganglion cells. The local currents generated by a light stimulus spreads from the photoreceptor cells to the bipolar cells, and then on to the innermost ganglion cells. The optic disc is the exit site of the retinal ganglion axons which then bundle into the optic nerve

Photoreceptors consist of rods and cones which are the photosensitive cells of the retina. Each rod and cone elaborates a specialized cilium, called the outer segment, that contains the phototransduction machinery. The rods contain a specific light-absorbing visual pigment, rhodopsin. In humans, there are three classes of cones, each characterized by the expression of distinct visual pigments: the blue cone, green cone and red cone pigments. Each type of visual pigment protein is tuned to absorb light maximally at different wavelengths. The rod rhodopsin mediates scotopic vision (in dim light), whereas the cone pigments are responsible for photopic vision (in bright light). The red, blue and green pigments also form the basis of color vision.

NM_013941 can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of the retinal. These include, but are not limited to, diseases that affect the basic morphologic components as mentioned above, e.g., the outer and inner cell layers, and the optic nerve the retina. Diseases include, e.g.,

retinal degeneration, retinal degenerations such as retinitis pigmentosa, Bardet-Biedl syndrome, Bassen-Kornzweig syndrome (abetalipoproteinemia), Best disease (vitelliform dystrophy), choroidemia, gyrate atrophy, congenital amaurosis, Refsum syndrome, Stargardt disease, Usher syndrome, macular degeneration (dry and wet forms), diabetic retinopathy, peripheral vitreoretinopathies, photic retinopathies, surgery-induced retinopathies, viral retinopathies (such as HIV retinopathy related to AIDS), ischemic retinopathies, retinal detachment, traumatic retinopathy, optic neuropathy, optic neuritis, ischemic optic neuropathy, Leber optic neuropathy, diseases of Bruch's membrane, glaucoma, cancer, retinoblastoma, cancer- associated retinopathy syndrome (CAR syndrome), melanoma-associated retinopathy (MAR), etc. NM_013941 can also be used for staging and classifying conditions and diseases of the present invention, alone, or in combination with conventional staging and classification schemes.

Spleen Gene Cluster

5

10

15

20

25

30

The present invention relates to a cluster of transmembrane and GPCR-type receptor genes located at chromosomal band 11q12.2. The genes of the present invention are expressed predominantly in the spleen (e.g., Fig. 10, lane 19) (hence, "spleen gene" cluster), as well as other tissues of the immune and reticuloendothelial system (RES), establishing this region of the chromosome as a unique gene complex involved in spleen, lymphoid, and/or reticuloendothelial function. TMD1030 and TMD0621 are highly expressed in spleen tissue, with insignificant levels in other tissues. In addition to spleen. TMD1029 and TMD1029 show significant expression in the liver and lymphocytes, as well. Because of their selectivity for spleen, lymphoid, and/or reticuloendothelial tissues, the gene complex, and the chromosomal region which comprises it, are useful to assess spleen, lymphoid, and/or reticuloendothelial tissue function and for diagnostic, prognostic, therapeutic, and research purposes. Information on the genes is summarized in Tables 15-19.

The spatial organization of the gene complex is illustrated in Fig. 11. The complex spans about at least 100 kb, from about EST markers G62658, SHGC-82134, etc. (located at the end closest to the centromere and TMD1030) to SHGC-154002, SHGC-9433, etc. (located at the end furthest from the centromere and TMD0621). All the genes have the same orientation of transcription. TMD1799 (XM 166849) (SEQ ID NO 193-194), located at the

-37-

upper region, shows very high expression in lymphocytes, but only marginal expression in spleen, indicating that expression in lymphocytes may predominate at the boundaries of the gene complex. In the lower region, TMD1027 (XM_166856) (SEQ ID NO 195-196), spleen expression virtually disappears, while lymph node expression becomes very high. The present invention includes this entire region, and any parts thereof. For instance, the present invention includes any DNA fragments within it which confer the observed tissue specificities described herein.

The gene complex is involved in spleen, immune, and RES functions. The spleen is located in the left upper region of the abdomen. In the adult, it weights about 90-180 grams, and is about 15 by 7.5 cm in size. The spleen is anatomically and functionally compartmentalized into two distinct regions, the red and white pulp. The red pulp comprises blood vessels interwoven with connective tissue ("pulp cords") that is lined with reticuloendothelial cells. It possesses a blood filtering function, removing opsonized cells and trapping abnormal red blood cells. It also is a storage reservoir for platelets and other blood cells. In the fetus, the red pulp has a hematopoietic function. Inside the red pulp, is lymphoid tissue know as the white pulp. Antibodies are made inside the white pulp. Similar to other lymphatic tissues, B- and T-cell's mature inside the white pulp, where they are involved in antigen presentation and lymphocyte maturation. The white pulp is clustered around the periarteriolar lymphoid sheath, and is comprised of follicles and marginal zone.

Naive B-cells are located in the primary follicle, memory cells, macrophages, and dendritic cells in the secondary follicle, and macrophages and B-cells in the marginal zone. The integrins LFA-1 and alpha4-beta1 are involved in localization of the B-cells to the marginal zone of the white pulp (Lu and Cyster, Science, 297:409, 2002).

The reticuloendothelial system (RES) is a multi-organ phagocytic system involved in removing particulates from the blood. It is comprised of the spleen and liver. It has the ability to sequester inert particles and dyes. Cells of the RES system include, macrophages, liver Kuppfer cells, endothelial cells lining the sinusoids of the liver, spleen, and bone marrow, and reticular cells of lymphatic and bone marrow tissues.

The polynucleotides, polypeptides, and ligands thereto, of the present invention can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of spleen, lymphoid, and/or reticuloendothelial tissues. These include, but are not limited to, splenomegaly, hypersplenism, hemolytic anemis, hereditary

5

10

15

20

25

-38-

spherocytosis, hereditary eliptocytosis, thalassemia minor and major, autoimmune hemolytic anemia, thrombocytopenia, idiopathic thrombocytopenic purpura, immunologic thrombocytopenia associated with chronic lymphocytic leukemia or systemic lupus erythematosis, TTP, leukemia, lymphoma, primary and metastatic tumors, splenic cysts, infection, inflammatory diseases, anemias, blood cancers, etc. See, Table 19 for other examples.

In view of their selectivity and display on the cell surface, the genes of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to the cells (e.g., reticuloendothelial cells, macrophages, Kupffer cells, monocytes, B-lymphocytes, T-lymphocytes, etc) in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to treat breast cancer. They can also be used to detect metastatic cells in biopsies. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly. See, Table 16. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types. TMD1030 and TMD0621 are predominantly and selectively expressed in spleen tissue.

5

10

15

20

25

The expression patterns of the selectively expressed polynucleotides disclosed herein can be described as a "fingerprint" in that they are a distinctive pattern displayed by a tissue. Just as with a fingerprint, an expression pattern can be used as a unique identifier to characterize the status of a tissue sample. The list of expressed sequences disclosed herein provides an example of such a tissue expression profile. It can be used as a point of reference to compare and characterize samples. Tissue fingerprints can be used in many ways, e.g., to classify an unknown tissue, to determine the origin of metastatic cells, to assess the physiological status of a tissue, to determine the effect of a particular treatment regime on a tissue, to evaluate the toxicity of a compound on a tissue of interest, etc.

For example, the tissue-selective polynucleotides disclosed herein represent the configuration of genes expressed by a normal tissue. To determine the effect of a toxin on a tissue, a sample of tissue can be obtained prior to toxin exposure ("control") and then at one or more time points after toxin exposure ("experimental"). An array of tissue-selective probes can be used to assess the expression patterns for both the control and experimental samples. As discussed in more detail below, any suitable method can be used. For instance, a DNA microarray can be prepared having a set of tissue-selective genes arranged on to a small surface area in fixed and addressable positions. RNA isolated from samples can be labeled using reverse transcriptase and radioactive nucleotides, hybridized to the array, and then expression levels determined using a detection system. Several kinds of information can be extracted: presence or absence of expression, and the corresponding expression levels. The normal tissue would be expected to express substantially all the genes represented by the tissue-selective probes. The various experimental conditions can be compared to it to determine whether a gene is expressed, and how its levels match up to the normal control.

While the expression profile of the complete gene set represented by the sequences disclosed here may be most informative, a fingerprint containing expression information from less than the full collection can be useful, as well. In the same way that an incomplete fingerprint may contain enough of the pattern of whorls, arches, loops, and ridges, to identify the individual, a cell expression fingerprint containing less than the full complement may be adequate to provide useful and unique identifying and other information about the sample. Moreover, because of heterogeneity of the population, as well differences in the particular physiological state of the tissue, a tissue's "normal" expression profile is expected to differ

5

10

15

20

25

-40-

between samples, albeit in ways that do not change the overall expression pattern. As a result of these individual differences, each gene although expressed selectively in spleen, may not on its own 100% of the time be adequately enough expressed to distinguish said tissue. Thus, the genes can be used in any of the methods and processes mentioned above and below as a group, or one at a time.

The present invention relates to methods of detecting spleen, lymphoid, and/or reticuloendothelial cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include SEQ ID NOS 197-204 listed in Table 17, and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g., monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting a spleen, lymphoid, and/or reticuloendothelial cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by a polypeptide of the present invention, or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface. Detection can be useful for assessing spleen integrity, e.g., when it is suspected that the spleen is damaged and undergoing deterioration. The appearance of polypeptides of the present invention in body fluids, such as blood, can indicate spleen damage, including neoplastic and/or apoptotic changes.

As indicated above, binding partners can be used to deliver agents specifically to the spleen, lymphoid, and/or reticuloendothelial tissues, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to a spleen, lymphoid, and/or

5

10

15

20

25

WO 03/089583 PCT/US03/11497 -41-

reticuloendothelial cell can comprise, e.g., contacting a spleen, lymphoid, and/or reticuloendothelial cell with an agent coupled to a binding partner specific for a polypeptide coding for TMD1030 (XM 166853), TMD1029 (XM 166854), TMD1028 (XM 166855), or TMD0621 (XM 166205), whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the spleen, lymphoid, and/or reticuloendothelial tissue can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parenterally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers. Any cell expressing a polypeptide coded for by TMD1030 (XM 166853), TMD1029 (XM 166854), TMD1028 (XM 166855), or TMD0621 (XM 166205) can be targeted, including, e.g., reticuloendothelial cells, macrophages, Kupffer cells, lymphocytes, B-lymphocytes, T-lymphocytes, etc.

Antibodies (alone or conjugated to active agents) can be used to ablate spleen and other tissues. For instance, in diseases where splenectomy is indicated (e.g., immune thrombocytopenic purpura, autoimmune hemolytic anemia, blood cell disorders, myeloproliferative disorders, tumors, hypersplenism, etc.), antibodies to TMD1030 and TMD0621 can be used to ablate spleen tissue, or block spleen function.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintiographic imaging. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917, 6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The

5

10

15

20

25

10

15

20

25

30

methods described therein can be used generally to associate a partner with an agent for any desired purpose. See, Bruehlmeier et al., *Nucl. Med. Biol.*, 29:321-327, 2002, for imaging using labeled receptor ligands. Antibodies and other ligands to receptors of the present invention can be used analogously.

A cell (see above for examples of spleen, lymphoid, and/or reticuloendothelial cell types) can also be modulated in accordance with the present invention, e.g., by methods of modulating a spleen, lymphoid, and/or reticuloendothelial cell, comprising, e.g., contacting said cell with an agent effective to modulate TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), or the biological activity of a polypeptide encoded thereby (e.g., SEQ ID NOS 185-192), or a mammalian homolog thereof, whereby said spleen, lymphoid, and/or reticuloendothelial cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

Any activity or function of the spleen, lymphoid, and/or reticuloendothelial tissues can be modulated, including, e.g., immune modulation (e.g., modulating antigen presentation, antibody production and secretion, humoral and cellular responses, etc.), sequestration and removal of red blood cells, clearance of microorganisms and particular antigens from blood, migration into the marginal zone or other immune and RES compartments, etc.

The present invention also relates to polypeptide detection methods for assessing spleen, lymphoid, and/or reticuloendothelial tissue function, e.g., methods of assessing spleen, lymphoid, and/or reticuloendothelial function, comprising, detecting a polypeptide coded for by TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), fragments thereof, polymorphisms thereof, in a body fluid, whereby the level of said polypeptide in said fluid is a measure of spleen, lymphoid, and/or reticuloendothelial function. spleen, lymphoid, and/or reticuloendothelial function tests are usually performed to determine whether the spleen, lymphoid, and/or reticuloendothelial tissue is functioning normally as a way of diagnosing spleen, lymphoid, and/or reticuloendothelial disease. Various tests are commonly used, including, e.g., 99Tc-colloid

liver-spleen scan, computed tomography, ultrasound scanning of left upper quandrant, MRI,

liver enzymes, etc.

-43-

Detection of a polypeptide coded for by TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), provides an additional assessment tool, especially in diseases or disorders, such as splenomegaly, hypersplenism, or ruptured spleen, where said polypeptides can appear in the blood, stool, urine, and other body fluids. As with the other tests, elevated levels of said polypeptide in blood, or other fluids, can indicate impaired spleen, lymphoid, and/or reticuloendothelial function. Values can be determined routinely, as they are for other markers, such as those mentioned above. Detecting can be performed routinely (see below), e.g., using an antibody which is specific for said polypeptide, by RIA, ELISA, or Western blot, etc., in analogy to the tests for enzymes and other proteins in body fluids.

Promoter sequences obtained from genes of the present invention can be utilized to selectively express heterologous genes in cells. Methods of expressing a heterologous polynucleotide in cells, e.g., spleen, lymphoid, and/or reticuloendothelial cells can comprise, e.g., expressing a nucleic acid construct in spleen, lymphoid, and/or reticuloendothelial cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected SEQ ID NOS 205-213. In addition to the cell lines mentioned below, the construct can be expressed in primary cells or in established cell lines.

The genes and polypeptides of the present invention can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of the spleen, lymphoid, and/or reticuloendothelial tissues mentioned above. The present invention relates to methods of identifying a genetic basis for a disease or disease-susceptibility, comprising, e.g., determining the association of a spleen, lymphoid, and/or reticuloendothelial disease or spleen, lymphoid, and/or reticuloendothelial disease-susceptibility with the gene complex of the present invention, e.g., a nucleotide sequence present in the gene complex at 11q12.2. An association between a spleen, lymphoid, and/or reticuloendothelial disease or disease-susceptibility and nucleotide sequence includes, e.g., establishing (or finding) a correlation (or relationship) between a DNA marker (e.g., gene, VNTR, polymorphism, EST, etc.) and a particular disease state. Once a relationship is identified, the DNA marker can be utilized in diagnostic tests and as a drug target.

Any region of the gene can be used as a source of the DNA marker, exons, introns,

5

10

15

20

25

-44-

intergenic regions, or any DNA from the gene cluster of the present invention at chromosomal region 11q12.2, etc.

Human linkage maps can be constructed to establish a relationship between a gene and a spleen, lymphoid, and/or reticuloendothelial disease or condition. Typically, polymorphic molecular markers (e.g., STRP's, SNP's, RFLP's, VNTR's) are identified within the region, linkage and map distance between the markers is then established, and then linkage is established between phenotype and the various individual molecular markers. Maps can be produced for an individual family, selected populations, patient populations, etc. In general, these methods involve identifying a marker associated with the disease (e.g., identifying a polymorphism in a family which is linked to the disease) and then analyzing the surrounding DNA to identity the gene responsible for the phenotype.

The present invention also relates to methods of expressing a polynucleotide in spleen, lymphoid, and/or reticuloendothelial tissue, comprising, e.g., inserting a polynucleotide, which is operably linked to an expression control sequence, into the spleen, lymphoid, and/or reticuloendothelial gene complex at chromosomal location 11q12.2 of a target cell, and growing said cell under conditions effective to express said polynucleotide.

The polynucleotide of interest can be inserted into the target chromosomal region by any suitable method, including, e.g., by gene targeting methods, such as homologous recombination, or by random insertion methods where transformed cells are subsequently screened for insertion into the desired chromosomal site. Chromosome engineering methods are discussed in more detail below, e.g., in the section on transgenic animals. By the phrase "spleen, lymphoid, and/or reticuloendothelial gene complex," it is meant the region of the chromosome in which the cluster of genes, e.g., TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), and TMD0621 (XM_166205), of the present invention are located. Inserting an expressible polynucleotide (e.g., a polynucleotide operably linked to a promoter sequence) into this region confers the tissue expression selectivity which is characteristic of the gene cluster. Any polynucleotide of interest can be inserted into the chromosomal region, including, e.g., polynucleotides encoding polypeptides, antisense polynucleotides, etc.

A cell comprising a polynucleotide inserted into the target chromosomal location can be utilized in vitro or in vivo, e.g., in a transgenic animal. The cell is grown under conditions

5

10

15

20

25

which are suitable to achieve polynucleotide expression. These conditions depend upon the cell's environment, e.g., tissue culture cell, or in the form of a transgenic animal.

-45-

Pancreas membrane protein genes

5

10

15

20

25

30

The present invention relates to all facets of pancreas membrane protein genes, polypeptides encoded by them, antibodies and specific binding partners thereto, and their applications to research, diagnosis, drug discovery, therapy, clinical medicine, forensic science and medicine, etc. The polynucleotides and polypeptides are useful in variety of ways, including, but not limited to, as molecular markers, as drug targets, and for detecting, diagnosing, staging, monitoring, prognosticating, preventing or treating, determining predisposition to, etc., diseases and conditions, such as pancreatic cancer, diabetes, pancreatitis, and other disorders especially relating to the pancreas and the functions its performs. The identification of specific genes, and groups of genes, expressed in pathways physiologically relevant to pancreas tissue permits the definition of functional and disease pathways, and the delineation of targets in these pathways which are useful in diagnostic, therapeutic, and clinical applications. The present invention also relates to methods of using the polynucleotides and related products (proteins, antibodies, etc.) in business and computer-related methods, e.g., advertising, displaying, offering, selling, etc., such products for sale, commercial use, licensing, etc.

The function, structure, and diseases of the pancreas were described previously. The polynucleotides, polypeptides, and ligands thereto, of the present invention can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of pancreas. These include, but are not limited to, e.g., acute and chronic pancreatitis, pancreatic abscess, pancreatic pseudocyst, nonalcoholic pancreatitis, alcoholic pancreatitis, classic acute hemorrhagic pancreatitis, chronic calcifying pancreatitis, familial hereditary pancreatitis, carcinomas of the pancreas, primary (idiopathic) diabetes (e.g., Type I (insulin dependent diabetes mellitus, IDDM) [insulin deficiency, beta cell depletion], Type II (non-insulin dependent diabetes mellitus, NIDDM) [insulin resistance, relative insulin deficiency, mild beta cell depletion]), nonobese NIDDM, obese NIDDM, maturity-onset diabetes of the young (MODY), islet cell tumors, diffuse hyperplasia of the islets of Langerhans, benign adenomas, malignant islet tumors, hyperfunction of the islets of Langerhans, hyperinsulinism and hypoglycemia, Zollinger-Ellison syndrome, beta cell

tumors (insulinoma), alpha cell tumors (glucagonoma), delta cell tumors (somatostatinoma), vipoma (diarrheogenic islet cell tumor), pancreatic cancers, pancreatic carcinoid tumors, multihormonal tumors, multiple endocrine neoplasia (MEN), MEN I (Wermer syndrome), MEN II (Sipple syndrome), MEN III or IIb, pancreatic endocrine tumors, etc.

For example, five different pancreatic tumor samples were examined (Nos. 1, 2, 3, 4, and 5). TMD0639 was up-regulated in about 1/5 pancreatic cancers (No. 4), TMD0645 was up-regulated in about 3/5 pancreatic cancers (Nos. 2, 3, and 5), and TMD1127 was up-regulated in about 2/5 pancreatic cancers (Nos. 1 and 4). These results indicate that the probes can be used in combination in order to maximize the detection of different types of pancreatic cancers and tumors. Thus, a sample from a patient can be assesses for expression of both TMD0645 and TMD1127 to increase the probability that the pancreas cancer will be detected.

In view of their selectivity and display on the cell surface, the membrane proteins of the present invention are useful targets for histological, diagnostic, and therapeutic applications relating to the cells (e.g., pancreatic progenitor, exocrine, endocrine, acinar, islet, alpha, beta, delta, F, D1, enterochromaffin, etc.) in which they are expressed. Antibodies and other protein binding partners (e.g., ligands, aptamers, small peptides, etc.) can be used to selectively target agents to a tissue for any purpose, included, but not limited to, imaging, therapeutic, diagnostic, drug delivery, gene therapy, etc. For example, binding partners, such as antibodies, can be used to treat carcinomas in analogy to how c-erbB-2 antibodies are used to breast cancer. They can also be used to detect metastatic cells in biopsies and other tissue samples. The genes and polypeptides encoded thereby can also be used in tissue engineering to identify tissues as they appear during the differentiation process, to target tissues, to modulate tissue growth (e.g., from starting stem cell populations), etc. Useful antibodies or other binding partners include those that are specific for parts of the polypeptide which are exposed extracellularly as indicated in Table 21. Any of the methods described above and below can be accomplished in vivo, in vitro, or ex vivo.

When expression is described as being "predominantly" in a given tissue, this indicates that the gene's mRNAs levels are highest in this tissue as compared to the other tissues in which it was measured. Expression can also be "selective," where expression is observed. By the phrase "selectively expressed," it is meant that a nucleic acid molecule

5

10

15

20

25

-47-

comprising the defined sequence of nucleotides, when produced as a transcript, is characteristic of the tissue or cell-type in which it is made. This can mean that the transcript is expressed only in that tissue and in no other tissue-type, or it can mean that the transcript is expressed preferentially, differentially, and more abundantly (e.g., at least 5-fold, 10-fold, etc., or more) in that tissue when compared to other tissue-types.

Table 20 is a summary of the genes of the present invention which are expressed selectively and/or predominantly in pancreas tissue. Fig. 12 is an illustration of these expression patterns. Each gene is associated with a Clone ID and Accession Number ("ACCN"). The Clone ID is an arbitrary identification number for the clone, and the accession number is the number by which it is listed in GenBank. Although specific sequences are disclosed herein, and listed in GenBank by an accession number), the present invention includes all forms of the gene, including polymorphisms, allelic variations, SNPs, splice variants, and any full-length versions when the disclosed or Genbank version is partial. For convenience, these genes, and their homologs in other species, are referred to throughout the disclosure in shorthand as "the genes of Table 20," "a gene of Table 20," "polynucleotides of Table 20," "tolly peptides of Table 20," etc..., because Table 20 contains a listing of the genes by accession number and clone ID.

The expression patterns of the selectively and/or predominantly expressed polynucleotides disclosed herein can be described as a "fingerprint" in that they are a distinctive pattern displayed by pancreas tissue. Just as with a fingerprint, an expression pattern can be used as a unique identifier to characterize the status of a tissue sample. The list of expressed sequences disclosed herein provides an example of such a tissue expression profile. It can be used as a point of reference to compare and characterize samples. Tissue fingerprints can be used in many ways, e.g., to classify an unknown tissue, to determine the origin of metastatic cells, to assess the physiological status of a tissue, to determine the effect of a particular treatment regime on a tissue, to evaluate the toxicity of a compound on a tissue of interest, etc.

For example, the pancreas-selective polynucleotides disclosed herein represent the configuration of genes expressed by a normal pancreas tissue. To determine the effect of a toxin on a tissue, a sample of tissue can be obtained prior to toxin exposure ("control") and then at one or more time points after toxin exposure ("experimental"). An array of pancreas-

5

10

15

20

25

10

15

20

25

30

selective probes can be used to assess the expression patterns for both the control and experimental samples. As discussed in more detail below, any suitable method can be used. For instance, a DNA microarray can be prepared having a set of pancreas-selective genes arranged on to a small surface area in fixed and addressable positions. RNA isolated from samples can be labeled using reverse transcriptase and radioactive nucleotides, hybridized to the array, and then expression levels determined using a detection system. Several kinds of information can be extracted: presence or absence of expression, and the corresponding expression levels. The normal tissue would be expected to express substantially all the genes represented by the tissue-selective probes. The various experimental conditions can be compared to it to determine whether a gene is expressed, and how its levels match up to the normal control.

While the expression profile of the complete gene set represented by the sequences disclosed here may be most informative, a fingerprint containing expression information from less than the full collection can be useful, as well. In the same way that an incomplete fingerprint may contain enough of the pattern of whorls, arches, loops, and ridges, to identify the individual, a cell expression fingerprint containing less than the full complement may be adequate to provide useful and unique identifying and other information about the sample. Moreover, because of heterogeneity of the population, as well differences in the particular physiological state of the tissue, a tissue's "normal" expression profile is expected to differ between samples, albeit in ways that do not change the overall expression pattern. As a result, a complete match with a particular tissue expression profile, as shown herein, is not necessary.

The present invention relates to methods of detecting pancreas cells, comprising one or more of the following steps, e.g., contacting a sample comprising cells with a polynucleotide specific for a gene of Table 20, or a mammalian homolog thereof, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization. Detecting can be accomplished by any suitable method and technology, including, e.g., any of those mentioned and discussed below, such as Northern blot and PCR. Specific polynucleotides include the primer sequences shown in Table 23, and complements thereto.

Detection can also be achieved using binding partners, such as antibodies (e.g.,

monoclonal or polyclonal antibodies) that specifically recognize polypeptides coded for by genes of the present invention. Thus, the present invention relates to methods of detecting a pancreas cell, comprising, one or more the following steps, e.g. contacting a sample comprising cells with a binding partner (e.g. an antibody, an Fab fragment, a single-chain antibody, an aptamer) specific for a polypeptide coded for by a polypeptide of Table 20, or a mammalian homolog thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding. Protein binding assays can be accomplished routinely, e.g., using immunocytochemistry, ELISA format, Western blots, etc. Useful epitopes include those exposed to the surface.

As indicated above, binding partners can be used to deliver agents specifically to the pancreas, e.g., for diagnostic, therapeutic, and prognostic purposes. Methods of delivering an agent to a pancreas cell can comprise, e.g., contacting a pancreas cell with an agent coupled to a binding partner specific for a polypeptide coding for a gene of Table 20, whereby said agent is delivered to said cell. Any type of agent can be used, including, therapeutic and imaging agents. Contact with the pancreas can be achieved in any effective manner, including by administering effective amounts of the agent to a host orally, parentally, locally, systemically, intravenously, etc. The phrase "an agent coupled to binding partner" indicates that the agent is associated with the binding partner in such a manner that it can be carried specifically to the target site. Coupling includes, chemical bonding, covalent bonding, noncovalent bonding (where such bonding is sufficient to carry the agent to the target), present in a liposome or in a lipid membrane, associated with a carrier, such as a polymeric carrier, etc. The agent can be directly linked to the binding partner, or via chemical linkers or spacers. Any cell expressing a polypeptide coded for by a gene of Table 20 can be targeted, including, e.g., pancreatic progenitor, exocrine, endocrine, secretory, acinar, islet, alpha, beta, delta, F, D1, enterochromaffin, etc.

Imaging of specific organs can be facilitated using tissue selective antibodies and other binding partners that selectively target contrast agents to a specific site in the body. Various imaging techniques have been used in this context, including, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic. A reporter agent can be conjugated or associated routinely with a binding partner. Ultrasound contrast agents combined with binding partners, such as antibodies, are described in, e.g., U.S. Pat. Nos, 6,264,917,

5

10

15

20

25

10

15

20

25

30

6,254,852, 6,245,318, and 6,139,819. MRI contrast agents, such as metal chelators, radionucleotides, paramagnetic ions, etc., combined with selective targeting agents are also described in the literature, e.g., in U.S. Pat. Nos. 6,280,706 and 6,221,334. The methods described therein can be used generally to associate a partner with an agent for any desired purpose. See, Bruehlmeier et al., *Nucl. Med. Biol.*, 29:321-327, 2002, for imaging pancreas using labeled receptor ligands. Antibodies and other ligands to receptors of the present invention can be used analogously.

A pancreas cell (see above for examples of pancreas cell types) can also be modulated in accordance with the present invention, e.g., by methods of modulating a pancreas cell, comprising, e.g., contacting said cell with an agent effective to modulate a gene of Table 20, or the biological activity of a polypeptide encoded thereby (e.g., SEQ ID NO 215, 217, 219, 221, 223, 225, 227, 229, 231, 233, 235, 237, 239, 241, 243, 245, 247, 249, 251, 253, and 255), or a mammalian homolog thereof, whereby said pancreas cell is modulated. Modulation as used throughout includes, e.g., stimulating, increasing, agonizing, activating, amplifying, blocking, inhibiting, reducing, antagonizing, preventing, decreasing, diminishing, etc.

An activity or function of the pancreas cell can be modulated, including, e.g., regulation of blood sugar, modulation of all aspects of the various secreted polypeptides (hormones, enzymes, etc.) produced by the pancreas, ligand-binding, exocytosis, amylase (and any of the other 20 or so digestive enzymes produced by the pancreas) secretion, autocrine responses, apoptosis (e.g., in the survival of beta-islet cells), etc.

The present invention also relates to polypeptide detection methods for assessing pancreas function, e.g., methods of assessing pancreas function, comprising, detecting a polypeptide coded for by a gene of Table 20, fragments thereof, polymorphisms thereof, in a body fluid, whereby the level of said polypeptide in said fluid is a measure of pancreas function. Pancreas function tests are usually performed to determine whether the pancreas is functioning normally as a way of diagnosing pancreas disease. Various tests are commonly used, including, e.g., assays for the presence of pancreatic enzymes in body fluids (e.g., amylase, serum lipase, serum trypsin-like immuoreactivity), studies of pancreatic structure (e.g., using x-ray, sonography, CT-scan, angiography, endoscopic retrograde cholangiopancreatography), and tests for pancreatic function (e.g., secretin-pancreozymin

-51-

(CCK) tst, Lundh meal test, Bz-Ty-PABA test, chymotrypsin in feces, etc). Detection of a polypeptide coded for by a gene of Table 20 provides an additional assessment tool, especially in diseases such as pancreatitis and pancreatic cancer where pancreatic markers can appear in the blood, stool, urine, and other body fluids. As with the other tests, elevated levels of said polypeptide in blood, or other fluids, can indicate impaired pancreas function. Values can be determined routinely, as they are for other markers, such as those mentioned above. Detecting can be performed routinely (see below), e.g., using an antibody which is specific for said polypeptide, by RIA, ELISA, or Western blot, etc., in analogy to the tests for pancreatic enzymes in body fluids.

Promoter sequences obtained from genes of the present invention can be utilized to selectively express heterologous genes in pancreas cells. Methods of expressing a heterologous polynucleotide in pancreas cells can comprise, e.g., expressing a nucleic acid construct in pancreas cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected SEQ ID NO 258, 261, 262, 265-267, 270-272, 275, 278, 279, 282-284, 287, 290-293, 296, 297, 303, 306, 309-314, 317-320, 323-326, 329, 332-333, 336-338, 341, and 344 as shown in Table 23. In addition to the cell lines mentioned below, the construct can be expressed in primary cells or in established cell lines.

The genes and polypeptides of Table 20 can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions of the pancreas as mentioned above. The present invention relates to methods of identifying a pancreatic disease or pancreatic disease-susceptibility, comprising, e.g., determining the association of a pancreatic disease or pancreatic disease-susceptibility with a nucleotide sequence present within the pancreatic gene complex. An association between a pancreas disease or disease-susceptibility and nucleotide sequence includes, e.g., establishing (or finding) a correlation (or relationship) between a DNA marker (e.g., gene, VNTR, polymorphism, EST, etc.) and a particular disease state. Once a relationship is identified, the DNA marker can be utilized in diagnostic tests and as a drug target.

Human linkage maps can be constructed to establish a relationship between the cytogenetic locus as shown in Table 22 and a pancreatic disease or condition. Typically, polymorphic molecular markers (e.g., STRP's, SNP's, RFLP's, VNTR's) are identified

5

10

15

20

25

-52-

within the region, linkage and map distance between the markers is then established, and then linkage is established between phenotype and the various individual molecular markers. Maps can be produced individual family, selected populations, patient populations, etc. In general, these methods involve identifying a marker associated with the disease (e.g., identifying a polymorphism in a family which is linked to the disease) and then analyzing the surrounding DNA to identity the gene responsible for the phenotype.

Nucleic acids

5

10

15

20

25

30

A mammalian polynucleotide, or fragment thereof, of the present invention is a polynucleotide having a nucleotide sequence obtainable from a natural source. When the species name is used, e.g., a human, it indicates that the polynucleotide or polypeptide is obtainable from a natural source. It therefore includes naturally-occurring normal, naturally-occurring mutant, and naturally-occurring polymorphic alleles (e.g., SNPs), differentially-spliced transcripts, splice-variants, etc. By the term "naturally-occurring," it is meant that the polynucleotide is obtainable from a natural source, e.g., animal tissue and cells, body fluids, tissue culture cells, forensic samples. Natural sources include, e.g., living cells obtained from tissues and whole organisms, tumors, cultured cell lines, including primary and immortalized cell lines. Naturally-occurring mutations can include deletions (e.g., a truncated amino- or carboxy-terminus), substitutions, inversions, or additions of nucleotide sequence. These genes can be detected and isolated by polynucleotide hybridization according to methods which one skilled in the art would know, e.g., as discussed below.

A polynucleotide according to the present invention can be obtained from a variety of different sources. It can be obtained from DNA or RNA, such as polyadenylated mRNA or total RNA, e.g., isolated from tissues, cells, or whole organism. The polynucleotide can be obtained directly from DNA or RNA, from a cDNA library, from a genomic library, etc. The polynucleotide can be obtained from a cell or tissue (e.g., from an embryonic or adult tissues) at a particular stage of development, having a desired genotype, phenotype, disease status, etc.

The polynucleotides described herein can be partial sequences that correspond to full-length, naturally-occurring transcripts. The present invention includes, as well, full-length polynucleotides that comprise these partial sequences, e.g., genomic DNAs and polynucleotides comprising a start and stop codon, a start codon and a polyA tail, a

-53-

transcription start and a polyA tail, etc. These sequences can be obtained by any suitable method, e.g., using a partial sequence as a probe to select a full-length cDNA from a library containing full-length inserts. A polynucleotide which "codes without interruption" refers to a polynucleotide having a continuous open reading frame ("ORF") as compared to an ORF which is interrupted by introns or other noncoding sequences.

Polynucleotides and polypeptides can be excluded as compositions from the present invention if, e.g., listed in a publicly available databases on the day this application was filed and/or disclosed in a patent application having an earlier filing or priority date than this application and/or conceived and/or reduced to practice earlier than a polynucleotide in this application.

As described herein, the phrase "an isolated polynucleotide which is SEQ ID NO," or "an isolated polynucleotide which is selected from SEQ ID NO," refers to an isolated nucleic acid molecule from which the recited sequence was derived (e.g., a cDNA derived from mRNA; cDNA derived from genomic DNA). Because of sequencing errors, typographical errors, etc., the actual naturally-occurring sequence may differ from a SEQ ID listed herein. Thus, the phrase indicates the specific molecule from which the sequence was derived, rather than a molecule having that exact recited nucleotide sequence, analogously to how a culture depository number refers to a specific cloned fragment in a cryotube.

As explained in more detail below, a polynucleotide sequence of the invention can contain the complete sequence as shown herein, degenerate sequences thereof, anti-sense, muteins thereof, genes comprising said sequences, full-length cDNAs comprising said sequences, complete genomic sequences, fragments thereof, homologs, primers, nucleic acid molecules which hybridize thereto, derivatives thereof, etc.

25 Genomic

5

10

15

20

30

The present invention also relates genomic DNA from which the polynucleotides of the present invention can be derived. A genomic DNA coding for a human, mouse, or other mammalian polynucleotide, can be obtained routinely, for example, by screening a genomic library (e.g., a YAC library) with a polynucleotide of the present invention, or by searching nucleotide databases, such as GenBank and EMBL, for matches. Promoter and other regulatory regions (including both 5' and 3' regions, as well introns) can be identified

upstream or downstream of coding and expressed RNAs, and assayed routinely for activity, e.g., by joining to a reporter gene (e.g., CAT, GFP, alkaline phosphatase, luciferase, galatosidase). A promoter obtained from a tissue selective gene can be used, e.g., in gene therapy to obtain tissue-specific expression of a heterologous gene (e.g., coding for a therapeutic product or cytotoxin). 5' and 3' sequences (including, UTRs and introns) can be used to modulate or regulate stability, transcription, and translation of nucleic acids, including the sequence to which is attached in nature, as well as heterologous nucleic acids.

Constructs

5

10

15

20

25

30

A polynucleotide of the present invention can comprise additional polynucleotide sequences, e.g., sequences to enhance expression, detection, uptake, cataloging, tagging, etc. A polynucleotide can include only coding sequence; a coding sequence and additional non-naturally occurring or heterologous coding sequence (e.g., sequences coding for leader, signal, secretory, targeting, enzymatic, fluorescent, antibiotic resistance, and other functional or diagnostic peptides); coding sequences and non-coding sequences, e.g., untranslated sequences at either a 5' or 3' end, or dispersed in the coding sequence, e.g., introns.

A polynucleotide according to the present invention also can comprise an expression control sequence operably linked to a polynucleotide as described above. The phrase "expression control sequence" means a polynucleotide sequence that regulates expression of a polypeptide coded for by a polynucleotide to which it is functionally ("operably") linked. Expression can be regulated at the level of the mRNA or polypeptide. Thus, the expression control sequence includes mRNA-related elements and protein-related elements. Such elements include promoters, enhancers (viral or cellular), ribosome binding sequences, transcriptional terminators, etc. An expression control sequence is operably linked to a nucleotide coding sequence when the expression control sequence is positioned in such a manner to effect or achieve expression of the coding sequence. For example, when a promoter is operably linked 5' to a coding sequence, expression of the coding sequence is driven by the promoter. Expression control sequences can include an initiation codon and additional nucleotides to place a partial nucleotide sequence of the present invention in-frame in order to produce a polypeptide (e.g., pET vectors from Promega have been designed to permit a molecule to be inserted into all three reading frames to identify the one that results

in polypeptide expression). Expression control sequences can be heterologous or endogenous to the normal gene.

-55-

A polynucleotide of the present invention can also comprise nucleic acid vector sequences, e.g., for cloning, expression, amplification, selection, etc. Any effective vector can be used. A vector is, e.g., a polynucleotide molecule which can replicate autonomously in a host cell, e.g., containing an origin of replication. Vectors can be useful to perform manipulations, to propagate, and/or obtain large quantities of the recombinant molecule in a desired host. A skilled worker can select a vector depending on the purpose desired, e.g., to propagate the recombinant molecule in bacteria, yeast, insect, or mammalian cells. The following vectors are provided by way of example. Bacterial: pQE70, pQE60, pQE-9 (Qiagen), pBS, pD10, Phagescript, phiX174, pBK Phagemid, pNH8A, pNH16a, pNH18Z, pNH46A (Stratagene); Bluescript KS+II (Stratagene); ptrc99a, pKK223-3, pKK233-3, pDR54 0, pRIT5 (Pharmacia). Eukaryotic: PWLNEO, pSV2CAT, pOG44, pXT1, pSG (Stratagene), pSVK3, PBPV, PMSG, pSVL (Pharmacia), pCR2.1/TOPO, pCRII/TOPO, pCR4/TOPO, pTrcHisB, pCMV6-XL4, etc. However, any other vector, e.g., plasmids, viruses, or parts thereof, may be used as long as they are replicable and viable in the desired host. The vector can also comprise sequences which enable it to replicate in the host whose genome is to be modified.

20 Hybridization

5

10

15

25

30

Polynucleotide hybridization, as discussed in more detail below, is useful in a variety of applications, including, in gene detection methods, for identifying mutations, for making mutations, to identify homologs in the same and different species, to identify related members of the same gene family, in diagnostic and prognostic assays, in therapeutic applications (e.g., where an antisense polynucleotide is used to inhibit expression), etc.

The ability of two single-stranded polynucleotide preparations to hybridize together is a measure of their nucleotide sequence complementarity, e.g., base-pairing between nucleotides, such as A-T, G-C, etc. The invention thus also relates to polynucleotides, and their complements, which hybridize to a polynucleotide comprising a nucleotide sequence as set forth herein and genomic sequences thereof. A nucleotide sequence hybridizing to the latter sequence will have a complementary polynucleotide strand, or act as a template for one

10

15

20

25

30

in the presence of a polymerase (i.e., an appropriate polynucleotide synthesizing enzyme). The present invention includes both strands of polynucleotide, e.g., a sense strand and an anti-sense strand.

Hybridization conditions can be chosen to select polynucleotides which have a desired amount of nucleotide complementarity with the nucleotide sequences set forth in herein and genomic sequences thereof. A polynucleotide capable of hybridizing to such sequence, preferably, possesses, e.g., about 70%, 75%, 80%, 85%, 87%, 90%, 92%, 95%, 97%, 99%, or 100% complementarity, between the sequences. The present invention particularly relates to polynucleotide sequences which hybridize to the nucleotide sequences set forth in the attached sequence disclosure or genomic sequences thereof, under low or high stringency conditions. These conditions can be used, e.g., to select corresponding homologs in non-human species.

Polynucleotides which hybridize to polynucleotides of the present invention can be selected in various ways. Filter-type blots (i.e., matrices containing polynucleotide, such as nitrocellulose), glass chips, and other matrices and substrates comprising polynucleotides (short or long) of interest, can be incubated in a prehybridization solution (e.g., 6X SSC, 0.5% SDS, 100 μg/ml denatured salmon sperm DNA, 5X Denhardt's solution, and 50% formamide), at 22-68°C, overnight, and then hybridized with a detectable polynucleotide probe under conditions appropriate to achieve the desired stringency. In general, when high homology or sequence identity is desired, a high temperature can be used (e.g., 65 °C). As the homology drops, lower washing temperatures are used. For salt concentrations, the lower the salt concentration, the higher the stringency. The length of the probe is another consideration. Very short probes (e.g., less than 100 base pairs) are washed at lower temperatures, even if the homology is high. With short probes, formamide can be omitted. See, e.g., *Current Protocols in Molecular Biology*, Chapter 6, Screening of Recombinant Libraries; Sambrook et al., *Molecular Cloning*, 1989, Chapter 9.

For instance, high stringency conditions can be achieved by incubating the blot overnight (e.g., at least 12 hours) with a polynucleotide probe in a hybridization solution containing, e.g., about 5X SSC, 0.1-0.5% SDS, 100 µg/ml denatured salmon sperm DNA and 50% formamide, at 42°C, or hybridizing at 42°C in 5X SSPE, 0.1-0.5% SDS, and 50%

10

15

20

25

30

formamide, 100 μ g/ml denatured salmon sperm DNA, and washing at 65°C in 0.1% SSC and 0.1% SDS.

Blots can be washed at high stringency conditions that allow, e.g., for less than 5% bp mismatch (e.g., wash twice in 0.1% SSC and 0.1% SDS for 30 min at 65°C), i.e., selecting sequences having 95% or greater sequence identity.

Other non-limiting examples of high stringency conditions includes a final wash at 65°C in aqueous buffer containing 30 mM NaCl and 0.5% SDS. Another example of high stringent conditions is hybridization in 7% SDS, 0.5 M NaPO₄, pH 7, 1 mM EDTA at 50°C, e.g., overnight, followed by one or more washes with a 1% SDS solution at 42°C. Whereas high stringency washes can allow for, e.g., less than 10%, less than 5% mismatch, etc., reduced or low stringency conditions can permit up to 20% nucleotide mismatch. Hybridization at low stringency can be accomplished as above, but using lower formamide

conditions, lower temperatures and/or lower salt concentrations, as well as longer periods of

incubation time.

Hybridization can also be based on a calculation of melting temperature (Tm) of the hybrid formed between the probe and its target, as described in Sambrook et al.. Generally, the temperature Tm at which a short oligonucleotide (containing 18 nucleotides or fewer) will melt from its target sequence is given by the following equation: Tm = (number of A's and T's) x 2°C + (number of C's and G's) x 4°C. For longer molecules, Tm = 81.5 + 16.6 log₁₀[Na⁺] + 0.41(%GC) - 600/N where [Na⁺] is the molar concentration of sodium ions, %GC is the percentage of GC base pairs in the probe, and N is the length. Hybridization can be carried out at several degrees below this temperature to ensure that the probe and target can hybridize. Mismatches can be allowed for by lowering the temperature even further.

Stringent conditions can be selected to isolate sequences, and their complements, . which have, e.g., at least about 90%, 95%, or 97%, nucleotide complementarity between the probe (e.g., a short polynucleotide of the sequences disclosed herein or genomic sequences thereof) and a target polynucleotide.

Other homologs of polynucleotides of the present invention can be obtained from mammalian and non-mammalian sources according to various methods. For example, hybridization with a polynucleotide can be employed to select homologs, e.g., as described in Sambrook et al., *Molecular Cloning*, Chapter 11, 1989. Such homologs can have varying

amounts of nucleotide and amino acid sequence identity and similarity to such polynucleotides of the present invention. Mammalian organisms include, e.g., mice, rats, monkeys, pigs, cows, etc. Non-mammalian organisms include, e.g., vertebrates, invertebrates, zebra fish, chicken, Drosophila, C. elegans, Xenopus, yeast such as S. pombe, S. cerevisiae, roundworms, prokaryotes, plants, Arabidopsis, artemia, viruses, etc. The degree of nucleotide sequence identity between human and mouse can be about, e.g. 70% or more, 85% or more for open reading frames, etc.

Alignment

5

10

15

20

25

30

Alignments can be accomplished by using any effective algorithm. For pairwise alignments of DNA sequences, the methods described by Wilbur-Lipman (e.g., Wilbur and Lipman, Proc. Natl. Acad. Sci., 80:726-730, 1983) or Martinez/Needleman-Wunsch (e.g., Martinez, Nucleic Acid Res., 11:4629-4634, 1983) can be used. For instance, if the Martinez/Needleman-Wunsch DNA alignment is applied, the minimum match can be set at 9, gap penalty at 1.10, and gap length penalty at 0.33. The results can be calculated as a similarity index, equal to the sum of the matching residues divided by the sum of all residues and gap characters, and then multiplied by 100 to express as a percent. Similarity index for related genes at the nucleotide level in accordance with the present invention can be greater than 70%, 80%, 85%, 90%, 95%, 99%, or more. Pairs of protein sequences can be aligned by the Lipman-Pearson method (e.g., Lipman and Pearson, Science, 227:1435-1441, 1985) with k-tuple set at 2, gap penalty set at 4, and gap length penalty set at 12. Results can be expressed as percent similarity index, where related genes at the amino acid level in accordance with the present invention can be greater than 65%, 70%, 75%, 80%, 85%, 90%, 95%, 99%, or more. Various commercial and free sources of alignment programs are available, e.g., MegAlign by DNA Star, BLAST (National Center for Biotechnology Information), BCM (Baylor College of Medicine) Launcher, etc. BLAST can be used to calculate amino acid sequence identity, amino acid sequence homology, and nucleotide sequence identity. These calculations can be made along the entire length of each of the target sequences which are to be compared.

After two sequences have been aligned, a "percent sequence identity" can be determined. For these purposes, it is convenient to refer to a Reference Sequence and a

10

15

20

25

30

Compared Sequence, where the Compared Sequence is *compared* to the Reference Sequence. Percent sequence identity can be determined according to the following formula: Percent Identity = 100 [1-(C/R)], wherein C is the number of differences between the Reference Sequence and the Compared Sequence over the length of alignment between the Reference Sequence and the Compared Sequence where (i) each base or amino acid in the Reference Sequence that does not have a corresponding aligned base or amino acid in the Compared Sequence, (ii) each gap in the Reference Sequence, (iii) each aligned base or amino acid in the Reference Sequence that is different from an aligned base or amino acid in the Compared Sequence, constitutes a difference; and R is the number of bases or amino acids in the Reference Sequence over the length of the alignment with the Compared Sequence with any gap created in the Reference Sequence also being counted as a base or amino acid.

Percent sequence identity can also be determined by other conventional methods, e.g., as described in Altschul et al., *Bull. Math. Bio.* 48: 603-616, 1986 and Henikoff and Henikoff, *Proc. Natl. Acad. Sci.* USA 89:10915-10919, 1992.

Specific polynucleotide probes

A polynucleotide of the present invention can comprise any continuous nucleotide sequence described herein, sequences which share sequence identity thereto, or complements thereof. The term "probe" refers to any substance that can be used to detect, identify, isolate, etc., another substance. A polynucleotide probe is comprised of nucleic acid can be used to detect, identify, etc., other nucleic acids, such as DNA and RNA.

These polynucleotides can be of any desired size that is effective to achieve the specificity desired. For example, a probe can be from about 7 or 8 nucleotides to several thousand nucleotides, depending upon its use and purpose. For instance, a probe used as a primer PCR can be shorter than a probe used in an ordered array of polynucleotide probes. Probe sizes vary, and the invention is not limited in any way by their size, e.g., probes can be from about 7-2000 nucleotides, 7-1000, 8-700, 8-600, 8-500, 8-400, 8-300, 8-150, 8-100, 8-75, 7-50, 10-25, 14-16, at least about 8, at least about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, or more, etc. The polynucleotides can have non-naturally-occurring nucleotides, e.g., inosine, AZT, 3TC, etc. The polynucleotides can have 100% sequence identity or complementarity to a sequence disclosed herein, or it can have mismatches or

10

15

20

25

30

nucleotide substitutions, e.g., 1, 2, 3, 4, or 5 substitutions. The probes can be single-stranded or double-stranded.

In accordance with the present invention, a polynucleotide can be present in a kit, where the kit includes, e.g., one or more polynucleotides, a desired buffer (e.g., phosphate, tris, etc.), detection compositions, RNA or cDNA from different tissues to be used as controls, libraries, etc. The polynucleotide can be labeled or unlabeled, with radioactive or non-radioactive labels as known in the art. Kits can comprise one or more pairs of polynucleotides for amplifying nucleic acids specific for tissue selective genes, e.g., comprising a forward and reverse primer effective in PCR. These include both sense and anti-sense orientations. For instance, in PCR-based methods (such as RT-PCR), a pair of primers are typically used, one having a sense sequence and the other having an antisense sequence.

Another aspect of the present invention is a nucleotide sequence that is specific to, or for, a selective polynucleotide. The phrases "specific for" or "specific to" a polynucleotide have a functional meaning that the polynucleotide can be used to identify the presence of one or more target genes in a sample and distinguish them from non-target genes. It is specific in the sense that it can be used to detect polynucleotides above background noise ("non-specific binding"). A specific sequence is a defined order of nucleotides (or amino acid sequences, if it is a polypeptide sequence) which occurs in the polynucleotide, e.g., in the nucleotide sequences of the present invention, and which is characteristic of that target sequence, and substantially no non-target sequences. A probe or mixture of probes can comprise a sequence or sequences that are specific to a plurality of target sequences, e.g., where the sequence is a consensus sequence, a functional domain, etc., e.g., capable of recognizing a family of related genes. Such sequences can be used as probes in any of the methods described herein or incorporated by reference. Both sense and antisense nucleotide sequences are included. A specific polynucleotide according to the present invention can be determined routinely.

A polynucleotide comprising a specific sequence can be used as a hybridization probe to identify the presence of, e.g., human or mouse polynucleotide, in a sample comprising a mixture of polynucleotides, e.g., on a Northern blot. Hybridization can be performed under high stringent conditions (see, above) to select polynucleotides (and their complements which

-61-

can contain the coding sequence) having at least 90%, 95%, 99%, etc., identity (i.e., complementarity) to the probe, but less stringent conditions can also be used. A specific polynucleotide sequence can also be fused in-frame, at either its 5' or 3' end, to various nucleotide sequences as mentioned throughout the patent, including coding sequences for enzymes, detectable markers, GFP, etc, expression control sequences, etc.

A polynucleotide probe, especially one that is specific to a polynucleotide of the present invention, can be used in gene detection and hybridization methods as already described. In one embodiment, a specific polynucleotide probe can be used to detect whether a particular tissue or cell-type is present in a target sample. To carry out such a method, a selective polynucleotide can be chosen which is characteristic of the desired target tissue. Such polynucleotide is preferably chosen so that it is expressed or displayed in the target tissue, but not in other tissues which are present in the sample. For instance, if detection of pancreas, or kidney, it may not matter whether the selective polynucleotide is expressed in other tissues, as long as it is not expressed in cells normally present in blood, e.g., peripheral blood mononuclear cells. Starting from the selective polynucleotide, a specific polynucleotide probe can be designed which hybridizes (if hybridization is the basis of the assay) under the hybridization conditions to the selective polynucleotide, whereby the presence of the selective polynucleotide can be determined.

Probes which are specific for polynucleotides of the present invention can also be prepared using involve transcription-based systems, e.g., incorporating an RNA polymerase promoter into a selective polynucleotide of the present invention, and then transcribing antisense RNA using the polynucleotide as a template. See, e.g., U.S. Pat. No. 5,545,522.

¹ Polynucleotide composition

A polynucleotide according to the present invention can comprise, e.g., DNA, RNA, synthetic polynucleotide, peptide polynucleotide, modified nucleotides, dsDNA, ssDNA, ssRNA, dsRNA, and mixtures thereof. A polynucleotide can be single- or double-stranded, triplex, DNA:RNA, duplexes, comprise hairpins, and other secondary structures, etc. Nucleotides comprising a polynucleotide can be joined via various known linkages, e.g., ester, sulfamate, sulfamide, phosphorothioate, phosphoramidate, methylphosphonate, carbamate, etc., depending on the desired purpose, e.g., resistance to nucleases, such as

5

10

15

20

25

RNAse H, improved in vivo stability, etc. See, e.g., U.S. Pat. No. 5,378,825. Any desired nucleotide or nucleotide analog can be incorporated, e.g., 6-mercaptoguanine, 8-oxo-guanine, etc.

Various modifications can be made to the polynucleotides, such as attaching detectable markers (avidin, biotin, radioactive elements, fluorescent tags and dyes, energy transfer labels, energy-emitting labels, binding partners, etc.) or moieties which improve hybridization, detection, and/or stability. The polynucleotides can also be attached to solid supports, e.g., nitrocellulose, magnetic or paramagnetic microspheres (e.g., as described in U.S. Pat. No. 5,411,863; U.S. Pat. No. 5,543,289; for instance, comprising ferromagnetic, supermagnetic, paramagnetic, superparamagnetic, iron oxide and polysaccharide), nylon, agarose, diazotized cellulose, latex solid microspheres, polyacrylamides, etc., according to a desired method. See, e.g., U.S. Pat. Nos. 5,470,967, 5,476,925, and 5,478,893.

Polynucleotide according to the present invention can be labeled according to any desired method. The polynucleotide can be labeled using radioactive tracers such as ³²P, ³⁵S, ³H, or ¹⁴C, to mention some commonly used tracers. The radioactive labeling can be carried out according to any method, such as, for example, terminal labeling at the 3' or 5' end using a radiolabeled nucleotide, polynucleotide kinase (with or without dephosphorylation with a phosphatase) or a ligase (depending on the end to be labeled). A non-radioactive labeling can also be used, combining a polynucleotide of the present invention with residues having immunological properties (antigens, haptens), a specific affinity for certain reagents (ligands), properties enabling detectable enzyme reactions to be completed (enzymes or coenzymes, enzyme substrates, or other substances involved in an enzymatic reaction), or characteristic physical properties, such as fluorescence or the emission or absorption of light at a desired wavelength, etc.

25

30

5

10

15

20

Nucleic acid detection methods

Another aspect of the present invention relates to methods and processes for detecting tissue selective genes. Detection methods have a variety of applications, including for diagnostic, prognostic, forensic, and research applications. To accomplish gene detection, a polynucleotide in accordance with the present invention can be used as a "probe." The term "probe" or "polynucleotide probe" has its customary meaning in the art, e.g., a polynucleotide

10

15

20

25

30

which is effective to identify (e.g., by hybridization), when used in an appropriate process, the presence of a target polynucleotide to which it is designed. Identification can involve simply determining presence or absence, or it can be quantitative, e.g., in assessing amounts of a gene or gene transcript present in a sample. Probes can be useful in a variety of ways, such as for diagnostic purposes, to identify homologs, and to detect, quantitate, or isolate a polynucleotide of the present invention in a test sample.

Assays can be utilized which permit quantification and/or presence/absence detection of a target nucleic acid in a sample. Assays can be performed at the single-cell level, or in a sample comprising many cells, where the assay is "averaging" expression over the entire collection of cells and tissue present in the sample. Any suitable assay format can be used, including, but not limited to, e.g., Southern blot analysis, Northern blot analysis, polymerase chain reaction ("PCR") (e.g., Saiki et al., Science, 241:53, 1988; U.S. Pat. Nos. 4,683,195, 4,683,202, and 6,040,166; PCR Protocols: A Guide to Methods and Applications, Innis et al., eds., Academic Press, New York, 1990), reverse transcriptase polymerase chain reaction ("RT-PCR"), anchored PCR, rapid amplification of cDNA ends ("RACE") (e.g., Schaefer in Gene Cloning and Analysis: Current Innovations, Pages 99-115, 1997), ligase chain reaction ("LCR") (EP 320 308), one-sided PCR (Ohara et al., Proc. Natl. Acad. Sci., 86:5673-5677, 1989), indexing methods (e.g., U.S. Pat. No. 5,508,169), in situ hybridization, differential display (e.g., Liang et al., Nucl. Acid. Res., 21:3269-3275, 1993; U.S. Pat. Nos. 5,262,311, 5,599,672 and 5,965,409; WO97/18454; Prashar and Weissman, Proc. Natl. Acad. Sci., 93:659-663, and U.S. Pat. Nos. 6,010,850 and 5,712,126; Welsh et al., Nucleic Acid Res., 20:4965-4970, 1992, and U.S. Pat. No. 5,487,985) and other RNA fingerprinting techniques, nucleic acid sequence based amplification ("NASBA") and other transcription based amplification systems (e.g., U.S. Pat. Nos. 5,409,818 and 5,554,527; WO 88/10315), polynucleotide arrays (e.g., U.S. Pat. Nos. 5,143,854, 5,424,186; 5,700,637, 5,874,219, and 6,054,270; PCT WO 92/10092; PCT WO 90/15070), Qbeta Replicase (PCT/US87/00880), Strand Displacement Amplification ("SDA"), Repair Chain Reaction ("RCR"), nuclease protection assays, subtraction-based methods, Rapid-Scan™, etc. Additional useful methods include, but are not limited to, e.g., template-based amplification methods, competitive PCR (e.g., U.S. Pat. No. 5,747,251), redox-based assays (e.g., U.S. Pat. No. 5,871,918), Taqmanbased assays (e.g., Holland et al., Proc. Natl. Acad, Sci., 88:7276-7280, 1991; U.S. Pat. Nos.

-64-

5,210,015 and 5,994,063), real-time fluorescence-based monitoring (e.g., U.S. Pat. 5,928,907), molecular energy transfer labels (e.g., U.S. Pat. Nos. 5,348,853, 5,532,129, 5,565,322, 6,030,787, and 6,117,635; Tyagi and Kramer, *Nature Biotech.*, 14:303-309, 1996). Any method suitable for single cell analysis of gene or protein expression can be used, including in situ hybridization, immunocytochemistry, MACS, FACS, flow cytometry, etc. For single cell assays, expression products can be measured using antibodies, PCR, or other types of nucleic acid amplification (e.g., Brady et al., *Methods Mol. & Cell. Biol.* 2, 17-25, 1990; Eberwine et al., 1992, *Proc. Natl. Acad. Sci.*, 89, 3010-3014, 1992; U.S. Pat. No. 5,723,290). These and other methods can be carried out conventionally, e.g., as described in the mentioned publications.

Many of such methods may require that the polynucleotide is labeled, or comprises a particular nucleotide type useful for detection. The present invention includes such modified polynucleotides that are necessary to carry out such methods. Thus, polynucleotides can be DNA, RNA, DNA:RNA hybrids, PNA, etc., and can comprise any modification or substituent which is effective to achieve detection.

Detection can be desirable for a variety of different purposes, including research, diagnostic, prognostic, and forensic. For diagnostic purposes, it may be desirable to identify the presence or quantity of a polynucleotide sequence in a sample, where the sample is obtained from tissue, cells, body fluids, etc. In a preferred method as described in more detail below, the present invention relates to a method of detecting a polynucleotide comprising, contacting a target polynucleotide in a test sample with a polynucleotide probe under conditions effective to achieve hybridization between the target and probe; and detecting hybridization.

Any test sample in which it is desired to identify a polynucleotide or polypeptide thereof can be used, including, e.g., blood, urine, saliva, stool (for extracting nucleic acid, see, e.g., U.S. Pat. No. 6,177,251), swabs comprising tissue, biopsied tissue, tissue sections, cultured cells, etc.

Detection can be accomplished in combination with polynucleotide probes for other genes, e.g., genes which are expressed in other disease states, tissues, cells, such as brain, heart, kidney, spleen, thymus, liver, stomach, small intestine, colon, muscle, lung, testis, placenta, pituitary, thyroid, skin, adrenal gland, pancreas, salivary gland, uterus, ovary,

5

10

15

20

25

WO 03/089583

-65-

PCT/US03/11497

prostate gland, peripheral blood cells (T-cells, lymphocytes, etc.), embryo, breast, fat, adult and embryonic stem cells, etc.

Polynucleotides can be used in wide range of methods and compositions, including for detecting, diagnosing, staging, grading, assessing, prognosticating, etc. diseases and disorders associated with tissue selective genes, for monitoring or assessing therapeutic and/or preventative measures, in ordered arrays, etc. Any method of detecting genes and polynucleotides can be used; certainly, the present invention is not to be limited how such methods are implemented.

Along these lines, the present invention relates to methods of detecting polynucleotides of the present invention in a sample comprising nucleic acid. Such methods can comprise one or more the following steps in any effective order, e.g., contacting said sample with a polynucleotide probe under conditions effective for said probe to hybridize specifically to nucleic acid in said sample, and detecting the presence or absence of probe hybridized to nucleic acid in said sample, wherein said probe is a polynucleotide which is described herein, a polynucleotide having, e.g., about 70%, 80%, 85%, 90%, 95%, 99%, or more sequence identity thereto, effective or specific fragments thereof, or complements thereto. The detection method can be applied to any sample, e.g., cultured primary, secondary, or established cell lines, tissue biopsy, blood, urine, stool, cerebral spinal fluid, and other bodily fluids, for any purpose.

Contacting the sample with probe can be carried out by any effective means in any effective environment. It can be accomplished in a solid, liquid, frozen, gaseous, amorphous, solidified, coagulated, colloid, etc., mixtures thereof, matrix. For instance, a probe in an aqueous medium can be contacted with a sample which is also in an aqueous medium, or which is affixed to a solid matrix, or vice-versa.

Generally, as used throughout the specification, the term "effective conditions" means, e.g., the particular milieu in which the desired effect is achieved. Such a milieu, includes, e.g., appropriate buffers, oxidizing agents, reducing agents, pH, co-factors, temperature, ion concentrations, suitable age and/or stage of cell (such as, in particular part of the cell cycle, or at a particular stage where particular genes are being expressed) where cells are being used, culture conditions (including substrate, oxygen, carbon dioxide, etc.). When hybridization is the chosen means of achieving detection, the probe and sample can be

5

10

15

20

25

-66-

combined such that the resulting conditions are functional for said probe to hybridize specifically to nucleic acid in said sample.

The phrase "hybridize specifically" indicates that the hybridization between single-stranded polynucleotides is based on nucleotide sequence complementarity. The effective conditions are selected such that the probe hybridizes to a preselected and/or definite target nucleic acid in the sample. For instance, if detection of a polynucleotide set forth herein is desired, a probe can be selected which can hybridize to such target gene under high stringent conditions, without significant hybridization to other genes in the sample. To detect homologs of a polynucleotide set forth in herein, the effective hybridization conditions can be less stringent, and/or the probe can comprise codon degeneracy, such that a homolog is detected in the sample.

As already mentioned, the methods can be carried out by any effective process, e.g., by Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, in situ hybridization, etc., as indicated above. When PCR based techniques are used, two or more probes are generally used. One probe can be specific for a defined sequence which is characteristic of a selective polynucleotide, but the other probe can be specific for the selective polynucleotide, or specific for a more general sequence, e.g., a sequence such as polyA which is characteristic of mRNA, a sequence which is specific for a promoter, ribosome binding site, or other transcriptional features, a consensus sequence (e.g., representing a functional domain). For the former aspects, 5' and 3' probes (e.g., polyA, Kozak, etc.) are preferred which are capable of specifically hybridizing to the ends of transcripts. When PCR is utilized, the probes can also be referred to as "primers" in that they can prime a DNA polymerase reaction.

In addition to testing for the presence or absence of polynucleotides, the present invention also relates to determining the amounts at which polynucleotides of the present invention are expressed in sample and determining the differential expression of such polynucleotides in samples. Such methods can involve substantially the same steps as described above for presence/absence detection, e.g., contacting with probe, hybridizing, and detecting hybridized probe, but using more quantitative methods and/or comparisons to standards.

5

10

15

20

25

-67-

The amount of hybridization between the probe and target can be determined by any suitable methods, e.g., PCR, RT-PCR, RACE PCR, Northern blot, polynucleotide microarrays, Rapid-Scan, etc., and includes both quantitative and qualitative measurements. For further details, see the hybridization methods described above and below. Determining by such hybridization whether the target is differentially expressed (e.g., up-regulated or down-regulated) in the sample can also be accomplished by any effective means. For instance, the target's expression pattern in the sample can be compared to its pattern in a known standard, such as in a normal tissue, or it can be compared to another gene in the same sample. When a second sample is utilized for the comparison, it can be a sample of normal tissue that is known not to contain diseased cells. The comparison can be performed on samples which contain the same amount of RNA (such as polyadenylated RNA or total RNA), or, on RNA extracted from the same amounts of starting tissue. Such a second sample can also be referred to as a control or standard. Hybridization can also be compared to a second target in the same tissue sample. Experiments can be performed that determine a ratio between the target nucleic acid and a second nucleic acid (a standard or control), e.g., in a normal tissue. When the ratio between the target and control are substantially the same in a normal and sample, the sample is determined or diagnosed not to contain cells. However, if the ratio is different between the normal and sample tissues, the sample is determined to contain, e.g., kidney, pancreas, or immune cells. The approaches can be combined, and one or more second samples, or second targets can be used. Any second target nucleic acid can be used as a comparison, including "housekeeping" genes, such as beta-actin, alcohol dehydrogenase, or any other gene whose expression does not vary depending upon the disease status of the cell.

25 Methods of identifying polymorphisms, mutations, etc.

Polynucleotides of the present invention can also be utilized to identify mutant alleles, SNPs, gene rearrangements and modifications, and other polymorphisms of the wild-type gene. Mutant alleles, polymorphisms, SNPs, etc., can be identified and isolated from subjects with diseases that are known, or suspected to have, a genetic component. Identification of such genes can be carried out routinely (see, above for more guidance), e.g., using PCR, hybridization techniques, direct sequencing, mismatch reactions (see, e.g.,

5

10

15

20

-68-

above), RFLP analysis, SSCP (e.g., Orita et al., Proc. Natl. Acad. Sci., 86:2766, 1992), etc., where a polynucleotide having a sequence selected from the polynucleotides of the present invention is used as a probe. The selected mutant alleles, SNPs, polymorphisms, etc., can be used diagnostically to determine whether a subject has, or is susceptible to a disorder associated with tissue selective genes disclosed herein, as well as to design therapies and predict the outcome of the disorder. Methods involve, e.g., diagnosing a disorder or determining susceptibility to a disorder, comprising, detecting the presence of a mutation in a gene represented by a polynucleotide selected from the sequences disclosed herein. The detecting can be carried out by any effective method, e.g., obtaining cells from a subject, determining the gene sequence or structure of a target gene (using, e.g., mRNA, cDNA, genomic DNA, etc), comparing the sequence or structure of the target gene to the structure of the normal gene, whereby a difference in sequence or structure indicates a mutation in the gene in the subject. Polynucleotides can also be used to test for mutations, SNPs, polymorphisms, etc., e.g., using mismatch DNA repair technology as described in U.S. Pat. No. 5,683,877; U.S. Pat. No. 5,656,430; Wu et al., Proc. Natl. Acad. Sci., 89:8779-8783, 1992.

The present invention also relates to methods of detecting polymorphisms in tissue selective genes, comprising, e.g., comparing the structure of: genomic DNA comprising all or part of a tissue selective gene, mRNA comprising all or part of a tissue selective gene, cDNA comprising all or part of a tissue selective gene, or a polypeptide comprising all or part of a tissue selective gene, with the structure the polynucleotides set forth herein. The methods can be carried out on a sample from any source, e.g., cells, tissues, body fluids, blood, urine, stool, hair, egg, sperm, cerebral spinal fluid, biopy samples, serum, etc.

These methods can be implemented in many different ways. For example, "comparing the structure" steps include, but are not limited to, comparing restriction maps, nucleotide sequences, amino acid sequences, RFLPs, Dnase sites, DNA methylation fingerprints (e.g., U.S. Pat. No. 6,214,556), protein cleavage sites, molecular weights, electrophoretic mobilities, charges, ion mobility, etc., between standard and a test genes. The term "structure" can refer to any physical characteristics or configurations which can be used to distinguish between nucleic acids and polypeptides. The methods and instruments used to accomplish the comparing step depends upon the physical characteristics which are to be

5

10

15

20

25

-69-

compared. Thus, various techniques are contemplated, including, e.g., sequencing machines (both amino acid and polynucleotide), electrophoresis, mass spectrometer (U.S. Pat. Nos. 6,093,541, 6,002,127), liquid chromatography, HPLC, etc.

To carry out such methods, "all or part" of the gene or polypeptide can be compared. For example, if nucleotide sequencing is utilized, the entire gene can be sequenced, including promoter, introns, and exons, or only parts of it can be sequenced and compared, e.g., exon 1, exon 2, etc.

Mutagenesis

5

10

15

20

25

30

Mutated polynucleotide sequences of the present invention are useful for various purposes, e.g., to create mutations of the polypeptides they encode, to identify functional regions of genomic DNA, to produce probes for screening libraries, etc. Mutagenesis can be carried out routinely according to any effective method, e.g., oligonucleotide-directed (Smith, M., Ann. Rev. Genet. 19:423-463, 1985), degenerate oligonucleotide-directed (Hill et al., Method Enzymology, 155:558-568, 1987), region-specific (Myers et al., Science, 229:242-246, 1985; Derbyshire et al., Gene, 46:145, 1986; Ner et al., DNA, 7:127, 1988), linkerscanning (McKnight and Kingsbury, Science, 217:316-324, 1982), directed using PCR, recursive ensemble mutagenesis (Arkin and Yourvan, Proc. Natl. Acad. Sci., 89:7811-7815, 1992), random mutagenesis (e.g., U.S. Pat. Nos. 5,096,815; 5,198,346; and 5,223,409), sitedirected mutagenesis (e.g., Walder et al., Gene, 42:133, 1986; Bauer et al., Gene, 37:73, 1985; Craik, Bio Techniques, January 1985, 12-19; Smith et al., Genetic Engineering: Principles and Methods, Plenum Press, 1981), phage display (e.g., Lowman et al., Biochem. 30:10832-10837, 1991; Ladner et al., U.S. Pat. No. 5,223,409; Huse, WIPO Publication WO 92/06204), etc. Desired sequences can also be produced by the assembly of target sequences using mutually priming oligonucleotides (Uhlmann, Gene, 71:29-40, 1988). For directed mutagenesis methods, analysis of the three-dimensional structure of the polypeptide can be used to guide and facilitate making mutants which effect polypeptide activity. Sites of substrate-enzyme interaction or other biological activities can also be determined by analysis of crystal structure as determined by such techniques as nuclear magnetic resonance, crystallography or photoaffinity labeling. See, for example, de Vos et al., Science 255:306-312, 1992; Smith et al., J. Mol. Biol. 224:899-904, 1992; Wlodaver et al., FEBS Lett.

309:59-64, 1992.

5

10

15

20

25

30

In addition, libraries of genes and fragments thereof can be used for screening and selection of genes variants. For instance, a library of coding sequences can be generated by treating a double-stranded DNA with a nuclease under conditions where the nicking occurs, e.g., only once per molecule, denaturing the double-stranded DNA, renaturing it to for double-stranded DNA that can include sense/antisense pairs from different nicked products, removing single-stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting DNAs into an expression vector. By this method, expression libraries can be made comprising "mutagenized" tissue selective genes. The entire coding sequence or parts thereof can be used.

Polynucleotide expression, polypeptides produced thereby, and specific-binding partners thereto.

A polynucleotide according to the present invention can be expressed in a variety of different systems, in vitro and in vivo, according to the desired purpose. For example, a polynucleotide can be inserted into an expression vector, introduced into a desired host, and cultured under conditions effective to achieve expression of a polypeptide coded for by the polynucleotide, to search for specific binding partners. Effective conditions include any culture conditions which are suitable for achieving production of the polypeptide by the host cell, including effective temperatures, pH, medium, additives to the media in which the host cell is cultured (e.g., additives which amplify or induce expression such as butyrate, or methotrexate if the coding polynucleotide is adjacent to a dhfr gene), cycloheximide, cell densities, culture dishes, etc. A polynucleotide can be introduced into the cell by any effective method including, e.g., naked DNA, calcium phosphate precipitation, electroporation, injection, DEAE-Dextran mediated transfection, fusion with liposomes, association with agents which enhance its uptake into cells, viral transfection. A cell into which a polynucleotide of the present invention has been introduced is a transformed host cell. The polynucleotide can be extrachromosomal or integrated into a chromosome(s) of the host cell. It can be stable or transient. An expression vector is selected for its compatibility with the host cell. Host cells include, mammalian cells, e.g., COS, CV1, BHK, CHO, HeLa, LTK, NIH 3T3, insect cells, such as Sf9 (S. frugipeda) and Drosophila, bacteria, such as E.

-71-

coli, Streptococcus, bacillus, yeast, such as Sacharomyces, S. cerevisiae, fungal cells, plant cells, embryonic or adult stem cells (e.g., mammalian, such as mouse or human),

immune system cell lines, HH (ATCC CRL 2105), MOLT-4 (ATCC CRL 1582), MJ (ATCC CRL-8294), SK7 (ATCC HB-8584), SK8 (ATCC HB-8585), HM1 (HB-8586), H9 (ATCC HTB-176), HuT 78 (ATCC TIB-161), HuT 102 (ATCC TIB-162), Jurkat,

B-cell lines, B-cell precursor lines, NALM-36, B-cell and other lymphocyte lines immortalized with Epstein-Barr virus (transformed B lymphoblastoid), stromal cell lines, myelomas, HBM-Noda, WEHI231,

reticuloendothelial cells, endothelial cells, white blood cells, macrophages, antigenresenting cells, lymphocytes, GDM-1 (ATCC CRL-2627), THP-1 (ATCC TIB-202), HL-60 (ATCC CCL-240), and derivatives thereof, including primary and established cell lines thereof,

kidney cell lines, 293, G-402 (ATCC CRL-1440), ACHN (ATCC CRL-1611), Vero (ATCC CCL-81), 786-O (ATCC CRL-1932), 769-P (ATCC CRL-1933), CCD 1103 KIDTr (ATCC CRL-2304), CCD 1105 KIDTr (ATCC CRL-2305), Hs 835.T (ATCC CRL-7569), Hs 926.T (ATCC CRL-7678), Caki-1 (ATCC HTB-46), Caki-2 (ATCC HTB-47), SW 839 (ATCC HTB-49), LLC-MK2 (ATCC CCL-7), BHK-21 (ATCC CCL-10), MDCK, CV-1, (ATCC CRL-1573), KNRK (ATCC CRL-1569), NRK-49F (ATCC CRL-1570), A-704 (ATCC HTB-45), etc., established and primary kidney cells,

pancreas cell lines, , insulinoma cell lines, INS-H1, MIN6N8, RIN 1046-38, RIN-5AH, RIN-A12, RINm5F, capan-1, capan-2, MIA PaCa-2 (ATCC CRL-1420), PANC-1 (ATCC CRL-1469), AsPC-1 (ATCC CRL-1682), SU-86.86 (ATCC CRL-1837), CFPAC-1 (ATCC CRL-1918), HPAF-II (ATCC CRL-1937), TGP61 (ATCC CRL-2135) and other TGP lines, SW 1990 (ATCC CRL-2172), Mpanc-96 (ATCC CRL-2380), MS1 VEGF (ATCC CRL-2460), Beta-TC-6 (ATCC CRL-11506), LTPA (ATCC CRL-2389), 266-6 (ATCC CRL-2151), MS1 (ATCC CRL-2779), SVR (ATCC CRL-2280), NIT-2 (ATCC CRL-2364), alphaTC1 Clone 9 (ATCC CRL-2350), ATCC CRL-1492, BxPC-3 (ATCC CRL-1687), HPAC (ATCC CRL-2119), U.S. Pat. Nos. 6,110743, 5,928,942, 5,888,816, 5,888,705, and 5,723,333, etc., established and primary pancreas cells (e.g., according to Hellerstrom et al., *Diabetes*, 28:769-76, 1979),

5

10

15

20

25

5

10

15

20

25

30

retinal cell lines, RF/6A (CRL 1780), ARPE-19 (CRL-2302), ARPE-19/HPV-16 (CRL-2502), Y79 (HTB-18), WERI-Rb-1 (HTB-169), RPE-J (CRL-2240), SO-Rb50 (retinoblastoma cell line), RBL, HER-Xho1-CC2, WERI-Rb24 (Sery et al., *J. Pediatr. Ophthalmol. Strabismus*, 4:212-217, 1990), WERI-Rb27 (Sery et al., *J. Pediatr. Ophthalmol. Strabismus*, 4:212-217, 1990), HXO-Rb44, fetal retina cells, retinoblastoma cells, choroidal endothelial cells (e.g., Chor 55), etc., established and primary retinal cells (For other cell lines and methods thereof, see, also, Griege et al, *Differentiation*, 45:250-7, 1990; Bernstein et al., *Invest. Ophthalmol. Vis. Sci.*, 35:3931-3937, 1994; Howes et al., *Invest. Ophthalmol. Vis. Sci.*, 35:342-351, 1994).

Expression control sequences are similarly selected for host compatibility and a desired purpose, e.g., high copy number, high amounts, induction, amplification, controlled expression. Other sequences which can be employed include enhancers such as from SV40, CMV, RSV, inducible promoters, cell-type specific elements, or sequences which allow selective or specific cell expression. Promoters that can be used to drive its expression, include, e.g., the endogenous promoter, MMTV, SV40, trp, lac, tac, or T7 promoters for bacterial hosts; or alpha factor, alcohol oxidase, or PGH promoters for yeast. RNA promoters can be used to produced RNA transcripts, such as T7 or SP6. See, e.g., Melton et al., *Polynucleotide Res.*, 12(18):7035-7056, 1984; Dunn and Studier. *J. Mol. Bio.*, 166:477-435, 1984; U.S. Pat. No. 5,891,636; Studier et al., *Gene Expression Technology, Methods in Enzymology*, 85:60-89, 1987. In addition, as discussed above, translational signals (including in-frame insertions) can be included.

When a polynucleotide is expressed as a heterologous gene in a transfected cell line, the gene is introduced into a cell as described above, under effective conditions in which the gene is expressed. The term "heterologous" means that the gene has been introduced into the cell line by the "hand-of-man." Introduction of a gene into a cell line is discussed above. The transfected (or transformed) cell expressing the gene can be lysed or the cell line can be used intact.

For expression and other purposes, a polynucleotide can contain codons found in a naturally-occurring gene, transcript, or cDNA, for example, e.g., as set forth in herein or it can contain degenerate codons coding for the same amino acid sequences. For instance,

-73-

it may be desirable to change the codons in the sequence to optimize the sequence for expression in a desired host. See, e.g., U.S. Pat. Nos. 5,567,600 and 5,567,862.

A polypeptide according to the present invention can be recovered from natural sources, transformed host cells (culture medium or cells) according to the usual methods, including, detergent extraction (e.g., non-ionic detergent, Triton X-100, CHAPS, octylglucoside, Igepal CA-630), ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, hydroxyapatite chromatography, lectin chromatography, gel electrophoresis. Protein refolding steps can be used, as necessary, in completing the configuration of the mature protein. Finally, high performance liquid chromatography (HPLC) can be employed for purification steps. Another approach is express the polypeptide recombinantly with an affinity tag (Flag epitope, HA epitope, myc epitope, 6xHis, maltose binding protein, chitinase, etc) and then purify by anti-tag antibody-conjugated affinity chromatography.

The present invention also relates to specific-binding partners. These include antibodies which are specific for polypeptides encoded by polynucleotides of the present invention, as well as other binding-partners which interact with polynucleotides and polypeptides of the present invention. Protein-protein interactions between polypeptides and binding partners can be identified using any suitable methods, e.g., protein binding assays (e.g., filtration assays, chromatography, etc.), yeast two-hybrid system (Fields and Song, *Nature*, 340: 245-247, 1989), protein arrays, gel-shift assays, FRET (fluorescence resonance energy transfer) assays, etc. Nucleic acid interactions (e.g., protein-DNA or protein-RNA) can be assessed using gel-shift assays, e.g., as carried out in U.S. Pat. No. 6,333,407 and 5,789,538.

Antibodies, e.g., polyclonal, monoclonal, recombinant, chimeric, humanized, single-chain, Fab, and fragments thereof, can be prepared according to any desired method. Antibodies, and immune responses, can also be generated by administering naked DNA See, e.g., U.S. Pat. Nos. 5,703,055; 5,589,466; 5,580,859. Antibodies can be used from any source, including, goat, rabbit, mouse, chicken (e.g., IgY; see, Duan, W0/029444 for methods of making antibodies in avian hosts, and harvesting the antibodies from the eggs). An antibody specific for a polypeptide means that the antibody recognizes a defined sequence of

5

10

15

20

25

5

10

15

20

25

amino acids within or including the polypeptide. Other specific binding partners include, e.g., aptamers and PNA. Antibodies can be prepared against specific epitopes or domains.

Antibodies can also be humanized, e.g., where they are to be used therapeutically. Methods for obtaining human antibodies, e.g., from transgenic mice are described, e.g., in Green et al., Nature Genet. 7:13 (1994); Lonberg et al., Nature 368:856 (1994); and Taylor et al., Int. Immunol. 6:579 (1994). Antibody fragments of the present invention can be prepared by any suitable method, Fab and Fc fragments. sinbgle-chain antibodies can also be used. Another form of an antibody fragment is a peptide coding for a single complementarity-determining region (CDR). CDR peptides ("minimal recognition units") can be obtained by constructing genes encoding the CDR of an antibody of interest.

The term "antibody" as used herein includes intact molecules as well as fragments thereof, such as Fab, F(ab')2, and Fv which are capable of binding to an epitopic determinant present in Bin1 polypeptide. Such antibody fragments retain some ability to selectively bind with its antigen or receptor. The term "epitope" refers to an antigenic determinant on an antigen to which the paratope of an antibody binds. Epitopic determinants usually consist of chemically active surface groupings of molecules such as amino acids or sugar side chains and usually have specific three dimensional structural characteristics, as well as specific charge characteristics. Antibodies can be prepared against specific epitopes or polypeptide domains.

Antibodies which bind to polypeptides of the present invention can be prepared using an intact polypeptide or fragments containing small peptides of interest as the immunizing antigen. For example, it may be desirable to produce antibodies that specifically bind to the N- or C-terminal domains of the tissue selective polypeptides of the present invention. The polypeptide or peptide used to immunize an animal which is derived from translated cDNA or chemically synthesized which can be conjugated to a carrier protein, if desired. Such commonly used carriers which are chemically coupled to the immunizing peptide include keyhole limpet hemocyanin (KLH), thyroglobulin, bovine serum albumin (BSA), and tetanus toxoid.

30 Methods of detecting polypeptides

-75-

Polypeptides coded for by genes of the present invention can be detected, visualized, determined, quantitated, etc. according to any effective method. useful methods include, e.g., but are not limited to, immunoassays, RIA (radioimmunassay), ELISA, (enzyme-linked-immunosorbent assay), immunoflourescence, flow cytometry, histology, electron microscopy, light microscopy, in situ assays, immunoprecipitation, Western blot, etc.

Immunoassays may be carried in liquid or on biological support. For instance, a sample (e.g., blood, serum, stool, urine, cells, tissue, cerebral spinal fluid, body fluids, etc.) can be brought in contact with and immobilized onto a solid phase support or carrier such as nitrocellulose, or other solid support that is capable of immobilizing cells, cell particles or soluble proteins. The support may then be washed with suitable buffers followed by treatment with the detectably labeled specific antibody. The solid phase support can then be washed with a buffer a second time to remove unbound antibody. The amount of bound label on solid support may then be detected by conventional means.

A "solid phase support or carrier" includes any support capable of binding an antigen, antibody, or other specific binding partner. Supports or carriers include glass, polystyrene, polypropylene, polyethylene, dextran, nylon, amylases, natural and modified celluloses, polyacrylamides, and magnetite. A support material can have any structural or physical configuration. Thus, the support configuration may be spherical, as in a bead, or cylindrical, as in the inside surface of a test tube, or the external surface of a rod. Alternatively, the surface may be flat such as a sheet, test strip, etc. Preferred supports include polystyrene beads

One of the many ways in which gene peptide-specific antibody can be detectably labeled is by linking it to an enzyme and using it in an enzyme immunoassay (EIA). See, e.g., Voller, A., "The Enzyme Linked Immunosorbent Assay (ELISA)," 1978, Diagnostic Horizons 2, 1-7, Microbiological Associates Quarterly Publication, Walkersville, Md.); Voller, A. et al., 1978, J. Clin. Pathol. 31, 507-520; Butler, J. E., 1981, Meth. Enzymol. 73, 482-523; Maggio, E. (ed.), 1980, Enzyme Immunoassay, CRC Press, Boca Raton, Fla.. The enzyme which is bound to the antibody will react with an appropriate substrate, preferably a chromogenic substrate, in such a manner as to produce a chemical moiety that can be detected, for example, by spectrophotometric, fluorimetric or by visual means. Enzymes that can be used to detectably label the antibody include, but are not limited to, malate

5

10

15

20

25

-76-

dehydrogenase, staphylococcal nuclease, delta-5-steroid isomerase, yeast alcohol dehydrogenase, .alpha.-glycerophosphate, dehydrogenase, triose phosphate isomerase, horseradish peroxidase, alkaline phosphatase, asparaginase, glucose oxidase, .beta.-galactosidase, ribonuclease, urease, catalase, glucose-6-phosphate dehydrogenase, glucoamylase and acetylcholinesterase. The detection can be accomplished by colorimetric methods that employ a chromogenic substrate for the enzyme. Detection may also be accomplished by visual comparison of the extent of enzymatic reaction of a substrate in comparison with similarly prepared standards.

Detection may also be accomplished using any of a variety of other immunoassays. For example, by radioactively labeling the antibodies or antibody fragments, it is possible to detect peptides through the use of a radioimmunoassay (RIA). See, e.g., Weintraub, B., Principles of Radioimmunoassays, Seventh Training Course on Radioligand Assay Techniques, The Endocrine Society, March, 1986. The radioactive isotope can be detected by such means as the use of a gamma counter or a scintillation counter or by autoradiography.

It is also possible to label the antibody with a fluorescent compound. When the fluorescently labeled antibody is exposed to light of the proper wave length, its presence can then be detected due to fluorescence. Among the most commonly used fluorescent labeling compounds are fluorescein isothiocyanate, rhodamine, phycoerythrin, phycocyanin, allophycocyanin, o-phthaldehyde and fluorescamine. The antibody can also be detectably labeled using fluorescence emitting metals such as those in the lanthanide series. These metals can be attached to the antibody using such metal chelating groups as diethylenetriaminepentacetic acid (DTPA) or ethylenediaminetetraacetic acid (EDTA).

The antibody also can be detectably labeled by coupling it to a chemiluminescent compound. The presence of the chemiluminescent-tagged antibody is then determined by detecting the presence of luminescence that arises during the course of a chemical reaction. Examples of useful chemiluminescent labeling compounds are luminol, isoluminol, theromatic acridinium ester, imidazole, acridinium salt and oxalate ester.

Likewise, a bioluminescent compound may be used to label the antibody of the present invention. Bioluminescence is a type of chemiluminescence found in biological systems in which a catalytic protein increases the efficiency of the chemiluminescent reaction. The presence of a bioluminescent protein is determined by detecting the presence of

5

10

15

20

25

-77-

luminescence. Important bioluminescent compounds for purposes of labeling are luciferin, luciferase and aequorin.

Diagnostic

5

10

15

20

25

30

The present invention also relates to methods and compositions for diagnosing a disorder, or determining susceptibility to a disorder, using polynucleotides, polypeptides, and specific-binding partners of the present invention to detect, assess, determine, etc., a tissue selective gene. In such methods, the gene can serve as a marker for the disorder, e.g., where the gene, when mutant, is a direct cause of the disorder; where the gene is affected by another gene(s) which is directly responsible for the disorder, e.g., when the gene is part of the same signaling pathway as the directly responsible gene; and, where the gene is chromosomally linked to the gene(s) directly responsible for the disorder, and segregates with it. Many other situations are possible. To detect, assess, determine, etc., a probe specific for the gene can be employed as described above and below. Any method of detecting and/or assessing the gene can be used, including detecting expression of the gene using polynucleotides, antibodies, or other specific-binding partners.

The phrase "diagnosing" indicates that it is determined whether the sample has the disorder. A "disorder" means, e.g., any abnormal condition as in a disease or malady. "Determining a subject's susceptibility to a disease or disorder" indicates that the subject is assessed for whether s/he is predisposed to get such a disease or disorder, where the predisposition is indicated by abnormal expression of the gene (e.g., gene mutation, gene expression pattern is not normal, etc.). Predisposition or susceptibility to a disease may result when a such disease is influenced by epigenetic, environmental, etc., factors. Diagnosing includes prenatal screening where samples from the fetus or embryo (e.g., via amniocentesis or CV sampling) are analyzed for the expression of the gene.

By the phrase "assessing expression of a gene or polynucleotide," it is meant that the functional status of the gene is evaluated. This includes, but is not limited to, measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene. Thus, the term "assessing expression" includes evaluating the all aspects of the transcriptional and translational machinery of the gene. For

-78-

instance, if a promoter defect causes, or is suspected of causing, the disorder, then a sample can be evaluated (i.e., "assessed") by looking (e.g., sequencing or restriction mapping) at the promoter sequence in the gene, by detecting transcription products (e.g., RNA), by detecting translation product (e.g., polypeptide). Any measure of whether the gene is functional can be used, including, polypeptide, polynucleotide, and functional assays for the gene's biological activity.

In making the assessment, it can be useful to compare the results to a normal gene, e.g., a gene which is not associated with the disorder. The nature of the comparison can be determined routinely, depending upon how the assessing is accomplished. If, for example, the mRNA levels of a sample is detected, then the mRNA levels of a normal can serve as a comparison, or a gene which is known not to be affected by the disorder. Methods of detecting mRNA are well known, and discussed above, e.g., but not limited to, Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, etc. Similarly, if polypeptide production is used to evaluate the gene, then the polypeptide in a normal tissue sample can be used as a comparison, or, polypeptide from a different gene whose expression is known not to be affected by the disorder. These are only examples of how such a method could be carried out.

The genes and polypeptides of the present invention can be used to identify, detect, stage, determine the presence of, prognosticate, treat, study, etc., diseases and conditions as mentioned above. The present invention relates to methods of identifying a genetic basis for a disease or disease-susceptibility, comprising, e.g., determining the association of a disease or disease-susceptibility with a gene of the present invention. An association between a disease or disease-susceptibility and nucleotide sequence includes, e.g., establishing (or finding) a correlation (or relationship) between a DNA marker (e.g., gene, VNTR, polymorphism, EST, etc.) and a particular disease state. Once a relationship is identified, the DNA marker can be utilized in diagnostic tests and as a drug target. Any region of the gene can be used as a source of the DNA marker, exons, introns, intergenic regions, etc.

Human linkage maps can be constructed to establish a relationship between a gene and a disease or condition. Typically, polymorphic molecular markers (e.g., STRP's, SNP's, RFLP's, VNTR's) are identified within the region, linkage and map distance between the markers is then established, and then linkage is established between phenotype and the

5

10

15

20

25

-79-

various individual molecular markers. Maps can be produced for an individual family, selected populations, patient populations, etc. In general, these methods involve identifying a marker associated with the disease (e.g., identifying a polymorphism in a family which is linked to the disease) and then analyzing the surrounding DNA to identity the gene responsible for the phenotype. See, e.g., Kruglyak et al., Am. J. Hum. Genet., 58, 1347-1363, 1996; Matise et al., Nat. Genet., 6(4):384-90, 1994.

Assessing the effects of the rapeutic and preventative interventions (e.g., administration of a drug, chemotherapy, radiation, etc.) on disorders is a major effort in drug discovery, clinical medicine, and pharmacogenomics. The evaluation of therapeutic and preventative measures, whether experimental or already in clinical use, has broad applicability, e.g., in clinical trials, for monitoring the status of a patient, for analyzing and assessing animal models, and in any scenario involving disease treatment and prevention. Analyzing the expression profiles of polynucleotides of the present invention can be utilized as a parameter by which interventions are judged and measured. Treatment of a disorder can change the expression profile in some manner which is prognostic or indicative of the drug's effect on it. Changes in the profile can indicate, e.g., drug toxicity, return to a normal level, etc. Accordingly, the present invention also relates to methods of monitoring or assessing a therapeutic or preventative measure (e.g., chemotherapy, radiation, anti-neoplastic drugs, antibodies, etc.) in a subject having a disorder, or, susceptible to such a disorder, comprising, e.g., detecting the expression levels of one or more tissue selective genes. A subject can be a cell-based assay system, non-human animal model, human patient, etc. Detecting can be accomplished as described for the methods above and below. By "therapeutic or preventative intervention," it is meant, e.g., a drug administered to a patient, surgery, radiation, chemotherapy, and other measures taken to prevent, treat, or diagnose a disorder.

The present invention also relates to methods of using binding partners, such as antibodies, to deliver active agents to the tissue (e.g., kidney or pancreas or an immune cells) for a variety of different purposes, including, e.g., for diagnostic, therapeutic, and research purposes. Methods can involve delivering or administering an active agent to the tissue, comprising, e.g., administering to a subject in need thereof, an effective amount of an active agent coupled to a binding partner specific for a tissue selective polypeptide, wherein said binding partner is effective to deliver said active agent specifically to the target tissue.

5

10

15

20

25

-80-

Any type of active agent can be used in combination with it, including, therapeutic, cytotoxic, cytostatic, chemotherapeutic, anti-neoplastic, anti-proliferative, anti-biotic, etc., agents. A chemotherapeutic agent can be, e.g., DNA-interactive agent, alkylating agent, antimetabolite, tubulin-interactive agent, hormonal agent, hydroxyurea, Cisplatin, Cyclophosphamide, Altretamine, Bleomycin, Dactinomycin, Doxorubicin, Etoposide, Teniposide, paclitaxel, cytoxan, 2-methoxy-carbonyl-amino-benzimidazole, Plicamycin, Methotrexate, Fluorouracil, Fluorodeoxyuridin, CB3717, Azacitidine, Floxuridine, Mercapyopurine, 6-Thioguanine, Pentostatin, Cytarabine, Fludarabine, etc. Agents can also be contrast agents useful in imaging technology, e.g., X-ray, CT, CAT, MRI, ultrasound, PET, SPECT, and scintographic.

An active agent can be associated in any manner with a binding partner which is effective to achieve its delivery specifically to the target. Specific delivery or targeting indicates that the agent is provided to the tissue, without being substantially provided to other tissues. This is useful especially where an agent is toxic, and specific targeting to the tissue enables the majority of the toxicity to be aimed at the tissue, with as small as possible effect on other tissues in the body. The association of the active agent and the binding partner ("coupling") can be direct, e.g., through chemical bonds between the binding partner and the agent, or, via a linking agent, or the association can be less direct, e.g., where the active agent is in a liposome, or other carrier, and the binding partner is associated with the liposome surface. In such case, the binding partner can be oriented in such a way that it is able to bind to tissue selective polypeptide, e.g., exposed on the cell surface. Methods for delivery of DNA via a cell-surface receptor is described, e.g., in U.S. Pat. No. 6,339,139.

Identifying agent methods

The present invention also relates to methods of identifying agents, and the agents themselves, which modulate tissue selective genes. These agents can be used to modulate the biological activity of the polypeptide encoded for the gene, or the gene, itself. Agents which regulate the gene or its product are useful in variety of different environments, including as medicinal agents to treat or prevent disorders associated with genes and as research reagents to modify the function of tissues and cell.

5

10

15

20

25

-81-

Methods of identifying agents generally comprise steps in which an agent is placed in contact with the gene, its transcription product, its translation product, or other target, and then a determination is performed to assess whether the agent "modulates" the target. The specific method utilized will depend upon a number of factors, including, e.g., the target (i.e., is it the gene or polypeptide encoded by it), the environment (e.g., in vitro or in vivo), the composition of the agent, etc.

For modulating the expression of tissue selective genes, a method can comprise, in any effective order, one or more of the following steps, e.g., contacting a gene (e.g., in a cell population) with a test agent under conditions effective for said test agent to modulate the expression of tissue selective genes, and determining whether said test agent modulates said genes. An agent can modulate expression of a tissue selective gene at any level, including transcription (e.g., by modulating the promoter), translation, and/or perdurance of the nucleic acid (e.g., degradation, stability, etc.) in the cell.

For modulating the biological activity of polypeptides, a method can comprise, in any effective order, one or more of the following steps, e.g., contacting a polypeptide (e.g., in a cell, lysate, or isolated) with a test agent under conditions effective for said test agent to modulate the biological activity of said polypeptide, and determining whether said test agent modulates said biological activity.

Contacting a gene or polypeptide with the test agent can be accomplished by any suitable method and/or means that places the agent in a position to functionally control expression or biological activity. Functional control indicates that the agent can exert its physiological effect through whatever mechanism it works. The choice of the method and/or means can depend upon the nature of the agent and the condition and type of environment in which the gene or polypeptide is presented, e.g., lysate, isolated, or in a cell population (such as, in vivo, in vitro, organ explants, etc.). For instance, if the cell population is an in vitro cell culture, the agent can be contacted with the cells by adding it directly into the culture medium. If the agent cannot dissolve readily in an aqueous medium, it can be incorporated into liposomes, or another lipophilic carrier, and then administered to the cell culture. Contact can also be facilitated by incorporation of agent with carriers and delivery molecules and complexes, by injection, by infusion, etc.

Agents can be directed to, or targeted to, any part of the polypeptide which is

5

10

15

20

25

-82-

effective for modulating it. For example, agents, such as antibodies and small molecules, can be targeted to cell-surface, exposed, extracellular, ligand binding, functional, etc., domains of the polypeptide. Agents can also be directed to intracellular regions and domains, e.g., regions where the polypeptide couples or interacts with intracellular or intramembrane binding partners.

After the agent has been administered in such a way that it can gain access, it can be determined whether the test agent modulates expression or biological activity. Modulation can be of any type, quality, or quantity, e.g., increase, facilitate, enhance, up-regulate, stimulate, activate, amplify, augment, induce, decrease, down-regulate, diminish, lessen, reduce, etc. The modulatory quantity can also encompass any value, e.g., 1%, 5%, 10%, 50%, 75%, 1-fold, 2-fold, 5-fold, 10-fold, 100-fold, etc. To modulate expression means, e.g., that the test agent has an effect on its expression, e.g., to effect the amount of transcription, to effect RNA splicing, to effect translation of the RNA into polypeptide, to effect RNA or polypeptide stability, to effect polyadenylation or other processing of the RNA, to effect post-transcriptional or post-translational processing, etc. To modulate biological activity means, e.g., that a functional activity of the polypeptide is changed in comparison to its normal activity in the absence of the agent. This effect includes, increase, decrease, block, inhibit, enhance, etc.

A test agent can be of any molecular composition, e.g., chemical compounds, biomolecules, such as polypeptides, lipids, nucleic acids (e.g., antisense), carbohydrates, antibodies, ribozymes, double-stranded RNA, aptamers, etc. For example, if a polypeptide to be modulated is a cell-surface molecule, a test agent can be an antibody that specifically recognizes it and, e.g., causes the polypeptide to be internalized, leading to its down regulation on the surface of the cell. Such an effect does not have to be permanent, but can require the presence of the antibody to continue the down-regulatory effect. Antibodies can also be used to modulate the biological activity of a polypeptide in a lysate or other cell-free form.

Additional cell-based test systems suitable for the analysis of GPCR polypeptides are summarized in Marchese et al. (1999, Trends in Pharmacol. Sci. 20: 370-375) and comprise so-called "ligand screening assays." For example in yeast cells the pheromon receptor can be replaced by a GPCR according to the invention. The effect of test substances on the receptor

5

10

15

20

25

· -83-

can be determined upon modulation of histidine synthesis, i.e. by growing in histidine-free medium. In addition using cells transfected with nucleic acids according to the invention it can be analyzed whether test substances mediate translocation of a detectable arrestins, for example of a arrestin-GFP-fusion protein. Moreover, it can be analyzed whether test substances mediate GPCR-mediated dispersion or aggregation of Xenopus laevis melanophores. Another test system utilizes the universal adapter G-protein G alphal6, which mobilizes Ca.sup.2+. Other screening test systems are described in Lemer et al., supra; WO96/41169; U.S. Pat. No. 5,482,835; WO99/06535; EP 0 939 902; WO99/66326; WO98/34948; EP 0 863 214; U.S. Pat. No. 5,882,944 and U.S. Pat. No. 5,891,641.

10 Therapeutics

5

15

20

25

30

Selective polynucleotides, polypeptides, and specific-binding partners thereto, can be utilized in therapeutic applications, especially to treat diseases and conditions described herein. Useful methods include, but are not limited to, immunotherapy (e.g., using specific-binding partners to polypeptides), vaccination (e.g., using a selective polypeptide or a naked DNA encoding such polypeptide), protein or polypeptide replacement therapy, gene therapy (e.g., germ-line correction, antisense), etc.

Various immunotherapeutic approaches can be used. For instance, unlabeled antibody that specifically recognizes a tissue-specific antigen can be used to stimulate the body to destroy or attack a cancer or other diseased tissue, to cause down-regulation, to produce complement-mediated lysis, to inhibit cell growth, etc., of target cells which display the antigen, e.g., analogously to how c-erbB-2 antibodies are used to treat breast cancer. In addition, antibody can be labeled or conjugated to enhance its deleterious effect, e.g., with radionuclides and other energy emitting entitities, toxins, such as ricin, exotoxin A (ETA), and diphtheria, cytotoxic or cytostatic agents, immunomodulators, chemotherapeutic agents, etc. See, e.g., U.S. Pat. No. 6,107,090.

An antibody or other specific-binding partner can be conjugated to a second molecule, such as a cytotoxic agent, and used for targeting the second molecule to a tissue-antigen positive cell (Vitetta, E. S. et al., 1993, Immunotoxin therapy, in DeVita, Jr., V. T. et al., eds, Cancer: Principles and Practice of Oncology, 4th ed., J. B. Lippincott Co., Philadelphia, 2624-2636). Examples of cytotoxic agents include, but are not limited to, antimetabolites, alkylating agents, anthracyclines, antibiotics, anti-mitotic agents, radioisotopes and

-84-

chemotherapeutic agents. Further examples of cytotoxic agents include, but are not limited to ricin, doxorubicin, daunorubicin, taxol, ethidium bromide, mitomycin, etoposide, tenoposide, vincristine, vinblastine, colchicine, dihydroxy anthracin dione, actinomycin D, 1-dehydrotestosterone, diptheria toxin, Pseudomonas exotoxin (PE) A, PE40, abrin, elongation factor-2 and glucocorticoid. Techniques for conjugating therapeutic agents to antibodies are well.

In addition to immunotherapy, polynucleotides and polypeptides can be used as targets for non-immunotherapeutic applications, e.g., using compounds which interfere with function, expression (e.g., antisense as a therapeutic agent), assembly, etc. RNA interference can be used in vitro and in vivo to silence a gene when its expression contributes to a disease (but also for other purposes, e.g., to identify the gene's function to change a developmental pathway of a cell, etc.). See, e.g., Sharp and Zamore, *Science*, 287:2431-2433, 2001; Grishok et al., *Science*, 287:2494, 2001.

Delivery of therapeutic agents can be achieved according to any effective method, including, liposomes, viruses, plasmid vectors, bacterial delivery systems, orally, systemically, etc. Therapeutic agents of the present invention can be administered in any form by any effective route, including, e.g., oral, parenteral, enteral, intraperitoneal, topical, transdermal (e.g., using any standard patch), intravenously, ophthalmic, nasally, local, non-oral, such as aerosal, inhalation, subcutaneous, intramuscular, buccal, sublingual, rectal, vaginal, intra-arterial, and intrathecal, etc. They can be administered alone, or in combination with any ingredient(s), active or inactive.

In addition to therapeutics, per se, the present invention also relates to methods of treating a disease showing altered expression of a tissue selective gene, comprising, e.g., administering to a subject in need thereof a therapeutic agent which is effective for regulating expression of said gene and/or which is effective in treating said disease. The term "treating" is used conventionally, e.g., the management or care of a subject for the purpose of combating, alleviating, reducing, relieving, improving the condition of, etc., of a disease or disorder. By the phrase "altered expression," it is meant that the disease is associated with a mutation in the gene, or any modification to the gene (or corresponding product) which affects its normal function. Thus, expression refers to, e.g., transcription, translation, splicing, stability of the mRNA or protein product, activity of the gene product, differential

5

10

15

20

25

expression, etc.

Any agent which "treats" the disease can be used. Such an agent can be one which regulates the expression of a tissue selective gene. Expression refers to the same acts already mentioned, e.g. transcription, translation, splicing, stability of the mRNA or protein product, activity of the gene product, differential expression, etc. For instance, if the condition was a result of a complete deficiency of the gene product, administration of gene product to a patient would be said to treat the disease and regulate the gene's expression. Many other possible situations are possible, e.g., where the gene is aberrantly expressed, and the therapeutic agent regulates the aberrant expression by restoring its normal expression pattern.

10

15

20

25

30

5

Antisense

Antisense polynucleotide (e.g., RNA) can also be prepared from a polynucleotide according to the present invention. Antisense polynucleotide can be used in various ways, such as to regulate or modulate expression of the polypeptides they encode, e.g., inhibit their expression, for in situ hybridization, for therapeutic purposes, for making targeted mutations (in vivo, triplex, etc.) etc. For guidance on administering and designing anti-sense, see, e.g., U.S. Pat. Nos. 6,200,960, 6,200,807, 6,197,584, 6,190,869, 6,190,661, 6,187,587, 6,168,950, 6,153,595, 6,150,162, 6,133,246, 6,117,847, 6,096,722, 6,087,343, 6,040,296, 6,005,095, 5,998,383, 5,994,230, 5,891,725, 5,885,970, and 5,840,708. An antisense polynucleotides can be operably linked to an expression control sequence. A total length of about 35 bp can be used in cell culture with cationic liposomes to facilitate cellular uptake, but for *in vivo* use, preferably shorter oligonucleotides are administered, e.g. 25 nucleotides.

Antisense polynucleotides can comprise modified, nonnaturally-occurring nucleotides and linkages between the nucleotides (e.g., modification of the phosphate-sugar backbone; methyl phosphonate, phosphorothioate, or phosphorodithioate linkages; and 2'-O-methyl ribose sugar units), e.g., to enhance in vivo or in vitro stability, to confer nuclease resistance, to modulate uptake, to modulate cellular distribution and compartmentalization, etc. Any effective nucleotide or modification can be used, including those already mentioned, as known in the art, etc., e.g., disclosed in U.S. Pat. Nos. 6,133,438; 6,127,533; 6,124,445; 6,121,437; 5,218,103 (e.g., nucleoside thiophosphoramidites); 4,973,679; Sproat et al., "2'-O-Methyloligoribonucleotides: synthesis and applications," Oligonucleotides and Analogs A

Practical Approach, Eckstein (ed.), IRL Press, Oxford, 1991, 49-86; Iribarren et al., "2'O-Alkyl Oligoribonucleotides as Antisense Probes," Proc. Natl. Acad. Sci. USA, 1990, 87, 7747-7751; Cotton et al., "2'-O-methyl, 2'-O-ethyl oligoribonucleotides and phosphorothioate oligodeoxyribonucleotides as inhibitors of the in vitro U7 snRNP-dependent mRNA processing event," Nucl. Acids Res., 1991, 19, 2629-2635.

Arrays

5

10

15

20

30

The present invention also relates to an ordered array of polynucleotide probes and specific-binding partners (e.g., antibodies) for detecting the expression of tissue selective genes or polypeptides encoded thereby, in a sample, comprising, one or more polynucleotide probes or specific binding partners associated with a solid support or in separate receptacles, wherein each probe is specific for a tissue selective gene or a specific-binding partner which is specific for a polypeptide.

The phrase "ordered array" indicates that the probes are arranged in an identifiable or position-addressable pattern, e.g., such as the arrays disclosed in U.S. Pat. Nos. 6,156,501, 6,077,673, 6,054,270, 5,723,320, 5,700,637, WO09919711, WO00023803. The probes are associated with the solid support in any effective way. For instance, the probes can be bound to the solid support, either by polymerizing the probes on the substrate, or by attaching a probe to the substrate. Association can be, covalent, electrostatic, noncovalent, hydrophobic, hydrophilic, noncovalent, coordination, adsorbed, absorbed, polar, etc. When fibers or hollow filaments are utilized for the array, the probes can fill the hollow orifice, be absorbed into the solid filament, be attached to the surface of the orifice, etc. Probes can be of any effective size, sequence identity, composition, etc., as already discussed.

25 Transgenic animals

The present invention also relates to transgenic animals comprising tissue selective genes, and homologs thereof. (Methods of making transgenic animals, and associated recombinant technology, can be accomplished conventionally, e.g., as described in *Transgenic Animal Technology*, Pinkert et al., 2nd Edition, Academic Press, 2002.) Such genes, as discussed in more detail below, include, but are not limited to, functionally-disrupted genes, mutated genes, ectopically or selectively-expressed genes, inducible or

-87-

regulatable genes, etc. These transgenic animals can be produced according to any suitable technique or method, including homologous recombination, mutagenesis (e.g., ENU, Rathkolb et al., *Exp. Physiol.*, 85(6):635-644, 2000), and the tetracycline-regulated gene expression system (e.g., U.S. Pat. No. 6,242,667). The term "gene" as used herein includes any part of a gene, i.e., regulatory sequences, promoters, enhancers, exons, introns, coding sequences, etc. The nucleic acid present in the construct or transgene can be naturally-occurring wild-type, polymorphic, or mutated. Where the animal is a non-human animal, its homolog can be used instead. Transgenic animals can have structural and/or functional defects in any of the tissues described herein, e.g., pancreas, kidney, retina, and immune cells, as well as having or being susceptible to any of the associated disorders or diseases mentioned herein.

Along these lines, polynucleotides of the present invention can be used to create transgenic animals, e.g. a non-human animal, comprising at least one cell whose genome comprises a functional disruption of one or tissue selective genes, or homologs thereof (e.g., a mouse homolog when a mouse is used). By the phrases "functional disruption" or "functionally disrupted," it is meant that the gene does not express a biologically-active product. It can be substantially deficient in at least one functional activity coded for by the gene. Expression of a polypeptide can be substantially absent, i.e., essentially undetectable amounts are made. However, polypeptide can also be made, but which is deficient in activity, e.g., where only an amino-terminal portion of the gene product is produced.

The transgenic animal can comprise one or more cells. When substantially all its cells contain the engineered gene, it can be referred to as a transgenic animal "whose genome comprises" the engineered gene. This indicates that the endogenous gene loci of the animal has been modified and substantially all cells contain such modification.

Functional disruption of the gene can be accomplished in any effective way, including, e.g., introduction of a stop codon into any part of the coding sequence such that the resulting polypeptide is biologically inactive (e.g., because it lacks a catalytic domain, a ligand binding domain, etc.), introduction of a mutation into a promoter or other regulatory sequence that is effective to turn it off, or reduce transcription of the gene, insertion of an exogenous sequence into the gene which inactivates it (e.g., which disrupts the production of a biologically-active polypeptide or which disrupts the promoter or other transcriptional

5

10

15

20

25

-88-

machinery), deletion of sequences from the gene (or homolog thereof), etc. Examples of transgenic animals having functionally disrupted genes are well known, e.g., as described in U.S. Pat. Nos. 6,239,326, 6,225,525, 6,207,878, 6,194,633, 6,187,992, 6,180,849, 6,177,610, 6,100,445, 6,087,555, 6,080,910, 6,069,297, 6,060,642, 6,028,244, 6,013,858, 5,981,830, 5,866,760, 5,859,314, 5,850,004, 5,817,912, 5,789,654, 5,777,195, and 5,569,824. A transgenic animal which comprises the functional disruption can also be referred to as a "knock-out" animal, since the biological activity of its gene has been "knocked-out." Knock-outs can be homozygous or heterozygous.

For creating functionally disrupted genes, and other gene mutations, homologous recombination technology is of special interest since it allows specific regions of the genome to be targeted. Using homologous recombination methods, genes can be specificallyinactivated, specific mutations can be introduced, and exogenous sequences can be introduced at specific sites. These methods are well known in the art, e.g., as described in the patents above. See, also, Robertson, Biol. Reproduc., 44(2):238-245, 1991. Generally, the genetic engineering is performed in an embryonic stem (ES) cell, or other pluripotent cell line (e.g., adult stem cells, EG cells), and that genetically-modified cell (or nucleus) is used to create a whole organism. Nuclear transfer can be used in combination with homologous recombination technologies. For example, a gene locus can be disrupted in mouse ES cells using a positive-negative selection method (e.g., Mansour et al., Nature, 336:348-352, 1988). In this method, a targeting vector can be constructed which comprises a part of the gene to be targeted. A selectable marker, such as neomycin resistance genes, can be inserted into a an exon present in the targeting vector, disrupting it. When the vector recombines with the ES cell genome, it disrupts the function of the gene. The presence in the cell of the vector can be determined by expression of neomycin resistance. See, e.g., U.S. Pat. No. 6,239,326. Cells having at least one functionally disrupted gene can be used to make chimeric and germline animals, e.g., animals having somatic and/or germ cells comprising the engineered gene. Homozygous knock-out animals can be obtained from breeding heterozygous knockout animals. See, e.g., U.S. Pat. No. 6,225,525.

The present invention also relates to non-human, transgenic animal whose genome comprises recombinant tissue selective nuccleic acid (and homologs thereof) operatively linked to an expression control sequence effective to express said coding sequence in a target

5

10

15

20

25

WO 03/089583 PCT/US03/11497
-89-

tissue. Such a transgenic animal can also be referred to as a "knock-in" animal since an exogenous gene has been introduced, stably, into its genome. "Operable linkage" has the meaning used through the specification, i.e., placed in a functional relationship with another nucleic acid. When a gene is operably linked to an expression control sequence, as explained above, it indicates that the gene (e.g., coding sequence) is joined to the expression control sequence (e.g., promoter) in such a way that facilitates transcription and translation of the coding sequence. As described above, the phrase "genome" indicates that the genome of the cell has been modified. In this case, the recombinant gene has been stably integrated into the genome of the animal. The nucleic acid (e.g., a coding sequence) in operable linkage with the expression control sequence can also be referred to as a construct or transgene.

Any expression control sequence can be used depending on the purpose. For instance, if selective expression is desired, then expression control sequences which limit its expression can be selected. These include, e.g., tissue or cell-specific promoters, introns, enhancers, etc. For various methods of cell and tissue-specific expression, see, e.g., U.S. Pat. Nos. 6,215,040, 6,210,736, and 6,153,427. These also include the endogenous promoter, i.e., the coding sequence can be operably linked to its own promoter. Inducible and regulatable promoters can also be utilized.

The present invention also relates to a transgenic animal which contains a functionally disrupted and a transgene stably integrated into the animals genome. Such an animal can be constructed using combinations any of the above- and below-mentioned methods. Such animals have any of the aforementioned uses, including permitting the knock-out of the normal gene and its replacement with a mutated gene. Such a transgene can be integrated at the endogenous gene locus so that the functional disruption and "knock-in" are carried out in the same step.

In addition to the methods mentioned above, transgenic animals can be prepared according to known methods, including, e.g., by pronuclear injection of recombinant genes into pronuclei of 1-cell embryos, incorporating an artificial yeast chromosome into embryonic stem cells, gene targeting methods, embryonic stem cell methodology, cloning methods, nuclear transfer methods. See, also, e.g., U.S. Patent Nos. 4,736,866; 4,873,191; 4,873,316; 5,082,779; 5,304,489; 5,174,986; 5,175,384; 5,175,385; 5,221,778; Gordon et al., Proc. Natl. Acad. Sci., 77:7380-7384, 1980; Palmiter et al., Cell, 41:343-345, 1985; Palmiter

5

10

15

20

25

et al., Ann. Rev. Genet., 20:465-499, 1986; Askew et al., Mol. Cell. Bio., 13:4115-4124, 1993; Games et al. Nature, 373:523-527, 1995; Valancius and Smithies, Mol. Cell. Bio., 11:1402-1408, 1991; Stacey et al., Mol. Cell. Bio., 14:1009-1016, 1994; Hasty et al., Nature, 350:243-246, 1995; Rubinstein et al., Nucl. Acid Res., 21:2613-2617,1993; Cibelli et al., 5 Science, 280:1256-1258, 1998. For guidance on recombinase excision systems, see, e.g., U.S. Pat. Nos. 5,626,159, 5,527,695, and 5,434,066. See also, Orban, P.C., et al., "Tissueand Site-Specific DNA Recombination in Transgenic Mice," Proc. Natl. Acad. Sci. USA, 89:6861-6865 (1992); O'Gorman, S., et al., "Recombinase-Mediated Gene Activation and Site-Specific Integration in Mammalian Cells," Science, 251:1351-1355 (1991); Sauer, B., et 10 al., "Cre-stimulated recombination at loxP-Containing DNA sequences placed into the mammalian genome," Polynucleotides Research, 17(1):147-161 (1989); Gagneten, S. et al. (1997) Nucl. Acids Res. 25:3326-3331; Xiao and Weaver (1997) Nucl. Acids Res. 25:2985-2991; Agah, R. et al. (1997) J. Clin. Invest. 100:169-179; Barlow, C. et al. (1997) Nucl. Acids Res. 25:2543-2545; Araki, K. et al. (1997) Nucl. Acids Res. 25:868-872; Mortensen, R. N. et al. (1992) Mol. Cell. Biol. 12:2391-2395 (G418 escalation method); Lakhlani, P. P. 15 et al. (1997) Proc. Natl. Acad. Sci. USA 94:9950-9955 ("hit and run"); Westphal and Leder (1997) Curr. Biol. 7:530-533 (transposon-generated "knock-out" and "knock-in"); Templeton, N. S. et al. (1997) Gene Ther. 4:700-709 (methods for efficient gene targeting, allowing for a high frequency of homologous recombination events, e.g., without selectable 20 markers); PCT International Publication WO 93/22443 (functionally-disrupted).

A polynucleotide according to the present invention can be introduced into any non-human animal, including a non-human mammal, mouse (Hogan et al., Manipulating the Mouse Embryo: A Laboratory Manual, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1986), pig (Hammer et al., Nature, 315:343-345, 1985), sheep (Hammer et al., Nature, 315:343-345, 1985), cattle, rat, or primate. See also, e.g., Church, 1987, Trends in Biotech. 5:13-19; Clark et al., Trends in Biotech. 5:20-24, 1987); and DePamphilis et al., BioTechniques, 6:662-680, 1988. Transgenic animals can be produced by the methods described in U.S. Pat. No. 5,994,618, and utilized for any of the utilities described therein.

30 Database

The present invention also relates to electronic forms of polynucleotides,

WO 03/089583 PCT/US03/11497 -91-

polypeptides, etc., of the present invention, including computer-readable medium (e.g., magnetic, optical, etc., stored in any suitable format, such as flat files or hierarchical files) which comprise such sequences, or fragments thereof, e-commerce-related means, etc. Along these lines, the present invention relates to methods of retrieving nucleic acid and/or polypeptide sequences from a computer-readable medium, comprising, one or more of the following steps in any effective order, e.g., selecting a cell or gene expression profile, e.g., a profile that specifies that said gene is differentially expressed in a tissue as described herein, and retrieving said differentially expressed nucleic acid or polypeptide.

A "gene expression profile" means the list of tissues, cells, etc., in which a defined gene is expressed (i.e, transcribed and/or translated). A "cell expression profile" means the genes which are expressed in the particular cell type. The profile can be a list of the tissues in which the gene is expressed, but can include additional information as well, including level of expression (e.g., a quantity as compared or normalized to a control gene), and information on temporal (e.g., at what point in the cell-cycle or developmental program) and spatial expression. By the phrase "selecting a gene or cell expression profile," it is meant that a user decides what type of gene or cell expression pattern he is interested in retrieving, e.g., he may require that the gene is differentially expressed in a tissue, or he may require that the gene is not expressed in blood, but must be expressed in pancreas. Any pattern of expression preferences may be selected. The selecting can be performed by any effective method. In general, "selecting" refers to the process in which a user forms a query that is used to search a database of gene expression profiles. The step of retrieving involves searching for results in a database that correspond to the query set forth in the selecting step. Any suitable algorithm can be utilized to perform the search query, including algorithms that look for matches, or that perform optimization between query and data. The database is information that has been stored in an appropriate storage medium, having a suitable computer-readable format. Once results are retrieved, they can be displayed in any suitable format, such as HTML.

For instance, the user may be interested in identifying genes that are differentially expressed in a pancreas or kidney. He may not care whether small amounts of expression occur in other tissues, as long as such genes are not expressed in peripheral blood lymphocytes. A query is formed by the user to retrieve the set of genes from the database

5

10

15

20

25

having the desired gene or cell expression profile. Once the query is inputted into the system, a search algorithm is used to interrogate the database, and retrieve results.

Advertising, licensing, etc., methods

The present invention also relates to methods of advertising, licensing, selling, purchasing, brokering, etc., genes, polynucleotides, specific-binding partners, antibodies, etc., of the present invention. Methods can comprises, e.g., displaying tissue selective polynucleotide or polypeptide sequences, or antibody specific thereto, in a printed or computer-readable medium (e.g., on the Web or Internet), accepting an offer to purchase said gene, polypeptide, or antibody.

Other

5

10

15

20

25

30

A polynucleotide, probe, polypeptide, antibody, specific-binding partner, etc., according to the present invention can be isolated. The term "isolated" means that the material is in a form in which it is not found in its original environment or in nature, e.g., more concentrated, more purified, separated from component, etc. An isolated polynucleotide includes, e.g., a polynucleotide having the sequenced separated from the chromosomal DNA found in a living animal, e.g., as the complete gene, a transcript, or a cDNA. This polynucleotide can be part of a vector or inserted into a chromosome (by specific gene-targeting or by random integration at a position other than its normal position) and still be isolated in that it is not in a form that is found in its natural environment. A polynucleotide, polypeptide, etc., of the present invention can also be substantially purified. By substantially purified, it is meant that polynucleotide or polypeptide is separated and is essentially free from other polynucleotides or polypeptides, i.e., the polynucleotide or polypeptide is the primary and active constituent. A polynucleotide can also be a recombinant molecule. By "recombinant," it is meant that the polynucleotide is an arrangement or form which does not occur in nature. For instance, a recombinant molecule comprising a promoter sequence would not encompass the naturally-occurring gene, but would include the promoter operably linked to a coding sequence not associated with it in nature, e.g., a reporter gene, or a truncation of the normal coding sequence.

-93-

The term "marker" is used herein to indicate a means for detecting or labeling a target. A marker can be a polynucleotide (usually referred to as a "probe"), polypeptide (e.g., an antibody conjugated to a detectable label), PNA, or any effective material.

The topic headings set forth above are meant as guidance where certain information can be found in the application, but are not intended to be the only source in the application where information on such topic can be found. Reference materials

For other aspects of the polynucleotides, reference is made to standard textbooks of molecular biology. See, e.g., Hames et al., <u>Polynucleotide Hybridization</u>, IL Press, 1985; Davis et al., <u>Basic Methods in Molecular Biology</u>, Elsevir Sciences Publishing, Inc., New York, 1986; Sambrook et al., <u>Molecular Cloning</u>, CSH Press, 1989; Howe, <u>Gene Cloning and Manipulation</u>, Cambridge University Press, 1995; Ausubel et al., <u>Current Protocols in Molecular Biology</u>, John Wiley & Sons, Inc., 1994-1998.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever. The entire disclosure of all applications, patents and publications, cited above and in the figures are hereby incorporated by reference in their entirety, including U.S. Application Serial Nos. 60/372,669 April 16, 2003, 60/374,823 filed April 24, 2002, 60/376,558 filed May 1, 2002, 60/381,366 filed May 20, 2002, 60/403,648 filed August 16, 2002, 60/411,882 filed September 20, 2002, and 60/424,336 filed November 7, 2002.

5

10

15

TABLE 1

(Himmen) (I sad	ACCIN	Precomment eles of expression	Olici et designisies	Cyregenetic levere
TMD0024	XM_060945	thymus	none	1q22
TMD1779	XM_060946	thymus and PBL	none	1q22
TMD0884	XM_060947	thymus	skin and ovary	1q22
TMD0025	XM_060948	thymus	none	1q22
TMD1780	XM_089422	thymus	none	1q22
TMD1781	XM_089421	PBL	thymus	1q22
TMD0304	XM_060956	bone marrow and muscle	testis	1q22
TMD0888	XM_060957	bone marrow	lung, muscle and testis	1q22
TMD0890	XM_060959	bone marrow	lung and PBL	1q22

TABLE 2

5				
	Clone ID (gene code)	ACCN	Protein seq length	Domain Description
10	TMD1779	XM_060946	264	Transmembrane domain: 26 - 48 Transmembrane domain: 55 - 77 Transmembrane domain: 92 - 114 Transmembrane domain: 134 - 156 Transmembrane domain: 197 - 219
15	TMD0024	XM_060945	268	Transmembrane domain: 16 - 38 Transmembrane domain: 53 - 75 Transmembrane domain: 96 - 118 Transmembrane domain: 156 - 178 Transmembrane domain: 191 - 213 Transmembrane domain: 228 - 246
25	TMD0025	XM_060948	313	Transmembrane domain: 29 - 51 Transmembrane domain: 58 - 77 Transmembrane domain: 92 - 114 Transmembrane domain: 135 - 157 Transmembrane domain: 197 - 219 Transmembrane domain: 240 - 262 Transmembrane domain: 272 - 294
30 35	TMD0304	XM_060956	319	Transmembrane domain: 28 - 50 Transmembrane domain: 63 - 82 Transmembrane domain: 102 - 124 Transmembrane domain: 144 - 166 Transmembrane domain: 205 - 227 Transmembrane domain: 240 - 262 Transmembrane domain: 272 - 294
40	TMD0884	XM_060947	299	Transmembrane domain: 20 - 42 Transmembrane domain: 54 - 76 Transmembrane domain: 91 - 113 Transmembrane domain: 126 - 148

	W O 03/00/30	33		PC1/USU3/1149
			-95-	
				Transmembrane domain: 183 - 205
				Transmembrane domain: 226 - 248
				Transmembrane domain: 258 - 277
5	TMD0888	XM_060957	312	Transmembrane domain: 25 - 47
				Transmembrane domain: 59 - 78
				Transmembrane domain: 98 - 120
				Transmembrane domain: 141 - 163
				Transmembrane domain: 207 - 229
10				Transmembrane domain: 241 - 260
				Transmembrane domain: 270 - 292
	TMD0890	XM 060959	280	Transmembrane domain: 26 - 48
		_		Transmembrane domain: 122 - 144
15				Transmembrane domain: 180 - 202
				Transmembrane domain: 215 - 237
				Transmembrane domain: 252 - 269
	TMD1780	XM 089422	491	Transmembrane domain: 20 - 42
20		_		Transmembrane domain: 54 - 76
				Transmembrane domain: 91 - 113
				Transmembrane domain: 137 - 159
				Transmembrane domain: 190 - 212
		<u>.</u>		Transmembrane domain: 231 - 253
25				Transmembrane domain: 266 - 283
				Transmembrane domain: 304 - 326
				Transmembrane domain: 336 - 358
				Transmembrane domain: 379 - 401
30				Transmembrane domain: 437 - 459
30	TMD1781	XM_089421	91	Transmembrane domain: 63 - 85

	TMD0024 XM_060945	TMD1779 XM_060946	TMD0884 XM_060947	TMD0025 XM_060948	TMD1780 XM_089422	TMD1781 XM_089421	TMD0304 XM_060956	TMD0888 XM_060957
TMD0024 XM_060945		4						
TMD1779 XM_060946	no significant similarity							
TMD0884 XM_060947	74%(371nt)	no significant similarity	· Vone					
TMD0025 XM_060948	71%(222nt) 80%(73nt)	90%(605nt)	83%(54nt)					10
TMD1780 XM_089422	81%(114nt) 74%(186nt) 79%(113nt) 77%(99nt)	83%(71nt)	78%(90nt)	80%(84nt)	Constitution of the Consti			
TMD1781 XM_089421	91%(35nt) 77%(80nt)	no significant similarity	no significant similarity	no significant similarity	77%(179nt) 82%(46nt)	and the state of t		
TMD0304 XM_060956	no significant similarity	no significant no significant significant similarity similarity similarity	no significant similarity	no significant similarity	84%(39nt)	no significant similarity		
TMD0888 XM_060957	no significant similarity	no significant similarity	no significant similarity	84% (38nt)	no significant similarity	no significant similarity	73%(241nt)	All Victoria
TMD0890 XM_060959		no significant no significant no significant similarity similarity similarity	no significant similarity	no significant similarity	no significant similarity	no significant similarity	no significant similarity	84%(39nt)

LABLE 3

					<u> </u>				
TMD0888 XP_060957									46% (196aa)
TMD0304 XP_060956								50%(301aa)	36%(196aa)
TMD1781 XP_089421				3			34%(89aa)	41%(82aa)	38%(72aa)
TMD1780 XP_089422						51%(93aa) 49%(77aa)	39%(300aa)	45%(304aa) 43%(189aa)	42%(200aa)
TMD0025 XP_060948					52%(300aa)	37%(94aa)	39%(299aa)	40% (305aa)	36%(179aa)
TMD0884 XP_060947				46%(166aa)	55%(165aa) 47%(111aa)	52%(40aa)	36% (163aa)	41%(157aa)	32%(156aa)
TMD1779 XP_060946			36%(92aa)	73%(233aa)	46%(227aa) 46%(169aa)	35%(82aa)	37%(229aa)	37%(239aa)	32%(132aa)
TMD0024 XP_060945		47%(200aa)	62%(171aa)	53%(252aa)	59%(261aa) 59%(181aa)	40%(94aa)	40%(257aa)	49%(251aa)	41%(196)
	TMD0024 XP_060945	TMD1779 XP_060946	TMD0884 XP_060947	TMD0025 XP_060948	TMD1780 XP_089422	TMD1781 XP_089421	TMD0304 XP_060956	TMD0888 XP_060957	TMD0890 XP_060959

FABLE 4

-99

	િ-'હોતાહેંહ)	ાં સ્વાહ	SELECTION (SELECTION)
TMD1779	GGTCAATGAGACTGTGG	GTGTGGAA	YOY
(SEQ ID NO 1-2)			ID NO 5)
TMD0024	CCACCTGCTCTCAGACA	GGCACCATAATTACCAGGAT	GAGTGCCAAATATATAAAGAGGTATGTTCAATGCAACATGTTAAATGCAA
6-7)	(SEQ ID NO 8)	(SEQID NO 9)	VECETION OF STANDARD AND STANDARD STANDARD AND STANDARD AND STANDARD STANDARD STANDARD STANDARD
TMD0025	ACTCTGGGCA	CTGGTTGGAGGAGTGGAAG	TAATACTATGTAAAAATCCACTGGACTAGAATCAGCTGTCCTCATGTGCC
(SECTID NO 12-13)	(SEQ ID NO 14)	GGCAG (SEQ ID NO 15)	(SEQ ID NO 19) TACCTTTCTGTATATAAAAACATATAACTAATACACACAC
			(SEQ ID NO 16)
			CITCAGAAGIAAAIGAAAIGAAGACIGGATACCAGCAAGACATACTGAATG
			CCCTTGGAGATATAAAAGTTCCCAGTAAATAGATGTGTGCCTCACATCTT (SEQ ID NO 18)
TMD0304 (SEQ ID NO	CTCTATGTTCCCGCATGC GCACAG	GCAAGGTGGAAATCCATGCA ATCTCAG	OTCTATGITCCCGCATGC GCAAGGTGGAAATCCATGCAAGACAGACGTTAAAAAATGACCAAACCTACAGAAAATTICCAGATAAT GCACAG
20-21)	10 22)	. 1	
TMD0884	TGTCAATATCCTGGTGTT	CATCTACCCAGAACCTTTCT	GTCACTGGTGTATAAGCACGCAGTGCAAAGGAAATATTAAAACTAGAACC
(35-26) 25-26)	(SEQ ID NO 27)	(SEQ ID NO 28)	(SEGIONO 23) TITCHTCATHTATAACATGAGGGGGCTTGGCTAGATATHAACAGCCTGC (SEQ
			ID NO 30)
			GCTAGATATTTAACAGCCTGCCTGTATTGACCACTTATGCATCAGGAAAT (SEQTD NO 31)
			ATTTGAGTTATGTATGAGAGACTGGGTACATCACTTTTTACTTGTTTT (SEQID NO 32)
TMD0888	GGAACTGGAGCCAGGTA	GGAGCAGGGATCAGCAGG	ACACTGCAGTTATATATGGGTGGCCCAGGTAGTTGAGCTGGTGAAATTTGA
33-34)	(SEQ ID NO 35)	(SEQ ID NO 36)	(SECTIONS 2)
			(SECTIONO 36) ATTCAAATTATATATATATTGGTCCAGTACGGTATCAATATATTATCAGTA (SEQ
			ID NO 39)
TMD0890	TCACCACCACTGGGACC	GGCCACACCAATCACTGTGC	CAATCTGTTATTTATACGGCCTCTACATCCATCCAGTACCTGCTTATGTA (SEQ)
4041)	2)	(SEQ ID NO 43)	GTTCTCTTTTTATAAAAGGCTATGTGGGACTTGCAAAACTTCTAGTGGCC (SEQ
			D NO 43) AGCATGAAATATAAGTAGGGAGTATCTTGGGGTAGAAAGGATGCCGAG (SEQ ID NO 46)
TMD1780	O	SGGAAGCACAGGT	ATCAATATTGTTAAAATGGCCGTACTGTCAAAAGCAATTTACAGATTCAA
(SEC ID NO 47-48)	(SEQ ID NO 49)	(SEQ ID NO 50)	(SEGIDINOS)) ATATGAAACCAAAAAAGCCCTCAAATAGCCCAAGTAACCCTAAAGAAAAA
			(SEQ ID NO 52)
			COCCUTATIONAL AND TOTAL OF THE COCCUTATION OF THE COCCUTATION OF STATES OF THE COCCUTATION OF THE COCCUTATI
			CATAAGGGTTCTTAAAATTGGGAGAGAGAATCAGAAAGTCAGAGAAGGG (SEQ ID NO 54)
TMD1781	ATGACAGTTTATGATTCC	TGTGAACAATG	TTCCCTATTTAATAAATGGTGCTGGGAAAACTGGCTAGCCATATGTAGAA
(35-56) 55-56)	(SEQ ID NO 57)	(SEQ ID NO 58)	AACAACCCCATCAAAAGTGGGCCAAAGATATGAACAGACACTTCTCAAA
			(SEQ ID NO 60) AATGGGGATCATTAAAAAGTCAGGAAACAACAGGGGGGGG
			(SEO ID NO 61) CCCAGAGGATTATAAATCATGCTGTAAAGACACATGCCCACGTATGT
			(SEQ ID NO 62)

TABLE 5

PRIMER (FOR, REV)	(SEQ ID NO)	02,70
PROMOTER (SEQ ID NO)		65-68
GENBANK PREDOMINANT PROMOTER IDENTIFIER SITES OF (SEQ ID NO)	EXPRESSION	
SEQ GENE GENBANK ID NUMBER IDENTIFIER		63,64 TMD0785 XM 060310 kidney
GENE NUMBER		TMD0785
SEQ ID	NO	63,64

	XM_062147	XM_062147 XM_061676
outside	1-27	1-28
TM (1)	28-50	29-51
inside	51-61	52-62
TM (2)	62-84	63-85
outside	86-58	66-98
TM (3)	99-121	100-122
inside	122-140	123-133
TM (4)	141-163	134-156
outside	164-203	187-201
TM (5)	204-226	202-224
inside	227-237	225-236
TM (6)	238-260	237-259
outside	261-274	260-273
TM (7)	275-293	274-296
inside	294-313	297-314

LABLE

obno i Digeno දෙන්වා ් කිලියා මාස්තා විය විය මාස්තාවේ පි	AGGN WELL	DOOD	প্রক্রেজনাদ্রনা প্রক্রিজন্তি	Other expression sites
TMD0049	XM 057351	Homo sapiens similar to organic anion transpoter 4 like protein (LOC116085) mRNA	kidney	none
TMD0190	XM 087157	Homo sapiens similar to sodium-coupled ascorbic acid transporter 2(LOC151295), mRNA.	kidney	colon and liver
TMD0242	XM 088369	Homo sapiens similar to unnamed protein product (LOC157724) mRNA	kidney	none
rMD0335	096680_MX	Homo sapiens similar to sodium iodide symporter (LOC159963) mRNA	kidney	adrenal gland, heart, intestine(small), liver, muscle, testis
TMD0371 (new)	XM 089732	Homo sapiens similar to CG8271 gene product (LOC196023), mRNA.	kidney	pancreas and testis
TMD0374 (new)	XM_085595	Homo saplens similar to unnamed protein product (LOC146802) mRNA	kidney	brain, muscle, ovary, skin, lestis
TMD0469	XM_038736	Homo saplens solute carrier family 4 sodium bicarbonate cotransporter member 9 (SLC4A9) mRNA	kidney	none
FMD0719	XM 059548	Homo sapiens hypothetical gene supported by XM_059548 (LOC131920) mRNA	kidney	none
TMD0731	XM 059703	Homo sapiens similar to putative (H. sapiens) (LOC134288) mRNA	kidney	adrenal gland, muscle, thyroid
TMD0785	XM_060310	Homo sapiens similar to olfactory receptor MOR275-2 (LOC127069), mRNA	kidney	none
TMD0841	XM 060623	Homo sapiens similar to KIAA0711 gene product (H. sapiens) (LOC127707) mRNA	kidney	lung
TMD1114	NM 019841	Homo sapiens transient receptor potential cation channel subfamily V member 5 (TRPV5) mRNA	kidney	none
TMD1148	XM 087108	Homo sapiens similar to calcium channel voltage-dependent gamma subunit 6 (LOC151151) mRNA	kidney	none

TABLE 8

ABLE 9

ලිනෙල දැන්ල	SEG (D (NO)	ි ිි	domein desentation
TMD0049	2	332	Sugar (and other) transporter: 2 - 302
			Transmembrane domain: 12 - 34
			Transmembrane domain: 39 - 58
			Transmembrane domain: 131 - 153
			Transmembrane domain: 157 - 179
			Transmembrane domain: 186 - 205
			Transmembrane domain: 215 - 237
TMD0190	4	243	Permease family: 91 - 224
TMD0242	œ	470	AA-nermease: 27 - 356
			Transmembrane domain: 13 - 35
			Transmembrane domain: 50 - 72
			Transmembrane domain: 93 - 115
			Transmembrane domain: 137 - 154
			Transmembrane domain: 161 - 183
			Transmembrane domain: 207 - 229
			Transmembrane domain: 242 - 264
			Transmembrane domain: 286 - 308
			Transmembrane domain: 335 - 357
			Transmembrane domain; 362 - 379
			Transmembrane domain: 392 - 414
			Transmembrane domain: 420 - 442

TMD0469	14	983	HCO3- transporter family: 108 - 891
			Transmembrane domain: 413 - 435
			Transmembrane domain: 447 - 469
			Transmembrane domain: 498 - 520
			Transmembrane domain: 532 - 554
			Transmembrane domain: 623 - 645
			Transmembrane domain: 665 - 684
			Transmembrane domain: 712 - 731
			Transmembrane domain: 751 - 773
			Transmembrane domain: 813 - 832
			Transmembrane domain: 839 - 858
			Transmembrane domain: 897 - 919
TMD0719	16	146	Transmembrane domain: 7 - 29
			Transmembrane domain: 49 - 71
TMD0731	18	218	Transmembrane domain: 38 - 60
			Transmembrane domain: 70 - 92
1/-			
TMD0785	20	312	7 transmembrane receptor (rhodopsin family): 58 - 290
			Transmembrane domain: 29 - 51
'n	<u>. </u>		Transmembrane domain: 61 - 83
			Transmembrane domain; 140 - 162
			Transmembrane domain: 197 - 219
			Transmembrane domain: 240 - 262
			Transmembrane domain: 272 - 294
TMD0841	22	1161	Kelch motif: 850 - 895
			Kelch motif: 897 - 938

	1	•
ļ	1	2
ς.		-
•		•

	327 - 349	383 - 405	420 - 438	451 - 473	493 - 512	519 - 541	554 - 576	7 - 24	39 - 61	98 - 90	
	Transmembrane domain: 327 - 349	Transmembrane domain: 383 - 405	Transmembrane domain: 420 - 438	Transmembrane domain: 451 - 473	Transmembrane domain: 493 - 512	Transmembrane domain: 519 - 541	Transmembrane domain: 554 - 576	Transmembrane domain: 7 - 24	Transmembrane domain: 39 - 61	Transmembrane domain: 68 - 90	
	729							103			
	24							26			
	TMD1114							TMD1148			

Gione ID (gene code) ACCN		Ovogeneticio de P	Ovegeneticlocus diseasellinkage start in the sease in the
TMD0049	XM 057351	11912.1	osteoporosis-pseudoglioma syndrome; spastic paraplegia 1 7
TMD0190	XM_087157	2936.2	auou
TMD0242	XM 088369	8q21.2	euou
TMD0335	XM_089960	11p14.2	none
TMD0371A	XM_089732	10q23.33	epilepsy, partial, with auditory features: spastic paraplegia 9, autosomal dominant
TMD0374	XM_085595	17p11.2	smith-magenis syndrome
TMD0469	XM_038736	5q31	paget disease of bone 4
TMD0719	XM 059548	3q29	9000
TMD0731	XM_059703	5q13.2	spastic paraplegia 11, autosomal recessive; corpus callosum, agenesis of, with neuronopathy
TMD0785	XM_060310	1q44-tel	familial cold urticarla (FCU); Muckle-Wells syndrome (MWS); prostate cancer susceptibility
TMD0841	XM_060623	1p36.13	breast cancer, ductal, 2; prostate cancer/brain cancer susceptibility; melanoma, cutaneous
TMD1114	NM_019841		glaucoma 1, open angle, f
TMD1148	XM_087108	2914.1	motor neuronopathy, distal hereditary, with vocal cord paralysis; cardiomyopathy, dilated, 1h

ABLE 10

ල්ලමය් ((නූල්) දල (ලමා	gliceds (1930 to 100)	[1] A SERVICE (SERVICE NO.
TMD0049	GCGCTTCCGGACCTGTATCTCCAC (104)	aargagcctctaaagaagggttccagactaccaggagctcactggaaata (106)
(78,79)	CAAGCTCTGGGTCTCGGGCAGAAG (105)	
TMD0190	ACCATCCTGCAACTTGGATGGGC (107)	GCTITATGTATATGAAAACCCTGTTTATCTGAGCCTAGAACTGTCTTTGC (109)
(80,81)	aaggagccggaagacaggagg (108)	agtgatagtittaaatgggagggaataaagtctgcaaaatttccccatat (110)
TMD0242	GAGTCTCCCTGTGCGTTTGGGCTG (111)	AGTCCCAGCTTAAAAAAGAGACAGACAGACAGAGAGAGAG
(82,83)	AAGTGTAAAGCATGCCCGCCTGA (112)	TTAGTGATTTAAAAAAATGTGAAGAAGAGAGGCAGCAGTAAAAGGA (114)
TMD0335	GTTCGCTATGCTGCCACGGTCATC (115)	GATACAAATAATTAAAAGCCCAGGTTAAGGTAAATATTTAAAGACCAAG (117)
(84,85)	AGTCCTGGCAGTCCTGGCATTGTG (116)	ATCTCACGAATTAAAAATGCTGAGGTGGTAAATTGTTATCAATTCTATGT (118)
TMD0371	CAGGATTACGCACAAACGGCATGG (119)	CTAGACTATTTAAAAAACCCCTGGCTTGCACAGTGGCTCAAGCCTGTAA (121)
(86,87)	TGGGAGGCAGAGATAGCAGAGCCC (120)	
TMD0374	CTGGTCCTGGCCCCTGATAAGC (122)	AGCTGTCCTCATTAAAAGTGACCTGGAGTGAATGGATTCTTCTGCCTAT (124)
(88,89)	CCCAGGTCTGGTTGCAGTGCTCTC (123)	CCAATTCTTCTGAAAAACGGGAGTCACTGTGGGCACCATCACGCCCGGGT (125)
TMD0469	CTGAGGTGTCCCTCCCAAGCAGGT (126)	TAAACAAATACATAAATGAGGCAGTTACTAGTAGTGGTAACTGCTAGGAA (128)
(90,91)	TACGGCCGAGAAGCACTGGAGATG (127)	actaaaaatataaaaatcagccaggcctggtggcacatgtctgtaatctc (129)
		GGGATGCATTATAAATGCAACCAGCCCAGAGGGCCCCTGGCTTCAGAACCT (130)
TMD0719	GTCACCTCAGCGATCTCAACGATAGGG (131)	ATATACCTTGTTTAAAAGAGGGGTATTATCACAATAAAACAAGGAAAGCT (133)
(85, 93)	TGGAGCAGGAACAGGATATAGGTCAGGG (132)	ACCCCTACTTTTAAAGGCCTTGACAAACAGTGCTAAAGTTCTCACCTTAA (134)
TMD0731	GGGTGGGAAGGAAGAGAG (135)	TTATTGGGCATAAAAATATGAAGAGGTCCCCAGAGAGTCCCTAGGTTCT (137)
(94,95)	CCAGCTAGTTCATGCTTGGCGCAG (136)	
TMD0785 (96,97)	CTGTTGGGAATCTTCAGCCAGATCTCACAC (138) ATGGAGGTTTCTGCACGCTCAGCA (139)	AAGCAATTTGTTAAAAACTGGCATTACTTTACTCTTATGCTTTCTGTGTC (140) ACTTTAATTTTATAAAGAAGGTTCACATCAAGAAATTCCAAGTGAGGTTC (141)
TMD0841	GGGCCACTTCCACAGACAGGAAGC (142)	AAGGCTTCTTCAAAAAAGGGGGCTTGTTCTGGGCCAGAAAATCAGAGTG (144)
(68, 99)	TGGCCTGAGAGGTAGATTCCACATAGTAGTCGT (143)	
TMD1114	CTCCTTTCTGGTCAGAGACAAGACTGGGAC	CAGCGAGAGAAAAATGTCCCACAAAGTTGAGCCCTCCCCACTCCCAGTG (147)
	GIGATGTCTCGAGAATGAGTGCGGTTG	
(100, 101)	(140)	TARIRIARARINIARARING GCARCALINITATICCICCIGGIGII (148)
TMD1148	GCAGATGACCCGACCTGACTGTTCTTC (149)	GCCAGAGAGTTTAAATGAAGCCCTACTTTGGGGCAGGAGCGGGAGGAAAC (151)
(102, 103)	rescrerecascrascrassraccas (150)	

SEQ	GENE	GENBANK	PREDOMINANT	OTHER	PROMOTER	PRIMER
8	NUMBER	IDENTIFIER	SITES	SITES OF	(SEQ ID NO)	(FOR, REV)
NO			OF EXPRESSION	EXPRESSION		(SEQ ID NO)
152,	1MD0986	624190 MX	pancreas	low levels in	156-161	154,155
153				testis		
162,	TMD0987	XM_061780	pancreas	low levels in	166	164,165
163				testis		
167,	TMD0353	XM_061781	pancreas			169,170
168		•				
171,	TMD0989	XM_061784	pancreas			173,174
172						
175,	TMD058	XM_061785	pancreas	low levels in	179,180	177,178
176			-	testis		

TABLE 12

outside 1-23 1-25 1-24 TM (1) 24-46 26-48 23-45 1-24 inside 47-58 49-60 46-65 48-59 TM (2) 59-78 61-83 66-88 60-82 outside 79-97 84-97 89-97 83-96 TM (3) 98-120 98-120 98-120 97-115 inside 121-140 121-139 121-140 120-13 TM (4) 141-163 140-162 141-163 140-16 outside 164-198 163-202 164-203 163-20 TM (5) 199-221 203-25 204-256 202-22 TM (5) 241-260 238-260 238-260 237-23 TM (5) 241-269 261-272 261-272 outside 261-274 261-269 273-292 269-26 TM (7) 75-292 270-289 273-292 269-26 Tm (7) 275-27 273-27 273-27 273-27		WW_061779	XM_061780	XM_061781	XM_061779 XM_061780 XM_061781 XM_061784 XM_061785	XM_061785
24-46 26-48 23-45 47-58 49-60 46-65 59-78 61-83 66-88 79-97 84-97 89-97 98-120 98-120 98-120 121-140 121-139 121-140 141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	outside	1-23	1-25	1-22		1-24
47-58 49-60 46-65 59-78 61-83 66-88 79-97 84-97 89-97 98-120 98-120 98-120 121-140 121-139 121-140 141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (1)	24-46	26-48	23-45		25-47
59-78 61-83 66-88 79-97 84-97 89-97 98-120 98-120 98-120 121-140 121-139 121-140 141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	inside	47-58	49-60	46-65		48-59
79-97 84-97 89-97 98-120 98-120 98-120 121-140 121-139 121-140 141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (2)	82-28	61-83	88-99		60-82
98-120 98-120 121-140 121-139 141-163 140-162 141-163 140-162 141-163 140-162 164-198 163-202 199-221 203-25 222-240 226-237 241-260 238-260 241-260 238-260 261-274 261-269 261-274 261-269 253-314 290-318 293-314 290-318	outside	79-97	84-97	26-68		83-96
121-140 121-139 121-140 141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (3)	98-120	98-120	98-120		97-119
141-163 140-162 141-163 164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	inside	121-140	121-139	121-140		120-139
164-198 163-202 164-203 199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (4)	141-163	140-162	141-163		140-162
199-221 203-25 204-226 222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	outside	164-198	163-202	164-203		163-201
222-240 226-237 227-237 241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (5)		203-25	204-226		202-224
241-260 238-260 238-260 261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	inside	222-240	226-237	227-237		225-236
261-274 261-269 261-272 75-292 270-289 273-292 293-314 290-318 293-323	TM (6)	241-260	238-260	238-260		237-259
75-292 270-289 273-292 293-314 290-318 293-323	outside	261-274	261-269	261-272		260-268
293-314 290-318 293-323	TM (7)	75-292	270-289	273-292		269-291
	inside	293-314	290-318	293-323		292-311

FABLE 1:

GENBANK	MOUSE	061779	061780	061781	061784	061785
IDENTIFIER	HOMOLOG					
XM 061779			42% (63%)	36% (57%)	43% (64%) 40% (61%)	40% (61%)
XM_061780	MOR239-6 (AY073489) 90% (93%)	42% (63%)		41% (60%)	44% (62%)	46% (67%)
XM 061781		36% (57%) 41% (60%)	41% (60%)		43% (63%) 40% (61%)	40% (61%)

-110

KM 061784	MOROZ	130/ (610/)	(/003) /011	(/063) /061		(/050/ /010
	1410142)	(0/+0) 0/0+	(0/70) 0/44 (0/40) 0/64	45% (05%)		(%/8)%!8
	~>85%					
	MOR223	40% (61%)	46% (67%)	40% (61%)	81% (87%)	
	~>85%					8

TABLE 14

TABLE 15

ලියෙ (ලියෙ දෙයි)	AGGN	Preforment sites of expression	Outer expression elles	Gyloganellocus
TMD1030 (SEQ ID NO 185- 186)	XM_166853	spleen	liver	11q12.2
TMD1029 (SEQ ID NO 187- XM_166854-188)		spleen, lymphocytes, liver	brain, heart, lung, lymph node 11q12.2	11912.2
TMD1028 (SEQ ID NO 189- XM_166855 190)		spleen, lymphocytes	liver	11q12.2
TMD0621 (SEQ.ID NO 191- XM_166205	- 1	spleen	brain, heart, liver, lung and pancreas	11912.2

TABLE 16

	१ ग्वंसित सित्ति (हरू)	Conein description. 🚠 🔐
TMD1030 XM_166853	298	Transmembrane domain; 27 Transmembrane domain: 98 Transmembrane domain: 140 Transmembrane domain: 175

			Transmembrane domain: 238 - 260 Transmembrane domain: 275 - 292
TMD1029	XM_16684	306	Transmembrane domain: 26 - 48 Transmembrane domain: 61 - 78 Transmembrane domain: 98 - 120 Transmembrane domain: 140 - 162 Transmembrane domain: 196 - 218 Transmembrane domain: 238 - 260 Transmembrane domain: 275 - 292
TMD1028	FMD1028 XM_166855	173	Transmembrane domain: 18 - 40 Transmembrane domain: 61 - 83 Transmembrane domain: 103 - 125 Transmembrane domain: 137 - 156
TMD0621	FMD0621 XM_166205	109	Transmembrane domain: 9-31 Transmembrane domain: 69 - 91

TABLE 17

			,	
હ્યુંઓ	GAGCCTATAATATATGAGCCAGCTACGAGTTGGA (SEQ ID NO 198)	AAACCTGTTTGTACAGAGGCATTTATTGAGCC (SEQ ID NO 200)	CTCCAACCCAGTGAACATCAAGTTAAATCCCAC (SEQ ID NO 202)	CTCATTAATACGATGGCATAGATACATGTAAGAGAG (SEQ ID NO 204)
िन्दीष्ट्रिक संस्थान	TMD1030 XM_166853 GGGATTTGGTGTCCAACACGAATTTCA (SEQ ID NO 197)	TMD1029 XM_166854 GTCACTGAATTCTATCTTCTGGGATTTGGTGC (SEQ ID NO 199)	XM_166855 GATATCATTTTGGGGCTGCATGATACTATTGG (SEQ ID NO 201)	TMD0621 KM_166205 TTAAGCTATTAGTTAGTTCATATGTCATGGGTTTCC (SEQ ID NO 203)
KCCN	XM_166853	хм_166854	XM_166855	хм <u>_</u> 166205
(මානමෙම	TMD1030	TMD1029	TMD1028	TMD0621

TABLE 18

	ACCIN ^{SA} XM_166853		(0.94)	(SEQ	9	30 20	<u> </u>
TMD1029	XM_166854	ACAICCATTATATACAGGGTTAATATATGTGCAGGATTAGCACCTAGA (0.95) (SEQ ID NO 205) XM 166854 AAATGTATAAATTCTGCATGAAATTGGGGGTGGGGCTTGTACTACTTTTG (0.98) (SEQ ID NO 207)	(0.95)	(SEQ	8 8	10 20 30 30 30 30 30 30 30 30 30 30 30 30 30	(9)
1	XM 166855	TMD1028 XM 166855 ATGTTCCATCTAAATGAAGCCTGAGAAACCCAGCACTACCCACTTGTTAG (0.94) (SEO ID NO 208)	(0.94)	(SEO	QI	20	(8)
	1	ACATCCATTATATAACAGGGTTAATATACTTGTAAAGAATAGCACCTAGA (0.95) (SEQ ID NO 209)	(0.95)	(SEQ	2	10 20	

-51

(SEQ ID NO 210)	(SEQ ID NO 211)	(SEQ ID NO 212)	(SEQ ID NO 213)	
(66.0)	(0.97)	(1.00)	(0.91)	
[TMD062] XM 166205 AAATATATATTTTAAATTGGCCAGGCGGTGGCTCACGCCTATAAATCCC (0.99) (SEQ ID NO 210)	GGCTCACGCCTATAATCCCAGCACTTTGGGAGGCCGAGGCAGGTGGATCA (0.97) (SEQ ID NO 211)	TCCCAAATATATATATATACACACACACACACACACACA	CACACACACATATATACACACACATATATTATAATCATTTAACAAC	
XM 166205		• • • • •		
TMD0621				tramaras.

BNSDOCID: <WO, 03089583A2 1 >

TABLE 19

TABLE 19

(from Principles of Internal Medicine, Volume 1, Page 357, 12th Edition, McGraw-Hill Inc.)

WO 03/089583

euou 	pancreas and testis	Homo sapiens putative ion channel protein CATSPER2 (CATSPER2), mRNA.	NM_054020	TMD1127
none	pancreas and testis		NM_014386	TMD1111
sionnach	pancreas and testis	Homo sapiens similar to putative (H. sapiens) (LOC138240) mRNA	XM_059954	TMD0753
liver, lung, mammary gland, ovary, pituitary, prostate and stomach	pancreas and testis	Homo sapiens similar to putative (H. sapiens) (LOC135886) mRNA	XM_059812	TMD0739
stomach	pancreas and testis	Homo sapiens similar to testis-specific transporter TST1 (H. sapiens) (LOC133482) mRNA	XM_059654	TMD0727
adrenal gland. brain, prostate and stomach	pancreas and testis	Homo sapiens similar to hypothetical protein (H. sapiens) (LOC133309)	XM_059639	TMD0726
prostate and stomach	pancreas and testis	Homo sapiens similar to dJ39G22.2 (novel protein) (H. sapiens) (LOC127391) mRNA	XM_059140	TMD0677
prostate and stomach	pancreas and testis	Homo sapiens similar to putative (H. sapiens) (LOC127348) mRNA	XM_059134	TMD0675
brain, pituitary, prostate and stomach	pancreas and testis	Homo sapiens similar to RIKEN cDNA 4930549C01 gene (LOC127309) imRNA	XM_059132	TMD0674
bone marrow, brain, heart, kidney, liver, lung, lymph node, PBL, muscle, pituitary, prostate, skin, spleen, stomach and thymus	pancreas and testis	Homo sapiens LOC146225 (LOC146225), mRNA.	XM_085376	TMD0645
liver, PBL and prostate	pancreas and testis	Homo sapiens similar to data source:MGD, source key:MGI:96073, evidence:ISS-hexosaminidase A-putative (LOC204249), mRNA.	XM_058690	TMD0639
stomach	pancreas and testis	Homo sapiens similar to putative (H. sapiens) (LOC118670) mRNA	XM_058332	TMD0608
pituitary	brain and pancreas	Homo sapiens KIAA1910 protein (KIAA1910) mRNA	XM_055514	TMD0574
brain, kidney, lung, lymph node, PBL, mammary gland, pitultary, stomach, testis and thyroid	pancreas	Homo sapiens hypothetical protein DKFZp564A1164 (DKFZP564A1164) mRNA	XM_048304	TMD0530
опоп	pancreas and testis	Homo sapiens similar to unnamed protein product (LOC130644) mRNA	XM_065813	TMD0290
PBL, prostate, thymus and uterus	pancreas and testis	Homo sapiens similar to odorant receptor S18 gene (LOC120010) mRNA	XM_061815	TMD0271
adrenal gland, bone marrow, colon, heart, intestine(small), kidney, liver, pituitary, prostate, skin, stomach and thyroid	pancreas and testis	Homo sapiens similar to CG5281 gene product (LOC159371) mRNA	XM_089550	TMD0267
colon, stomach and testis	pancreas	Homo sapiens similar to beta-2 adrenergic receptor (LOC139760) mRNA	XM_066873	TMD0258
skin and testis	pancreas	receptor (LOC139478) mRNA	XM_066725	TMD0256
none	pancreas	Homo sapiens similar to olfactory receptor (LOC135941) mRNA	XM_069616	TMD0233
brain, heart and kidney	pancreas and testis	Homo sapiens olfactory receptor MOR212-1 (LOC219956), mRNA.	XM_166914	TMD0077
Comment of the respicasion selection of the selection of	Předomínaní ellesof.	Gene Nama/Dascription	AGGN	්ල්කාව [ව

Table 21

Cone (D	ACCN	Protein seq lenglin (aa)	Domain description
TMD0077	XM_166914	310	7 transmembrane receptor (rhodopsin family)
			Transmembrane domains: 27 - 49
			Transmembrane domains: 61 - 83
			Transmembrane domains: 98 - 120
			Transmembrane domains: 141 - 163
			Transmembrane domains: 202 - 224
			Transmembrane domains: 237 - 259
			Transmembrane domains: 274 - 291
TMD0233	XM_069616	310	7 transmembrane receptor (rhodopsin family)
	•		Transmembrane domain: 26 - 48
			Transmembrane domain: 60 - 77
			Transmembrane domain: 97 - 119
			Transmembrane domain: 140 - 162
			Transmembrane domain: 196 - 218
			Transmembrane domain: 239 - 261
			Transmembrane domain: 272 - 291
TMD0256	XM_066725	308	7 transmembrane receptor (rhodopsin family)
			Transmembrane domain: 27 - 49
			Transmembrane domain: 61 - 83
			Transmembrane domain: 98 - 120
			Transmembrane domain: 140 - 162
			Transmembrane domain: 196 - 218
			Transmembrane domain: 239 - 258
			Transmembrane domain: 273 - 291
TMD0258	XM_066873	335	7 transmembrane receptor (rhodopsin family)
			Transmembrane domain: 10 - 32
			Transmembrane domain: 39 - 61
			Transmembrane domain: 79 - 101
			Transmembrane domain: 121 - 143
			Transmembrane domain: 163 - 185
			Transmembrane domain: 226 - 248
			Transmembrane domain: 263 - 282
TMD0267	XM_089550	324	Integral membrane protein DUF6: 49-161
			Transmembrane domain: 59 - 78
			Transmembrane domain: 91 - 110
			Transmembrane domain: 115 - 137
			Transmembrane domain: 146 - 168
			Transmembrane domain: 183 - 201
			Transmembrane domain: 214 - 236
			Transmembrane domain: 246 - 265

	·		
			Transmembrane domain: 270 - 292
			Transmembrane domain: 297 - 316
			·
TMD0271	XM_061815	291	7 transmembrane receptor (rhodopsin family)
			Transmembrane domain: 29 - 51
			Transmembrane domain: 56 - 78
			Transmembrane domain: 83 - 105
			Transmembrane domain: 120 - 142
			Transmembrane domain: 163 - 185
			Transmembrane domain: 190 - 207
			Transmembrane domain: 220 - 239
			Transmembrane domain: 249 - 271
TMD0290	XM 065813	245	Transmembrane domain: 24 - 46
			Transmembrane domain: 61 - 83
			Transmembrane domain: 96 - 118
			Transmembrane domain: 128 - 150
			Transmembrane domain: 162 - 184
			Transmembrane domain: 221 - 243
TMD0530	XM 048304	· 708	Immunoglobulin domain: 139-206
			Immunoglobulin domain: 326-377
			Transmembrane domain: 511 - 533
TMD0574	XM 055514	696	Leucine rich repeat C-terminal domain: 212-262
			Leucine rich repeat C-terminal domain: 529-579
			Transmembrane domain: 621 - 643
TMD0608	XM 058332	105	Transmembrane domain: 13 - 35
TMD0639	XM 058690	127	Transmembrane domain: 12 - 34.
-			Transmembrane domain: 44 - 66
TMD0645	XM 085376	248	Transmembrane domain: 113 - 135
			Transmembrane domain: 150 - 169
			Transmembrane domain: 176 - 198
TMD0674	XM 059132	134	Transmembrane domain: 5 - 22
TMD0675	XM 059134	206	Transmembrane domain: 15 - 37
TMD0677	XM 059140	182	Transmembrane: 49 - 71
	2.50.7.0	<u></u>	
TMD0726	XM_059639	96	Transmembrane domain: 13 - 35
			Transmembrane domain: 50 - 72
	related to		Į ·

	· · · · · · · · · · · · · · · · · · ·		
			Transmembrane domain: 145 - 164
			Transmembrane domain: 171 - 193
			Transmembrane domain: 229 - 251
			Transmembrane domain: 264 - 286
			Transmembrane domain: 314 - 336
			Transmembrane domain: 421 - 443
			Transmembrane domain: 453 - 475
			Transmembrane domain: 580 - 602
			Transmembrane domain: 668 - 690
			Organic Anion Transporter Polypeptide (OATP) family, C-terminus: 125-473
			Organic Anion Transporter Polypeptide (OATP) family, N-terminus: 558-717
		<u>.:</u>	
TMD0739	XM_059812	265	Transmembrane domain: 126 - 148
			Transmembrane domain: 185 - 207
TMD0753	XM_059954	161	Transmembrane domain: 26 - 48
TMD1111	NM 014386	. 609	Ion transporter domain: 284-490
***************************************		- 000	Transmembrane domain: 34 - 56
			Transmembrane domain: 274 - 296
			Transmembrane domain: 315 - 337
			Transmembrane domain: 364 - 386
·····			Transmembrane domain: 407 - 429
			Transmembrane domain: 469 - 491
TMD1127	NM_054020	528	Ion transporter domain: 172-340
			Transmembrane domain: 113 - 132
			Transmembrane domain: 147 - 169
			Transmembrane domain: 176 - 198
			Transmembrane domain: 241 - 263
			Transmembrane domain: 276 - 295
			Transmembrane domain: 315 - 337

Table 22

@lone ID	ACCN	(Cytegeneticlecus	egsibil exessib
TMD0077	XM_166914	11q12.2	angioedema, hereditary, spastic paraplegia 17; osteoporosis- pseudoglioma syndrome; pancreatic tumor
TMD0233	XM_069616	7q35	glaucoma 1, open angle, f;
TMD0256	XM_066725	Xq26.1	x inactivation, familial skewed, 2; panhypopituitarism; thoracoabdominal syndrome; dandy-walker malformation with mental retardation, basal ganglia disease, and seizures; split-hand/foot malformation 2; mental retardation with optic atrophy, deafness
TMD0258	XM_066873	Xq26.1	x inactivation, familial skewed, 2; panhypopituitarism; thoracoabdominal syndrome; dandy-walker malformation with mental retardation, basal ganglia disease, and seizures; split-hand/foot malformation 2; mental retardation with optic atrophy, deafness
TMD0267	XM_089550	10q24.1	comeal dystrophy of bowman layer, type ii; alzheimer disease 6
TMD0271	XM_061815	11p15.4	charcot-marie-tooth disease, type 4b, form 2; deafness, neurosensory, autosomal recessive 18;
TMD0290	XM_065813	2p23.1	none
TMD0530	XM_048304	19q13.13	hypocalciuric hypercalcemia, familial, type iii; deafness, autosomal dominant nonsyndromic sensorineural 4; microcephaly, primary autosomal recessive. 2
TMD0574	XM 055514	13q31.1	microcoria, congenital; schizophrenia 7;
TMD0608	XM_058332	10q26.3	endometrial carcinoma
TMD0639	XM_058690	15q22.32	cataract, central saccular, with sutural opacities; obesity syndrome
TMD0645	XM_085376	16q23.1	dehydrated hereditary stomatocytosis; pancreatic acinar cancer
TMD0674	XM_059132	1p36.11	breast cancer, ductal, 2; prostate cancer/brain cancer susceptibility; melanoma, cutaneous malignant; inflammatory bowel disease 7;
TMD0675	XM_059134	1p33	carcinoma of pancreas
TMD0677	XM_059140	1p34.2	deafness, autosomal dominant nonsyndromic sensorineural 2; porphyria cutanea tarda; hypercholesterolemia, familial, plosis, hereditary congenital 1 ;
TMD0726	XM_059639	10q11.22	none
TMD0727	related to XM_059654	5q21.1	anemia, dyserythropoietic congenital, type iii; dyslexia, specific, 1; colorectal cancer, hereditary nonpolyposis, type 7; cataract, central saccular, with sutural opacities
TMD0739	XM_059812	7q11.23	autism, susceptibility to, 1; muscular dystrophy, limb-girdle, type 1d; aneurysm, intracrania l
TMD0753	XM_059954	9q21.12	hemophagocytic lymphohistiocytosis, familial, 1; amyotrophic lateral sclerosis with frontotemporal dementia
TMD1111	NM_014386	5q31	, none
TMD1127	NM_054020	15q13-q15	nanophthalmos 2; spastic paraplegia 11, autosomal recessive; corpus callosum, agenesis of, with neuronopathy; pancreatic acinar carcinoma

4	1800	TABLE 23	
CODE TMD0077 (SEQ ID NO 214- 215)	ACCN XM_166914	PRIMERS TCATGGATCACCAGCTCCACGCTC (Forward) (SEQ ID NO 256) CACCAGATCACCACCATGGAAGCA (Reverse) (SEQ ID NO 257)	GGATTCAGGCCTTTTAAAACCCCACTCAGTGGGTGCATGGCAGGGCTTTGA (0.88) (SEQ ID NO 258)
TMD0233 (SEQ ID NO 216- 217)	XM_069616	TGCTGACGAATCTTATGAACCAGG (Forward) (SEQ ID NO259) TCACGTCAGCCTCTCCTTCCTCAGTG (Reverse) (SEQ ID NO 260)	TCACAAATCATATAAATTAGGGGAAAGAGAGGGCAGGTATACTCTAAAAA (0.96) (SEQ ID NO 261) AATTTCTTATTTAAAAAGACCTCAGAAATGTCACCATGCTTAGTTATTTA (0.95) (SEQ ID NO 262)
TMD0256 (SEQ ID NO 218- · · 219)	XM_066725	GGCCATGGACAATGTCACAGCAG (Forward) (SEQ ID NO 263) AGCAGACACATACTGGGCCATTCATAACCAC (Reverse) (SEQ ID NO 264)	GGTACTATTCTATATTTTGGGCACACAGCAATGAAGAAACAGAAAAACC (0.93) (SEQ ID NO 265) CTGGGTTCATAAAATATGGAGCAGAAAGTTTTTACAAATATAGAACAGCA (0.92) (SEQ ID NO 266) TAGAATGTGTTATAAAAAAATGAAGCAGGGCTAGGGGGAAAGAGGTGGTGA (0.91)
TMD0258 (SEQ ID NO 220- 221)	XM_066873	CCTCATTGGCTTCCTCCCACTCG (Forward) (SEQ ID NO 268) GCCATCAAACTCTGAGCTGGAGATAGTGAC (Reverse) (SEQ ID NO 269)	CANGORACTITITAAAACTCCCATTGCACAGTTACCACCCAGAATAATTA (0.97) (SEQ ID NO 270 CATCCTGGAATATATTTGCGTCCAACTCTGCACCTTGCTCTCTATTCCCT (0.96) (SEQ ID NO 27) CTGGGGCCCTCAAAAAGCTCACCTTCCCTCACTTCCCACTTCAACTGAT (0.91)
TMD0267 (SEQ ID NO 222- 223)	XM_089550	TGGCCTCGTTGAAAGTGTCATCATCC (Forward) (SEQ ID NO 273 TTGGTACCATTTACGAATGGCCGC (Reverse) (SEO ID NO 274)	AAACGGCATTITAAAAATGCAGGTTTAAATTGTTATCCTCATCTATGGTT (0.98) (SEQ ID NO 275)
TMD0271 (SEQ ID NO 224- 225)	XM_061815	CTGGACTTGAGCAGTACCACGTCTGGATC (Forward) (SEQ ID NO 276) CATATTCCCACAGCAATTTTGACAATGG (Reverse) (SEQ ID NO 277)	ATTITGGTTATATATAGAGGAGTCTAGGAAAAGACTCGTGGGTCTGATTC (0.97) (SEQ ID NO 278) TACTCATATTTATATAGCAGCAACTTACATTGACCCAGGGAGAACTCAGT (0.94) (SEO ID NO 279)
TMD0290 (SEQ ID NO 226- 227)	XM_065813	GTTACCCACCCAACCGTCACGACC (Forward) (SEQ ID NO 280) CAGGCGATGCCAGAGAAGACGATG (Reverse) (SEQ ID NO 281)	CTGGGGGTCTATAGAAAAAGGACTGGAGGAGCTTTTGCAGCAACTTTGCAT (0.97) (SEQ ID NO 282) TITTCTTCTTTTAAAAACACGCTTTCACTCTCAAAACAGCAGGAGTGAA (0.98) (SEQ ID NO 283) AACTGGGGTCTATAAGAGAGCCAGGGCACTTATTCATCCAGGGCAGATG (0.99)
TMD0530 (SEQ ID NO 228- 229)	XM_048304	CTATGACTTCAACCCACACCTGGGCA (Forward) (SEQ ID NO 285) AAGGTCGCCAACTTGTCCTGGCTC (Reverse) (SEQ ID NO 286)	GGGCGGGAGTAAAAGGCAGAGTCCAATTCCACCGGCCCCCAGTGTGGGTG (0.86) (SEQ ID NO 287)

CODE	ACCN	PRIMERS	PROMOTER
TMD0574 (SEQ ID NO 230- 231)	XM_055514	TCAATGCCCAAACTGAGGA (Forward) (SEQ ID NO 288) CAACACCGAGATGGACACCCTGCT (Reverse) (SEQ ID NO 289)	CTTTTAAGGTTAAAAATGTGGGTTTTAGATGATTGTCCTTTCTAAACAGC (0.99) (SEQ ID NO 290) TCAGGATGTCTAAAAAAGATCTCTCTAGTGTACACACGTGCACACACA
TMD0608 (SEQ ID NO 232- 233)	XM_058332	CTCAGGACGAAGATCATGATCGGCATC (Forward) (SEQ ID NO 294) GAAGATTTTTGTGCCCAGCTTTCCCAAG (Reverse) (SEQ ID NO 295)	TATTCTCACTTATAAGTGGGAGCTAAGCCATGAGGGCACCAAGGCATAAG (0.99) (SEQ ID NO 296) TTACATATGTATACATGTGCCATGCTGGTGTGCTGCACCATTAACTCGT (0.96) (SEQ ID NO 297)
TMD0639 (SEQ ID NO 234- 235)	XM_058690	TCCATGCTCAGCTTCATCTCAGCTACC (Forward) (SEQ ID NO 298) TCCATCTCAGACCTTGGCCCTTCA (Reverse) (SEQ ID NO 299)	AAATAACCCCATTAAAAAGTGGGCAAAGGGCATGAACACTTTTCAAAAGA (1.00) (SEQ ID NO 300)
TMD0645 (SEQ ID NO 236- 237)	XM_085376	AGGACGGTAAGGAGCCATCGGACA (Forward) (SEQ ID NO 301) CTTGCCAGGTTCTGGTGGCTTGG (Reverse) (SEQ ID NO 302)	TCTTTTTGTCTATAAATAGGACTTTGATTTTCTGGACTAGAGAATTGTAT (0.94) (SEQ ID NO 303)
TMD0674 (SEQ ID NO 238- 239)	XM_059132	ACGACTCCAAGAACAGCCAGGCCG (Forward) (SEQ ID NO 304) AAGGTAACATCGGCAGAGGCCAGC (Reverse) (SEQ ID NO 305)	GCTAGCATTTTTAAAAGCTGATGTCTTCACTGGGCACGGGGACTCACAC (0.94) (SEQ ID NO 306)
TMD0675 (SEQ ID NO 240- 241)	XM_059134	CGGCCAGGTACCAAAGCTCAGCTG (Forward) (SEQ ID NO 307) GCCAGATTCAGGAGGAATGGAAGAGAAC (Reverse) (SEQ ID NO 308)	TGATCTACTTTTTAAAAGGATCATGCTGGCTGCTGGGGATTTAGGATA (0.91) (SEQ ID NO 309) TGATAGTGATAAAAAAAGTGGCCAGATTTTGGTTATTTTTGAAATAAA (0.99) (SEQ ID NO 310) TATAGTGATATTTAAAGCCAGGGGTCTGGGTGAGATAACTGATGGAATGA (0.93) (SEQ ID NO 311) ATTGGAGGACTATAAAGGGGGTCTTAAAATGGTGCTAAGAAGCTGA (0.96) (SEQ ID NO 312) AGAGGGACTATAAAATGGTGCTAAGAAGCTGAGCTGA (0.96) (SEQ ID NO 313) GSEQ ID NO 313) GACATTCCACCCAAAAAATGCCACTGGATGAAGTCCCTTCCATTAA (0.92) (SEQ ID NO 314)

CODE	ACCN	PRIMFRS	ATLUMUM
TMD0677 (SEQ ID NO 242- 243)	XM_059140	TTGGGAGAGACTAGTGCACCTCAGCA (Forward) (SEQ ID NO 315) GAGCAATCCCTCTCGTGGCAGGT (Reverse) (SEQ ID NO 316)	AAAAGTGCTTTTAAACAGGGGGGGTGGGGGCTTATGAGAAGGGGACCA (1.00) (SEQ ID NO 317) CCATTTCTACTAAAAAAAAAAAAAAAAAAAAAAAAAAA
TMD0726 (SEQ ID NO 244- 245)	619650_MX	ACTTCCAAACATCTACAÁCTCCTCAGAGTCTCATT (Forward) (SEQ ID NO 321) TGCAGCACCATCATGTAAGGGACAA (Reverse) (SEQ ID NO 322)	TTTTTTAAACTATAAAAAGTGGGGATCAGAAAACACAGTCATAAGGGAAA (0.97) (SEQ ID NO 323) GTATATGCTATATATATCAGGATTCACTTTAATGGCATTGAGTTCCAGGA (0.98) (SEQ ID NO 324) ATAAACAATTTAAAAATTAGCCCACCATGGTGGTACACACCTGTCGTTCT (0.99) (SEQ ID NO 325) AAAAAGTGAAAAAAAAAAAGGTGAGGAGACTTTAACTTTCTGAAATATATT (0.92) (SEQ ID NO 326)
TMD0727 (SEQ ID NO 246- 247)	XM_059654 (related to)	CCAAGAAGCCGGGAGAAGTGGATG (Forward) (SEQ ID NO 327) TGACAGAGCTAGGCATATGAGCACTGGA (Reverse) (SEQ ID NO 328)	CTAAAGAGCTTATATATCAGCCTAAGAAAAGAAAACCAATAAGAAGTTGC (0.96) (SEQ ID NO 329)
TMD0739 (SEQ ID NO 248- 249)	ZM_059812	GCAGTTGGTTCAGAACCGAGATCACC (Forward) (SEQ ID NO 330) GGCAGATGGGGATACATTTATTCTCTGGG (Reverse) (SEQ ID NO 331)	ACTAAAAATACAAAAAGTAGCCGGGTATGGTGGTAGGCGCCTATAATCC (0.93) (SEQ ID NO 332) GGTAGGCGCCTATAATCCCAGCTACTTGGGAGGCTGAGGCAGGAGAATTG (0.92) (SEQ ID NO 333)
TMD0753 (SEQ ID NO 250- 251)	XM_059954	TCGGCTTGGAAATCAGAATGAGAAGG (Forward) (SEQ ID NO 334) TGCACAAAGAATGATTGCAGCAGTGAGTAG (Reverse) (SEQ ID NO 335)	AAAAGGCTTATATAAAAGGGTTTTGTTTTGTTTTTGAGACGGAGTT (0.97) (SEQ D NO 336) GGCCAACTTATAAAAAGGTTTATGTTTTTGTTCTGATAATTTCGTTTCT (0.91) (SEQ D NO 337) AAGTTAAAAAAGAACAGGCTACAAAGTTATAGGTTATAGGTTGT (0.96) (SEQ D NO 338)
TMDIIII (SEQ ID NO 252- 253)	NM_014386	GGGCGGTGTAGTGCAGGTCCG (Forward) (SEQ ID NO 339) CCTCCAGTTGCAGGGAATTCTGCC (Reverse) (SEQ ID NO 340)	AATTCAAATATTTAAAACGGACTGTCTCCTCTTCACAAAAGTCTAGATCT (0.92) (SEQ ID NO 341)
TMD1127 (SEQ ID NO 254- 255)	NM_054020	GGCTGTTGAGCAGCTTCATGTGC (Forward) (SEQ ID NO 342) CTCCTCTGGATGATCTGCCGCTTG (Reverse) (SEQ ID NO 343)	ATTGGGTGCATATATATAGGATAGTTAGCTCTTCTTGTTGAATTGATC (0.89) (SEQ D NO 344)

CLAIMS:

5

- A method of detecting an immune system cell, comprising:
 contacting a sample comprising cells with a polynucleotide specific for TMD0024
 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025
 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304
 (XM_060956), TMD0888 (XM_060957), or TMD0890 (XM_060959) of claim 28, under
- (XM_060956), TMD0888 (XM_060957), or TMD0890 (XM_060959) of claim 28, under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization.
- A method of claim 1, wherein said detecting is performed by:
 Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR,
 RACE PCR, or in situ hybridization.
 - 3. A method of detecting an immune system cell, comprising:
- contacting a sample comprising cells with a binding partner specific for a polypeptide coded for by TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), or TMD0890 (XM_060959) of claim 28, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding.
 - 4. A method of claim 3, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.
- 25 5. A method of delivering an agent to an immune cell, comprising:

contacting an immune cell with an agent coupled to binding partner specific for a polypeptide coded for by TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), or TMD0890

- 30 (XM_060959) of claim 28, whereby said agent is delivered to said cell.
 - 6. A method of claim 5, wherein the agent is a therapeutic agent or an imaging agent.

10

15

20

25

30

- 7. A method of claim 5, wherein the agent is cytotoxic.
- 8. A method of claim 5, wherein the binding partner is an antibody.

A method of modulating the maturation of an immune system cell, comprising:
 contacting said cell with an agent effective to modulate a gene, or polypeptide
 encoded thereby, selected from TMD0024 (XM_060945), TMD1779 (XM_060946),
 TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781
 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890
 (XM_060959) of claim 28, whereby the maturation of an immune cell is modulated.

- 10. A method of modulating interactions between lymphoid and non-lymphoid immune system cells, comprising:
- contacting said cells with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28, whereby the interaction is modulated.

11. A method of expressing a heterologous polynucleotide in immune system cells, comprising:

expressing a nucleic acid construct in immune system cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected from SEQ ID NOS 5, 10, 11, 16-19, 29-32, 37-39, 44-46, 51-54, and 59-62.

12. A method of treating an immune system disease, comprising:

administering to a subject in need thereof a therapeutic agent which is effective for regulating expression of a gene, or polypeptide encoded thereby, selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025

(XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28.

- 13. A method of claim 12, wherein said agent is an antibody or an antisense which is effective to inhibit translation of said gene.
 - 14. A method of diagnosing an immune disease associated with abnormal gene expression, or determining a subject's susceptibility to such disease, comprising:

assessing the expression of a gene, or polypeptide encoded thereby, selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28 in a tissue sample comprising immune system cells.

15. A method of claim 14, wherein assessing is:

measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene.

20 16. A method of claim 14, wherein said assessing detecting is performed by:

Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or *in situ* hybridization, and

using a polynucleotide probe having a sequence selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28, or a polynucleotide probe having 95% sequence identity or more to a sequence set forth in SEQ ID NOS 1, 6, 12, 20, 25, 33, 40, 47, or 55, effective specific fragments thereof, or complements thereto.

17. A method of assessing a therapeutic or preventative intervention in a subject having an

25

10

15

20

25

30

immune system disease, comprising,

determining the expression levels of a gene, or polypeptide encoded thereby, selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28 in a tissue sample comprising immune system cells.

- 18. A method of claim 17, further comprising assessing the expression levels of a plurality of said genes or polypeptides.
- 19. A method for identifying an agent that modulates the expression of a gene or polypeptide in the immune system gene complex, comprising,

contacting an immune system cell with a test agent under conditions effective for said test agent to modulate the expression of a gene selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28, or the biological activity of a polypeptide encoded thereby, in said immune system cell, and

determining whether said test agent modulates said gene or polypeptide.

- 20. A method of claim 19, wherein said agent is an antisense polynucleotide which is effective to inhibit translation of said gene or an antibody specific for said polypeptide.
- 21. A method of detecting polymorphisms in a gene in the immune system gene complex, comprising: comparing the structure of:

genomic DNA or RNA or cDNA or a polypeptide comprising all or part of a gene selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) of claim 28 with the structure of SEQ ID NOS 1, 6, 12, 20, 25, 33, 40, 47, or 55.

- 22. A method of claim 20, wherein said polymorphism is a nucleotide deletion, substitution, inversion, or transposition.
- 5 23. A method of identifying a genetic basis for an immune disease or disease-susceptibility, comprising:

determining the association of an immune disease or disease-susceptibility with a nucleotide sequence present in a genome comprising the gene complex of claim 28.

- 10 24. A method of claim 23, wherein determining is performed by producing a human-linkage map of said complex.
 - 25. A method of claim 23, wherein determining is performed by comparing the nucleotide sequences between normal subjects and subjects having an immune system disease.
 - 26. A non-human, transgenic mammal, or a cell thereof, whose genome comprises a functional disruption of a gene selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888
- 20 (XM_060957), and TMD0890 (XM_060959) of claim 28, or a mouse homolog thereof, and which has a defect in immune system function.
 - 27. A method of selecting a gene predominantly expressed in immune system cells from a database comprising polynucleotide sequences for genes, comprising:

WO 03/089583 PCT/US03/11497

-131-

displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for a gene selected from TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959), or complements to the polynucleotides sequence,

wherein said displayed sequences have been retrieved from said database upon it selection by a user.

- 28. A composition consisting essentially of the 1q22 immune gene complex, comprising TMD0024 (XM_060945), TMD1779 (XM_060946), TMD0884 (XM_060947), TMD0025 (XM_060948), TMD1780 (XM_089422), TMD1781 (XM_089421), TMD0304 (XM_060956), TMD0888 (XM_060957), and TMD0890 (XM_060959) genes, or a fragment thereof comprising at least two said genes.
- 15 29. A composition of claim 28, wherein said complex consists essentially of the chromosome region between STS markers SHGC-81033 and SHGC-145403, or a fragment thereof comprising at least two said genes.
- 30. A composition of claim 28, wherein said complex consists essentially of the
 20 chromosome region between STS markers SHGC-81033 and D1S3249, G15944,
 GDB:191077, or GDB:196442, or a fragment thereof comprising at least two said genes.
 - 31. A composition of claim 28, wherein said complex consists essentially of the chromosome region between STS markers RH118729 and D1S2577 or SHGC-145403, or a fragment thereof comprising at least two said genes.
- 32. A method of detecting an immune system cell, comprising:
 contacting a sample comprising cells with a polynucleotide specific for a XM_062147
 (SEQ ID NO 63) or XM_061676 (SEQ ID NO 69) of claim 59 under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization.

5

10

- 33. A method of claim 32, wherein said detecting is performed by: Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or in situ hybridization.
- 34. A method of detecting an immune system cell, comprising: contacting a sample comprising cells with a binding partner specific for a polypeptide coded for XM_062147 (SEQ ID NO 64) or XM_061676 (SEQ ID NO 70) of claim 59 under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding.
 - 35. A method of claim 34, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.
- 36. A method of delivering an agent to an immune cell, comprising:
 contacting an immune cell with an agent coupled to binding partner specific for XM_062147 (SEQ ID NO 64) or XM_061676 (SEQ ID NO 70) of claim 59, whereby said agent is delivered to said cell.
 - 37. A method of claim 36, wherein the agent is a therapeutic agent or an imaging agent.
 - 38. A method of claim 36, wherein the agent is cytotoxic.
 - 39. A method of claim 36, wherein the binding partner is an antibody.
- 40. A method of modulating the maturation of an immune system cell, comprising: contacting said cell with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59, whereby the maturation of an immune cell is modulated.
- 41. A method of modulating interactions between lymphoid and non-lymphoid immune system cells, comprising:

contacting said cells with an agent effective to modulate a gene, or polypeptide encoded thereby, selected from XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59, whereby the interaction is modulated.

5 42. A method of expressing a heterologous polynucleotide in immune system cells, comprising:

expressing a nucleic acid construct in immune system cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is SEQ ID NOS 65, 66, 72, 73, 74, or 75.

10

43. A method of treating an immune system disease, comprising:

administering to a subject in need thereof a therapeutic agent which is effective for regulating expression of a gene, or polypeptide encoded thereby, selected from XM_062147 (SEQ ID NO 63 or 64) or XM 061676 (SEQ ID NO 69 or 70) of claim 59.

15

- 44. A method of claim 43, wherein said agent is an antibody or an antisense which is effective to inhibit translation of said gene.
- 45. A method of diagnosing an immune disease associated with abnormal gene expression,or determining a subject's susceptibility to such disease, comprising:

assessing the expression of a gene, or polypeptide encoded thereby, selected from XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59 in a tissue sample comprising immune system cells.

25 46. A method of claim 45, wherein assessing is:

measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene.

30 47. A method of claim 45, wherein said assessing detecting is performed by:
Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR,

RACE PCR, or in situ hybridization, and

using a polynucleotide probe having a sequence selected from SEQ ID NOS 67, 68, 76, and 77.

5 48. A method of assessing a therapeutic or preventative intervention in a subject having an immune system disease, comprising,

determining the expression levels of a gene, or polypeptide encoded thereby, selected from XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59 in a tissue sample comprising immune system cells.

10

- 49. A method of claim 48, further comprising assessing the expression levels of a plurality of said genes or polypeptides.
- 50. A method for identifying an agent that modulates the expression of a gene or polypeptide
 in the immune system gene complex, comprising,

contacting an immune system cell with a test agent under conditions effective for said test agent to modulate the expression of XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59, or a polypeptide encoded thereby, in said immune system cell, and

- determining whether said test agent modulates said gene.
 - 51. A method of claim 50, wherein said agent is an antisense polynucleotide to a target polynucleotide sequence selected from SEQ ID NOS 63 or 69 and which is effective to inhibit translation of said gene.

25

52. A method of detecting polymorphisms in a gene in the immune system gene complex, comprising:

comparing the structure of: genomic DNA or RNA or cDNA comprising all or part of an allele of XM_062147 or XM_061676 with SEQ ID NOS 63 or 69 of claim 59.

30

53. A method of claim 52, wherein said polymorphism is a nucleotide deletion, substitution,

10

15

20

inversion, or transposition.

- 54. A non-human, transgenic mammal whose genome comprises a functional disruption of a gene represented by XM_062147 (SEQ ID NO 63) or XM_061676 (SEQ ID NO 69) of claim 59, and which has a defect in immune system function.
- 55. A mammalian immune system cell whose genome comprises a functional disruption of a gene represented by XM_062147 (SEQ ID NO 63) or XM_061676 (SEQ ID NO 69) of claim 59, and which has a defect in immune system function.
- 56. A mammalian cell of claim 55, wherein said cell is a mouse cell.
- 57. A non-human, transgenic mammal, or a cell thereof, comprising a gene operatively linked to an expression control sequence effective to express said gene in immune system, wherein said sequence is SEQ ID NOS 65, 66, 71, 72, 73, 74, or 75.
- 58. A method of selecting a gene predominantly expressed in immune system cells from a database comprising polynucleotide sequences for genes, comprising:

displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for XM_062147 (SEQ ID NO 63 or 64) or XM_061676 (SEQ ID NO 69 or 70) of claim 59, or complements to the polynucleotides sequence,

wherein said displayed sequences have been retrieved from said database upon selection by a user.

25 59. A composition comprising:

bone marrow specific genes consisting essentially of XM_062147 (SEQ ID NO 63 or 64) and XM_061676 (SEQ ID NO 69 or 70), or polypeptides thereof.

60. A method of detecting a kidney cell, comprising:

ontacting a sample comprising cells with a polynucleotide specific for a polynucleotide, or a naturally-occurring polymorphisms thereof, of claim 81 under conditions effective for said polynucleotide to hybridize specifically to said gene, and

detecting specific hybridization.

- 61. A method of claim 60, wherein said detecting is performed by:
 - Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR,
- 5 RACE PCR, or in situ hybridization.
 - 62. A method of detecting an kidney cell, comprising:

contacting a sample comprising cells with a binding partner specific for a polypeptide coded for by a polynucleotide of claim 81, or a naturally-occurring polymorphism thereof, under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding.

- 63. A method of claim 62, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.
- 64. A method of delivering an agent to a kidney cell, comprising:

contacting a kidney cell with an agent coupled to binding partner specific for polypeptide coded for by a polynucleotide of claim 81, or a naturally-occurring polymorphism thereof, whereby said agent is delivered to said cell.

20

15

10

- 65. A method of claim 64, wherein the agent is a therapeutic agent, a cytotoxic agent, or an imaging agent.
- 66. A method of claim 64, wherein the binding partner is an antibody.

25

67. A method of modulating a kidney cell, comprising:

contacting said cell with an agent effective to modulate a polynucleotide, or polypeptide encoded thereby, or a naturally-occurring polymorphism thereof, of claim 81, whereby the kidney cell is modulated.

30

68. A method of assessing kidney function, comprising:

15

20

25

detecting a polypeptide coded for by a polynucleotide of claim 81, or a naturallyoccurring polymorphism thereof, or fragments thereof, in a body fluid, whereby the amount of said polypeptide in said fluid is a measure of kidney function.

- 5 69. A method of claim 68, wherein said detecting is performed using an antibody which is specific for said polypeptide.
 - 70. A method of claim 69, wherein said detecting is performed by RIA, ELISA, or Western blot.
 - 71. A method of expressing a heterologous polynucleotide in kidney cells, comprising: expressing a nucleic acid construct in kidney cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected from SEQ ID NOS. 106, 109, 110, 113, 114, 117, 118, 121, 124, 125, 128-130, 133, 134, 137, 140, 141, 144, 147, 148, and 151.
 - 72. A method of diagnosing a kidney disease associated with abnormal gene expression, or determining a subject's susceptibility to such disease, comprising:
 - assessing the expression of a polynucleotide of claim 81, or a polypeptide encoded thereby, or naturally-occurring polymorphisms thereof, in a tissue sample comprising kidney cells.
 - 73. A method of claim 72, wherein assessing is:
 - measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene.
 - 74. A method of assessing a therapeutic or preventative intervention in a subject having a kidney disease, comprising,
- determining the expression levels of a polynucleotide of claim 81, a naturallyoccurring polymorphism thereof, or polypeptide encoded thereby, in a tissue sample

comprising kidney cells.

75. A method of claim 74, further comprising assessing the expression levels of a plurality of said genes or polypeptides.

5

10

20

76. A method for identifying an agent that modulates the expression of a polynucleotide or polypeptide selectively expressed in kidney cells, comprising,

contacting an kidney cell with a test agent under conditions effective for said test agent to modulate the expression of a polynucleotide of claim 81, or a naturally-occurring polymorphism thereof, or the biological activity of a polypeptide encoded thereby, in said kidney cell, and

determining whether said test agent modulates said gene or polypeptide.

- 77. A non-human, transgenic mammal whose genome comprises a functional disruption of a
 gene represented by a polynucleotide of claim 81, or a homolog thereof, and which has a
 defect in kidney function.
 - 78. A mammalian kidney cell whose genome comprises a functional disruption of a gene represented by a polynucleotide of claim 81, or a homolog thereof, and which has a defect in kidney function.
 - 79. A mammalian cell of claim 78, wherein said cell is a mouse cell.
- 80. A method of selecting a gene predominantly expressed in kidney cells from a database comprising polynucleotide sequences for genes, comprising:

displaying, in a computer-readable medium, a polynucleotide sequence, or a polypeptide encoded thereby, of claim 81, or complements to the polynucleotides sequence, wherein said displayed sequences have been retrieved from said database upon selection by a user.

5

10

81. A composition comprising two or more of the following polynucleotides expressed; selectively in kidney:

TMD0049 (XM_057351), TMD0190 (XM_087157), TMD0242 (XM_088369), TMD0335 (XM_089960), TMD0371, TMD0374, TMD0469 (XM_038736), TMD0719 (XM_059548), TMD0731 (XM_059703), TMD0785 (XM_060310), TMD0841 (XM_060623), TMD1114 (NM_019841), and/or TMD 1148 (XM_087108).

- 82. A method of detecting a pancreas cell, comprising:
- contacting a sample comprising cells with a polynucleotide specific for TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, of claim 113 under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization.
 - 83. A method of claim 82, wherein said detecting is performed by:
- Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or *in situ* hybridization.
 - 84. A method of detecting a pancreas cell, comprising:

contacting a sample comprising cells with a binding partner specific for a polypeptide coded for by TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, of claim 113 under conditions effective for said binding partner bind specifically to said polypeptide, and, detecting specific binding.

85. A method of claim 84, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.

30

25

86. A method of delivering an agent to a pancreas cell, comprising:

contacting a pancreas cell with an agent coupled to binding partner specific for

TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, of claim 113, whereby said agent is delivered to said cell.

- 87. A method of claim 86, wherein the agent is a therapeutic agent or an imaging agent.
- 88. A method of claim 86, wherein the agent is cytotoxic.
- 89. A method of claim 86, wherein the binding partner is an antibody.
- 90. A method of modulating a pancreas cell, comprising: contacting said cell with an agent effective to modulate TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, or the biological activity of a polypeptide encoded thereby, of claim 113, whereby the pancreas cell is modulated.
- 91. A method of assessing pancreas function, comprising: detecting a polypeptide coded for TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, or fragments thereof, in a body fluid, whereby the amount of said polypeptide in said fluid is a measure of pancreas function.
- 92. A method of claim 91, wherein said detecting is performed using an antibody which is specific for said polypeptide.
 - 93. A method of claim 91, wherein said detecting is performed by RIA, ELISA, or Western blot.
 - 94. A method of expressing a heterologous polynucleotide in pancreas cells, comprising: expressing a nucleic acid construct in pancreas cells, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is SEQ ID NOS 156-161, 166, 179, or 180.
 - 95. A method of diagnosing a pancreas disease associated with abnormal gene expression,

25

or determining a subject's susceptibility to such disease, comprising:

assessing the expression of TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, or polypeptide encoded thereby, of claim 113 in a tissue sample comprising pancreas cells.

5

96. A method of claim 95, wherein assessing is:

measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene.

10

15

20

97. A method of claim 95, wherein said assessing is performed by:

Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or *in situ* hybridization, and

using a polynucleotide probe having a sequence selected from SEQ ID NOS 154, 155, 164, 165, 169, 170, 173, 174, 177, 178, or a complement thereto.

98. A method of assessing a therapeutic or preventative intervention in a subject having a pancreas disease, comprising,

determining the expression levels of TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, or a polypeptide encoded thereby, of claim 113 in a tissue sample comprising pancreas cells.

99. A method of claim 98, further comprising assessing the expression levels of a plurality of said genes or polypeptides.

25

30

100. A method for identifying an agent that modulates the expression of TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, or the biological activity of a polypeptide encoded thereby, comprising,

contacting a pancreas cell with a test agent under conditions effective for said test agent to modulate the expression of TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785 of claim 113, or the biological activity of a polypeptide encoded thereby, in said

pancreas cell, and

5

determining whether said test agent modulates said gene or polypeptide.

- 101. A method of claim 100, wherein said agent is an antisense polynucleotide to a target polynucleotide sequence selected from SEQ ID NO 152, 162, 167, 171, or 175 and which is effective to inhibit translation of said gene.
 - 102. A method of detecting polymorphisms in TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, comprising,
- comparing the structure of: genomic DNA or RNA or cDNA comprising all or part of an allele of TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785, with SEQ ID NOS 152, 153, 162, 163, 167, 168, 171, 172, 175, or 176 of claim 113.
- 103. A method of claim 102, wherein said polymorphism is a nucleotide deletion,substitution, inversion, or transposition.
 - 104. A method of identifying a pancreatic disease or pancreatic disease-susceptibility, comprising:
- determining the association of a pancreatic disease or pancreatic disease-susceptibility
 with a nucleotide sequence present within the pancreatic gene complex of claim 113.
 - 105. A method of claim 104, wherein the pancreatic gene complex is from LOC160025-LOC119954.
- 106. A method of claim 104, wherein determining is performed by producing a human-linkage map of said complex.
 - 107. A method of claim 104, wherein determining is performed by comparing the nucleotide sequences between normal subjects and subjects having a pancreas disorder.
- 108. A non-human, transgenic mammal whose genome comprises a functional disruption of a gene represented by TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785

5

15

20

of claim 113, and which has a defect in pancreas function.

- 109. A mammalian pancreas cell whose genome comprises a functional disruption of a gene represented by TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785 of claim 113, and which has a defect in pancreas function.
- 110. A mammalian cell of claim 109, wherein said cell is a mouse cell.
- 111. A pancreas cell, comprising a gene operatively linked to an expression control sequence effective to express said gene in pancreas, wherein said sequence is SEQ ID NOS 156-161, 179, or 180.
 - 112. A method of selecting a gene predominantly expressed in pancreas cells from a database comprising polynucleotide sequences for genes, comprising:
 - displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for TMD0986, XM_061780, XM_061781, XM_061784, or XM_061785 of claim 113, or complements to the polynucleotides sequence,

wherein said displayed sequences have been retrieved from said database upon selection by a user.

- 113. A composition comprising: a pancreas specific gene consisting essentially of TMD0986, XM_061780, XM_061781, XM_061784, and/or XM_061785, or a polypeptide encoded thereby.
- 25 114. An isolated polynucleotide comprising a polynucleotide sequence which codes without interruption for a human TMD0986 having an amino acid sequence set forth in SEQ ID NO 153, or a complement thereto.
 - 115. An isolated polynucleotide comprising,
- a human TMD0986 polynucleotide sequence having 90% or more nucleotide sequence identity to the polynucleotide sequence set forth in SEQ ID NO 152 along its entire

length, which codes without interruption for human TMD0986, or a complement thereto, and which has G-protein coupling activity.

- 116. An isolated humansTMD0986 polypeptide comprising the amino acid sequence of a
 human TMD0986 as set forth in SEQ ID NO 153.
 - 117. An isolated human TMD0986 polypeptide consisting essentially of amino acids 1-117 of a human TMD0986 as set forth in SEQ ID NO 153.
- 10 118. An isolated polypeptide which is human TMD0986 having 90% or more amino acid sequence identity to the amino acid sequence set forth in SEQ ID NO 153, and which has protein binding activity.
 - 119. An antibody specific for an epitope selected from the polypeptide of claim 117.
 - 120. A method of detecting an retinal cell, comprising:

contacting a sample comprising cells with a polynucleotide specific for NM_013941 (SEQ ID NO 181), or a naturally-occurring polymorphisms thereof, of claim 142 under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization.

121. A method of claim 120, wherein said detecting is performed by:

Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or *in situ* hybridization.

122. A method of detecting an retinal cell, comprising:

contacting a sample comprising cells with a binding partner specific for a polypeptide coded for by NM_013941 (SEQ ID NO 182), or a naturally-occurring polymorphism thereof, of claim 142 under conditions effective for said binding partner bind specifically to said polypeptide, and

detecting specific binding.

15

20

25

- 123. A method of claim 122, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.
- 124. A method of delivering an agent to a retinal cell, comprising:
- contacting a retinal cell with an agent coupled to binding partner specific for by NM_013941 (SEQ ID NO 182), or naturally-occurring polymorphism thereof, of claim 142, whereby said agent is delivered to said cell.
- 125. A method of claim 124, wherein the agent is a therapeutic agent or an imaging agent.
- 126. A method of claim 124, wherein the agent is cytotoxic.
- 127. A method of claim 124, wherein the binding partner is an antibody.
- 15 128. A method of modulating a retinal cell, comprising:

contacting said cell with an agent effective to modulate NM_013941 (SEQ ID NO 181 or 182), or the biological activity of a polypeptide encoded thereby, of claim 142, whereby the retinal cell is modulated.

20 129. A method of diagnosing a retinal disease associated with abnormal gene expression, or determining a subject's susceptibility to such disease, comprising:

assessing the expression of NM_013941, a polymorphism thereof, or polypeptide encoded thereby, of claim 142 in a tissue sample comprising retinal cells.

- 25 130. A method of claim 129, wherein assessing is:
 - measuring expression levels of said gene, determining the genomic structure of said gene, determining the mRNA structure of transcripts from said gene, or measuring the expression levels of polypeptide coded for by said gene.
- 30 131. A method of claim 129, wherein said assessing detecting is performed by:

 Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR,

5

RACE PCR, or in situ hybridization, and

using a polynucleotide probe having a sequence selected from SEQ ID NOS 183 or 184, or a complement thereto.

5 132. A method of assessing a therapeutic or preventative intervention in a subject having an retinal disease, comprising,

determining the expression levels of NM_013941, a polymorphism thereof, or polypeptide encoded thereby, of claim 142 in a tissue sample comprising retinal cells.

- 133. A method of claim 132, further comprising assessing the expression levels of a plurality of said genes or polypeptides.
 - 134. A method for identifying an agent that modulates the expression of NM_013941 or the biological activity of a polypeptide encoded thereby, comprising,

contacting an retinal cell with a test agent under conditions effective for said test agent to modulate the expression of NM_013941 or a polymorphism thereof, of claim 142, or the biological activity of a polypeptide encoded thereby, in said retinal cell, and determining whether said test agent modulates said gene or polypeptide.

- 135. A method of claim 134, wherein said agent is an antisense polynucleotide to a target polynucleotide sequence selected from SEQ ID NO 181 and which is effective to inhibit translation of said gene.
- 136. A method of detecting polymorphisms in NM_013941, comprising:
 comparing the structure of: genomic DNA or RNA or cDNA comprising all or part
 of an allele of NM_013941, with SEQ ID NOS 181 or 182 of claim 142.
 - 137. A method of claim 136, wherein said polymorphism is a nucleotide deletion, substitution, inversion, or transposition.
- 30 138. A non-human, transgenic mammal whose genome comprises a functional disruption of a gene represented by NM_013941 (SEQ ID NO 181) of claim 142, and which has a defect in

10

retinal function.

- 139. A mammalian retinal cell whose genome comprises a functional disruption of a gene represented by NM_013941 (SEQ ID NO 181) of claim 142, and which has a defect in retinal function.
- 140. A mammalian cell of claim 139, wherein said cell is a mouse cell.
- 141. A method of selecting a gene predominantly expressed in retinal cells from a database comprising polynucleotide sequences for genes, comprising:

displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for NM_013941 (SEQ ID NO 181 or 182) of claim 142, or complements to the polynucleotides sequence,

wherein said displayed sequences have been retrieved from said database upon selection by a user.

142. A composition comprising:

a retinal specific gene consisting essentially of NM_013941 (SEQ ID NO 181 or 182), or a polypeptide encoded thereby.

20

25

5

10

15

143. A method of detecting a spleen cell, comprising:

contacting a sample comprising cells with a polynucleotide specific for TMD1030 (XM_166853) or TMD0621 (XM_166205) of claim 170 under conditions effective for said polynucleotide to hybridize specifically to said gene, and

detecting specific hybridization.

144. A method of claim 143, wherein said detecting is performed by:

Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or *in situ* hybridization.

30

145. A method of detecting a spleen cell, comprising:

contacting a sample comprising cells with a binding partner specific for a polypeptide

coded for by TMD1030 (XM_166853) or TMD0621 (XM_166205) of claim 170 under conditions effective for said binding partner bind specifically to said polypeptide, and detecting specific binding.

- 5 146. A method of claim 145, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.
- 147. A method of delivering an agent to a spleen cell, comprising:

 contacting a spleen with an agent coupled to binding partner specific for TMD1030

 (XM_166853) or TMD0621 (XM_166205) of claim 170, whereby said agent is delivered to said cell.
 - 148. A method of claim 147, wherein the agent is a therapeutic agent or an imaging agent.
- 15 149. A method of claim 148, wherein the agent is cytotoxic.
 - 150. A method of claim 147, wherein the binding partner is an antibody.
- 151. A method of modulating a spleen, immune, or reticuloendothelial cell, comprising:

 20 contacting said cell with an agent effective to modulate TMD1030 (XM_166853),

 TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), or the

 biological activity of a polypeptide encoded thereby, of claim 170, whereby the cell is

 modulated.
- 25 152. A method of assessing spleen function, comprising: detecting a polypeptide coded for by TMD1030 (XM_166853) or TMD0621 (XM_166205) of claim 170, or fragments thereof, in a body fluid, whereby the amount of said polypeptide in said fluid is a measure of spleen function.
- 30 153. A method of claim 152, wherein said detecting is performed using an antibody which is specific for said polypeptide.

- 154. A method of claim 152, wherein said detecting is performed by RIA, ELISA, or Western blot.
- 5 155. A method of expressing a heterologous polynucleotide in spleen cells, comprising: expressing a nucleic acid construct in spleen cell, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is SEQ ID NO 205-213.
- 10 156. A method of assessing a therapeutic or preventative intervention in a subject having a spleen or lymphoid disease, comprising,

determining the expression levels of TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), or a polypeptide encoded thereby, of claim 170 in a tissue sample comprising spleen, lymphoid, or reticuloendothelial cells.

- 157. A method of claim 156, further comprising assessing the expression levels of a plurality of said genes or polypeptides.
- 158. A method for identifying an agent that modulates the expression of TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), comprising,

contacting a spleen, lymphoid, or reticuloendothelial cell, with a test agent under conditions effective for said test agent to modulate the expression of TMD1030

25 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), of claim 170, and

determining whether said test agent modulates said gene.

159. A method of claim 158, wherein said agent is an antisense which is effective to inhibit translation of said gene.

5

- 160. A method for identifying an agent that modulates the expression of a polypeptide coded for by TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205), comprising,
- contacting a polypeptide coded for by TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205) of claim 170, with a test agent under conditions effective for said test agent to modulate said polypeptide, and determining whether said test agent modulates said polypeptide.
- 161. A method of detecting polymorphisms in comprising, comparing the structure of:

 10 genomic DNA or RNA or cDNA comprising all or part of an allele of TMD1030
 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621
 (XM_166205), with SEQ ID NOS 185, 187, 189, or 191 of claim 170.
- 162. A method of claim 161, wherein said polymorphism is a nucleotide deletion,substitution, inversion, or transposition.
 - 163. A method of identifying a genetic basis for a spleen, lymphoid, and/or reticuloendothelial disease or disease-susceptibility, comprising: determining the association of a spleen, lymphoid, and/or reticuloendothelial disease or disease-susceptibility with a nucleotide sequence present in the gene complex of claim 170.
 - 164. A method of claim 163, wherein determining is performed by producing a human-linkage map of said complex.
- 25 165. A method of claim 163, wherein determining is performed by comparing the nucleotide sequences between normal subjects and subjects having a spleen, lymphoid, and/or reticuloendothelial disease.
- 166. A non-human, transgenic mammal, or a cell thereof. whose genome comprises a functional disruption of a gene represented by TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205) of claim 170, and

PCT/US03/11497

which has a defect in spleen, lymphoid, and/or reticuloendothelial disease function.

- 167. A mammalian cell of claim 166, wherein said cell is a mouse cell.
- 5 168. A spleen, lymphoid, and/or reticuloendothelial cell, comprising a gene operatively linked to an expression control sequence effective to express said gene in spleen, lymphoid, and/or reticuloendothelial, wherein said sequence is SEQ ID NO 205-213.
 - 169. A method of selecting a gene predominantly expressed in spleen, lymphoid, and/or reticuloendothelial cells from a database comprising polynucleotide sequences for genes, comprising:

displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), or TMD0621 (XM_166205) of claim 170, or complements to the polynucleotides sequence, wherein said displayed sequences have been retrieved from said database upon selection by a user.

- 170. A composition consisting essentially of the 11q12.2 spleen gene complex, comprising TMD1030 (XM_166853), TMD1029 (XM_166854), TMD1028 (XM_166855), and TMD0621 (XM_166205).
- 171. A composition of claim 170, wherein said complex consists essentially of the chromosome region between STS markers G62658 and SHGC-154002.
- 172. A method of detecting a pancreas cell, comprising:
 contacting a sample comprising cells with a polynucleotide specific TMD0077,
 TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530,
 TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677,
 TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127 of claim 199
 under conditions effective for said polynucleotide to hybridize specifically to said gene, and detecting specific hybridization.

10

15

10

- 173. A method of claim 172, wherein said detecting is performed by: Northern blot analysis, polymerase chain reaction (PCR), reverse transcriptase PCR, RACE PCR, or in situ hybridization.
- 5 174. A method of detecting a pancreas cell, comprising: contacting a sample comprising cells with a binding partner specific for a polypeptide coded for by TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or
- TMD1127 of claim 199 under conditions effective for said binding partner bind specifically to said polypeptide, and

detecting specific binding.

15 175. A method of claim 174, wherein said detecting is performed by: immunocytochemistry, immunoprecipitation, or Western blot.

176. A method of delivering an agent to a pancreas cell, comprising:

- contacting a pancreas with an agent coupled to binding partner specific for 20 TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127 of claim 199, whereby said agent is delivered to said cell.
- 25 177. A method of claim 176, wherein the agent is a therapeutic agent or an imaging agent.
 - 178. A method of claim 176, wherein the agent is cytotoxic.
 - 179. A method of claim 176, wherein the binding partner is an antibody.
 - 180. A method of modulating a pancreas, immune, or reticuloendothelial cell, comprising:

PCT/US03/11497 -153-

contacting said cell with an agent effective to modulate TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, or the biological activity of a polypeptide encoded thereby, of claim 199, whereby the cell is modulated.

181. A method of assessing pancreas function, comprising:

detecting a polypeptide coded for by TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127 of claim 199, or fragments thereof, in a body fluid, whereby the amount of said polypeptide in said fluid is a measure of pancreas function. 182. A method of claim 181, wherein said detecting is performed using an antibody which is specific for said polypeptide.

15

10

5

183. A method of claim 181, wherein said detecting is performed by RIA, ELISA, or Western blot.

20

expressing a nucleic acid construct in pancreas cell, said construct comprising a promoter sequence operably linked to said heterologous polynucleotide, wherein said promoter sequence is selected SEQ ID NO 258, 261, 262, 265-267, 270-272, 275, 278, 279, 282-284, 287, 290-293, 296, 297, 300, 303, 306, 309-314, 317-320, 323-326, 329, 332-333, 336-338, 341, and 344.

184. A method of expressing a heterologous polynucleotide in pancreas cells, comprising:

25

30

185. A method of assessing a therapeutic or preventative intervention in a subject having a pancreas or lymphoid disease, comprising,

determining the expression levels of TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, or a polypeptide encoded thereby, of claim 199 in a tissue sample comprising pancreas, lymphoid, or reticuloendothelial cells.

186. A method of claim 185, further comprising assessing the expression levels of a plurality of said genes or polypeptides.

5

187. A method for identifying an agent that modulates the expression of TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, comprising,

10

contacting a pancreas, lymphoid, or reticuloendothelial cell, with a test agent under conditions effective for said test agent to modulate the expression of TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, of claim 199, and determining whether said test agent modulates said gene.

15

188. A method of claim 187, wherein said agent is an antisense which is effective to inhibit translation of said gene.

20

189. A method for identifying an agent that modulates the expression of a polypeptide coded for by TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, comprising,

25

contacting a polypeptide coded for by TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127 of claim 199, with a test agent under conditions effective for said test agent to modulate said polypeptide, and

determining whether said test agent modulates said polypeptide.

with SEQ ID NOS of Table 23 of claim 199.

- 190. A method of claim 189, wherein said test agent is an antibody.
- 191. A method of detecting polymorphisms in comprising, comparing the structure of : genomic DNA or RNA or cDNA comprising all or part of an allele of TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127.
- 10 192. A method of claim 191, wherein said polymorphism is a nucleotide deletion, substitution, inversion, or transposition.
 - 193. A method of identifying a genetic basis for a pancreas disease or disease-susceptibility, comprising: determining the association of a pancreas disease or disease-susceptibility with a gene of claim 199.
 - 194. A method of claim 193, wherein determining is performed by producing a human-linkage map of said gene.
- 20 195. A method of claim 193, wherein determining is performed by comparing the nucleotide sequences between normal subjects and subjects having a pancreas disease.
 - 196. A non-human, transgenic mammal, or a cell thereof. whose genome comprises a functional disruption of a gene represented by TMD0077, TMD0233, TMD0256, TMD0258,
- TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, of claim 199, and which has a defect in pancreas, lymphoid, and/or reticuloendothelial disease function.
- 30 197. A mammalian cell of claim 196, wherein said cell is a mouse cell.
 - 198. A method of selecting a gene predominantly expressed in pancreas tissue from a

5

5

10

displaying, in a computer-readable medium, a polynucleotide sequence or polypeptide sequence for TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127, of claim 199, or complements to the polynucleotides sequence, wherein said displayed sequences have been retrieved from said database upon selection by a user.


199. A composition comprising genes and/or polypeptide which are expressed predominantly in pancreas tissue comprising:

TMD0077, TMD0233, TMD0256, TMD0258, TMD0267, TMD0271, TMD0290, TMD0530, TMD0574, TMD0608, TMD0639, TMD0645, TMD0674, TMD0675, TMD0677, TMD0726, TMD0727, TMD0739, TMD0753, TMD1111, and/or TMD1127.

WO 03/089583 PCT/US03/11497


ST S) 최	RNA∑)		Acen	locus	Gene code	
55920 K	RH79988 155 ISHGC-545185	920K	XM_	173381	LOC254079		
55930 K 55940 K 55950 K	! RH93840 155 - R004X45 155 E SHGC-82759 O183312 155 RH66323 155	940K 950K	хм_	086610	CDID		
55970 K- 55980 K- 55990 K- 56000 K-	155	5970K 5980K 5990K					
56020K	RH120517156		. XM_	048792	CDIA		
560 40 K- 560 50 K- 560 60 K- 560 70 K-	D1 53359 150	6040K 6050K 6060K 6070K	NM_	001765.1	CDIC		
560 80K	- SHGC-12795	6030K	NM	001764.1	CD1B		
56100K 56110K 56120K	 AL009528 ¹⁵ = RH70821 15	6100K4 6110K	NM_	030893.1	CDIE		
561 30K	15	6134K			. 051202(0	TMD0024	
561 50K 561 60K	1	6160K	• • •	060945	LOC128360	TMD0024	
561 76 K	15	6170K- 6180K- 6199A-		060946	LOC128361	TMD1779 TMD0884	
561 90 K 5620 CK 5621 OK	15	6200 K	·	060947	LOC128362 LOC128363	TMD0025	
56220K	SHGC-15585	6230 K	XM	_060948	EOC 128303		Dia 1
5624 0 K	15	6240K- 6250K- 6260K-	, XM	089422	LOC164169	TMD1780	Fig. 1
5626 K 5627 K 5628 K	15	6270K	/ XM	_089421	LOC149631	TMD1781	
56290K- 56300K-	,	6290K		_060949 _089420	LOC128364 LOC164168		
56310K- 56320K- 56230K-) 5	6310K 66320K 66330K	XM	_060950	LOC128365		
56340K	0153249 15	6340K	XM.	_060951	LOC128366		
56360 K 56370 K	GDB:191075 GDB:196446 	6370K	ХМ	_060952	1.OC128367		
56380K- 56390K- 56400K-	. RM65132	56390K	XM	060953	LOC128368		
56410K	, 15	56410K 56420K	NM	_003126.1	SPTAI		
56430K	GDB:1815						
56450K- 56460K- 56470K-	- RH118729	56450K 56460K 56470K	-· XM	_060954	LOC128369		
56+80K- 56+90K-	1	56480K		_ _060955	LOC128370		
56500K 56510K	1	56500K			1.OC128371	TMD0304	
56520K 56530K 56540K	•	56520K 56530K 56540K		_ i_06 0 957	LOC128372	TMD0888	
36550K		56550K	· ₋ XM	1_060958	LOC128373		
36570K 36580K	1 - D1\$2577 1	56570K	XM	1_060959	LOC128374	TMD0890	
36590K 36600K	-SHGC-1454	200011	: . NM	1_002432.1	MNDA	•	
56610K- 56630K-	1	56610K 56620K 56630K			.:		
56650K	1	56640K	Ì				
56660K 56670K	1	56660K	XM	1_089418	LOC149629		
5660•K 5669•K 5670•K	,	156680K- 156690K- 156700K-	I I				
5671 0K-	1	156720K	-				

Fig. 2



WO 03/089583 PCT/US03/11497

Fig. 3



XM_062147



XM_061676




FIG. 4

TMD0719

 \sim

TMD0841


 ∞

<u>&</u>


WO 03/089583 PCT/US03/11497



Fig. 6



FigIA



FigtB

XM_061779	XM_061780	XM_061781	XM_061784	XM_061785	
314	318	323	149	311	Fig. 8

WO 03/089583 PCT/US03/11497




Fig. 9A






Fig. 9B




Fig. 10

RNA	accession locus	gene code
55+5H	XM_166847 LOC219495	
	XM_166848 LOC219496	
 	XM_166849 LOC219497	TMD1799
55.6H	XM_166850 LOC245817	
	XM_166851 LOC219498	
55+7H	XM_166852 LOC219521	
	XM_166853 LOC219522 XM_166854 LOC219523 XM_166855 LOC219524	TMD1030 TMD1029 TMD1028
85. 81.	XM_166205 LOC219525	TMD0621
		
55.9#	XM_169164 LOC219526 XM_166856 LOC219527	TMD1027
<u> </u>	XM_166857 LOC219511 NM_005161.1 AGTRL1	· · · · · · · · · · · · · · · · · · ·

ig. 11

FIG. 12A

9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10 11 12 13 14 15 16 17 18 19 20 21 22 24		9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	17 18 19 20
TMD0077 = 1 2 3 4 5 6	TMD0233	TMD0256	TMD0258	TMD0267	1 2 3 4 5 6 TMD0271	1 2 3 4 5 6 TMD0290	1 2 3 4 5 6

FIG. 12B

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24
MII MIII	DI!			Mi		
TMD0530	TMD0608	TMD0639	TMD0645	TMD0674	TMD0675	



	24		24	24	24	24	24		24
	23		23	23	23	23	23		23.
_	22		22	22		22	22		22
	21		7	7	21	212	20 21		21
ł	20		20	20	20	20	20		20
	19		19	19	19	19	19	•	19
	18		& .	18	18	17 18			200
j.	17		17	17	17	. 11	17		17
-	16		91.	16	16	16	16		16
1	15	1	5.	15	15	2	15		15
**************************************	14		13 14	4	4	7	4	· ·	4
	13			2	13	13	13		13
	12		10 11 12	12	12	12	. 12	:	12
•			=	=	=	=	1=	:	
	10 11		10	0	0	10	10	;	10
1			0	6	6	6	6		0
1	6		∞	∞	∞	∞	∞		∞
•	∞		7	~ : \			7 8	į	_
	7		9	9	7	9	9	,	9
	9 9		v .	8	9	2	5		S
			- 4 2	4	4	; 4	4		4
	4		ന	6	7	m	8	•	m
	m.		8	8	8	7	7		7
	7				7	-		•	
:	-		1111	111	!				
TMD0677		TMD0726	TMD0727	TMD0739		TMD0753	TMD1111	TMD1127	

16U 200 PCT FINAL.ST25 · SEQUENCE LISTING

						~.	-205		J1 J 1 .							
<1	10>	Ori	Gene	Tec	hnol	ogies	s, I	nc								
<1	20>	TIS	SUE :	SPEC	IFIC	GENI	ES A	ND GI	ENE (CLUS	rers					
<1	30>	1 6 U	200	PCT												
	50> 51>		60/3° 2-04-		69											
	50> 51>		50/41 2-09-		32											
	50> 51>		50/42 2-11-		36											
	50> 51>		50/37 2-04-		23											
	50> 51>		50/37 2-05-		8											
	50> 51>		50/36 2-05-		56											
	50> 51>		50/40 2-08-		8											
<16	60>	344														
<17	70>	Pate	ntIn	ver	sion	3.1										
<21	10> 11> 12> 13>	1 795 DNA Homo	sap	iens												
<22 <22 <22 <22	21> 22>	CDS (1).	. (79	5)												
<40 atg Met 1	00> gag Glu	l cgg Arg	gtc Val	aat Asn 5	gag Glu	act Thr	gtg Val	gtg Val	aga Arg 10	gag Glu	gtc Val	atc Ile	ttc Phe	ctc Leu 15	ggc Gly	48
ttc Phe	tca Ser	tcc Ser	ctg Leu 20	gcc Ala	agg Arg	ctg Leu	cag Gln	cag Gln 25	ctg Leu	ctc Leu	ttt Phe	gtt Val	atc Ile 30	ttc Phe	ctg Leu	96
ctc Leu	ctc Leu	tac Tyr 35	ctg Leu	ttc Phe	act Thr	ctg Leu	ggc Gly 40	acc Thr	aat Asn	gca Ala	atc Ile	atc Ile 45	att Ile	tcc Ser	acc Thr	144
att Ile	gtc Val 50	ctg Leu	gac Asp	agg Arg	gcc Ala	ctt Leu 55	cat His	atc Ile	ccc Pro	atg Met	tac Tyr 60	ttc Phe	ttc Phe	ctt Leu	gcc Ala	192
atc Ile 65	ctc Leu	tct Ser	tgc Cys	tct Ser	gag Glu 70	att Ile	tgc Cys	tac Tyr	acc Thr	ttc Phe 75	atc Ile	att Ile	gta Val	ccc Pro	aag Lys 80	240
atg Met	ctg Leu	gtt Val	gac Asp	ctg Leu 85	ctg Leu	tcc Ser	cag Gln	aag Lys	aag Lys 90	acc Thr	att Ile	tct Ser	ttc Phe	ctg Leu 95	ggc Gly	288
tgt Cys	gcc Ala	atc Ile	caa Gln 100	atg Met	ttt Phe	tcc Ser	ttc Phe	ctc Leu 105	ttc Phe	ctt Leu	ggc Gly	tgc Cys	tct Ser 110	cac His	tcc Ser	336
ttt Phe	ctg Leu	ctg Leu 115	gca Ala	gtc Val	atg Met	ggt Gly	tat Tyr 120	gat Asp	cgt Arg	tac Tyr	ata Ile	gcc Ala 125	atc Ile	tgt Cys	aac Asn	384
cca Pro	ctg Leu 130	cgc Arg	tac Tyr	tca Ser	gtg Val	cta Leu 135	atg Met	gga Gly	cat His	GJ À ààà	gtg Val 140	tgt Cys	atg Met	gga Gly	cta Leu	432

										100	200	PCT	FINA	L.51	25		
								ttc Phe								48	0
								tat Tyr								52	8
								ctc Leu 185								57	6
								atg Met								62	4
								tcc Ser								67	2
								gga Gly								72	0
								tat Tyr								76	8
					ctg Leu			taa								79	5
<210 <211 <212 <213	> 2 ?> F	264 PRT	sapi	ens													
< 400)> 2	2															
Met 1	Glu	Arg	Val	Asn 5	Glu	Thr	Val	Val	Arg 10	Glu	Val	Ile	Phe	Leu 15	Gly		
Phe	Ser	Ser	Leu 20	Ala	Arg	Leu	Gln	Gln 25	Leu	Leu	Phe	Val	Ile 30	Phe	Leu		
Leu	Leu	Tyr 35	Leu	Phe	Thr	Leu	Gly 40	Thr	Asn	Ala	Ile	Ile 45	Ile	Ser	Thr		
Ile	Val 50	Leu	Asp	Arg	Ala	Leu 55	His	Ile	Pro	Met	Туг 60	Phe	Phe	Leu	Ala		
Ile 65	Leu	Ser	Cys	Ser	Glu 70	Ile	Суѕ	Туr	Thr	Phe 75	Ile	lle	Val	Pro	Lys 80		
Met	Leu	Val	Asp	Leu 85	Leu	Ser	Gln	Lys	Lys 90	Thr	Ile	Ser	Phe	Leu 95	Gly		
Cys	Ala	Ile	Gln 100	Met	Phe	Ser	Phe	Leu 105	Phe	Leu	G1 y	Cys	Ser 110	His	Ser		
Phe	Leu	Leu 115	Ala	Val	Met	Gly	Tyr 120	Asp	Arg	Tyr	Ile	Ala 125	Ile	Cys,			
Pro	Leu 130	Arg	Tyr	Ser	Val	Leu 135	Met	Gly	His	Gly	Val 140	Cys	Met	Gly	Leu		
Val 145	Ala	Ala	Ala	Cys	Ala 150	Cys	Gly	Phe	Thr	Val 155	Ala	Gln	Ile	Ile	Thr 160		

Ser Leu Val Phe His Leu Pro Phe Tyr Ser Ser Asn Gln Leu His His 165 170 175Phe Phe Cys Asp Ile Ala Pro Val Leu Lys Leu Ala Ser His His Asn His Phe Ser Gln Ile Val Ile Phe Met Leu Cys Thr Leu Val Leu Ala Ile Pro Leu Leu Leu lle Leu Val Ser Tyr Val His Ile Leu Ser Ala Ile Leu Gln Phe Pro Ser Thr Leu Gly Val Ile Ala Lys Arg Lys Phe His Asn Ser Asp Asp Phe Ser His Tyr Asn Ser Phe Gln Asp Pro Pro 250 Val Asn Lys Ser Leu Leu Ile Asp 260 <210> <211> 32 <212> DNA <213> Homo sapiens <400> 3 ggtcaatgag actgtggtga gagaggtcat ct 32 <210> 4 <211> 31 <212> DNA <213> Homo sapiens <400> 4 ctatcactcc cagtgtggaa ggaaactgaa g 31 <210> 5 <211> 50 <212> DNA <213> Homo sapiens <400> 5 ctctttcaga tttaaatggg ccagacttag ttttatgtgg tgcagacatt 50 <210> 6 <211> 807 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (1)..(807) <223> <400> 6 atg gcc gtt att cgc ttc agc tgg act ctc cac act ccc atg tat ggc Met Ala Val Ile Arg Phe Ser Trp Thr Leu His Thr Pro Met Tyr Gly 48 ttt cta ttc atc ctt tca ttt tct gag tcc tgc tac act ttt gtc atc Phe Leu Phe Ile Leu Ser Phe Ser Glu Ser Cys Tyr Thr Phe Val Ile 96 atc cct cag ctg ctg gtc cac ctg ctc tca gac acc aag acc atc tcc Ile Pro Gln Leu Leu Val His Leu Leu Ser Asp Thr Lys Thr Ile Ser 144

ttc atg gcc Phe Met Ala 50	tgt gcc ac Cys Ala Th	cc cag ctg or Gln Leu 55	ttc ttt Phe Phe	ttc ctt Phe Leu 60	ggc ttt Gly Phe	gct tge Ala Cy	192 s
acc aac tgc Thr Asn Cys 65		e Ala Val					
att tgt cac Ile Cys His							
ttg gag ttg Leu Glu Leu						Ala Le	
gtg gcc acc Val Ala Thr 115	Asn Leu I						
gtt aac cac Val Asn His 130					Lys Let		
act gac acc Thr Asp Thr 145	His Val Ly						u
gta att atg Val Ile Met							
gtt aac acc Val Asn Thr						Ala Ph	
gtc acc tgt Val Thr Cys 195	Ala Ser Hi		Val Val				
gcc tct ato Ala Ser Ile 210					Ala Se		
gat cag ttg Asp Gln Leu 225		al Thr Tyr					מ
cct ctt gtc Pro Leu Val				Val Lys			
aga gtt ctt Arg Val Leu							807
<210> 7 <211> 268 <212> PRT <213> Homo	sapiens						
<400> 7							
Met Ala Val	. Ile Arg Pi 5	ne Ser Trp	Thr Leu 10	His Thr	Pro Me	Tyr G1; 15	У
Phe Leu Phe	e Ile Leu Se 20	er Phe Ser	Glu Ser 25	Cys Tyr	Thr Pho	val Il	e ·
Ile Pro Glr 35	ı Leu Leu Va	al His Leu 40	Leu Ser	Asp Thr	Lys Th	r Ile Se	r
Phe Met Ala 50	ı Cys Ala Ti	or Gln Leu 55	Phe Phe	Phe Leu 60	Gly Ph	e Ala Cy	s

Thr 65	Asn	Cys	Leu	Leu	Ile 70	Ala	Val	Met	Gly	Туг 75	Asp	Arg	Tyr	Val	Ala 80		
Ile	Cys	His	Pro	Leu 85	Arg	Туr	Thr	Leu	11e 90	Ile	Asn	Lys	Arg	Leu 95	Gly		
Leu	Glu	Leu	Ile 100	Ser	Leu	Ser	Gly	Ala 105	Thr	Gly	Phe	Phe	Ile 110	Ala	Leu		
Val	Ala	Thr 115	Asn	Leu	lle	Cys	Asp 120	Met	Arg	Phe	Cys	Gly 125	Pro	Asn	Arg		
Val	Asn 130	His	Tyr	Phe	Cys	Asp 135	Met	Ala	Pro	Val	11e 140	Lys	Leu	Ala	Cys		
Thr 145	Asp	Thr	His	Val	Lys 150	Glu	Leu	Ala	Leu	Phe 155	Ser	Leu	Ser	Ile	Leu 160		
Val	Ile	Met	Val	Pro 165	Phe	Leu	Leu	Ile	Leu 170	Ile	Ser	Tyr	Gly	Phe 175	Ile		
Val	Asn	Thr	Ile 180	Leu	Lys	Ile	Pro	Ser 185	Ala	Glu	Gly	Lys	Lys 190	Ala	Phe		
Val	Thr	Cys 195	Ala	Ser	His	Leu	Thr 200	Val	Val	Phe	Val	His 205	Tyr	Gly	Cys		
Ala	Ser 210	Ile	Ile	Туr	Leu	Arg 215	Pro	Lys	Ser	Lys	Ser 220	Ala	Ser	Asp	Lys		
Asp 225	Gln	Leu	Val	Ala	Val 230	Thr	Tyr	Thr	Val	Val 235	Thr	Pro	Leu	Leu	Asn 240		
Pro	Leu	Val	Tyr	Ser 245		Arg	Asn	Lys	Glu 250	Val	Lys	Thr	Ala	Leu 255	Lys		
Arg	Val	Leu	Gly 260	Met	Pro	Val	Ala	Thr 265	Lys	Met	Ser				-		
<210 <211 <211 <211	1 > 2	8 25 DNA Homo	sap	iens													
<400 cca		B ctc 1	tcaga	acac	ca a	gacc											25
<210 <210 <210 <210	1> : 2> :	9 27 DNA Homo	san	i ens													
<40)>	9			a.ta.	-+	20										27
		taa 1	ccac	cayy	ac y	Lya	99							,		•	.,
<210 <210 <210 <210	1> 2>	10 50 DNA Homo	sap	iens													
<400 gag		10 aaa t	tata	taaa	ga go	gtate	gttc	a at	gcaa	catg	tta	aatg	_				50

<210> 11 <211> 50 <212> DNA <213> Home	o sapiens					
<400> 11 actccttaga	taaaaaaggg	cagatttatt	aaagaaccct	gatttaatca		50
<210> 12 <211> 4982 <212> DNA <213> Homo	? O sapiens					
<220> <221> CDS <222> (201 <223>	9)(2960)					
<400> 12 gtactccttc	agaatcagag	aattccagct	tccatggttt	acattattca	tcatattcag	60
tcaagtgagg	gcctagtggc	ggttaaaggt	tgattagttg	aaagaagatt	caaatgaaag	120
tcttttggga	aagcaatgag	gcaaggctaa	gcaatgacca	taagtttaga	tttcctcatt	180
gttttgaata	gacaggaaat	catttgtcca	gaaggaggta	ttatgtaggg	aaacttttac	240
ctttctgtat	ataaaaacat	ataactaata	cacacacact	catacacaaa	tatcaatgga	300
ggtatacatt	gtgtttactt	tttctatgtt	tatgtacaat	agtaatatct	ttatagttat	360
actaacgtta	ttaaaataag	taattatatt	aactaagttt	aggaccagtt	tctagtaagt	420
aagaaagaaa	aaaaatcatc	tccaaattct	atgaatagat	ataatgaatt	tcaagaatgc	480
ctgatgaatt	aacttaggat	tcaggaaaca	aaaaaagttg	ctattgaata	gaaaaatgga	540
aaagtaacag	caacaaaatt	ctggtagcag	atgccaataa	tttcccaaga	caaaatgatg	600
tagtaacttc	agaagtatat	aaatgaagac	tggataccag	caagacatac	tggatgattt	660
tgtatccaga	tagtgctttt	tttacttatt	aggttgggtt	attgaaaaat	gttccagtga	720
aaaaaattag	gcctaagatg	attttagaaa	taatttgtaa	tggcagtttg	caaaatattt	780
ttagtggcag	aatgttcaaa	agaaatctta	ttaacataac	aacatacaaa	agatacaaag	840
cctatggttt	acagcaggag	aggggaaact	ggcaaaattc	ccaagtgtgc	cattctctct	900
cacactctgt	agcaagctct	gtcatttcta	caaaactctt	atttctctga	gtttctccaa	960
gttagctcag	catggaaaag	tgaagtgtgt	tacaaaatgc	cacaaagtca	gtcatctctc	1020
tttaccaccc	tggtgactat	tctcttcctg	aaagaagaat	ttttttcttt	atactaatgc	1080
actaatgtta	tttattttta	ttttatttta	tttatttatt	tttgagacag	attctcactg	1140
tgtcacccag	tctggagtgc	agaggcacaa	tcttggctca	ctgcaacctc	cgcctcccgg	1200
gctcaagtga	atctcatgcc	tcagcctccc	gagtagctgg	gattacaggt	gtgtgctgcc	1260
atacctggct	aatttttgta	cttttagtaa	agaccaggtt	ttgccatgtt	gccgaggctg	1320
gtcttgaacc	cctggcctca	agcaatccac	ccaccttggc	ttctcaaagt	gctgggatta	1380
caggtgtgag	ccaccacatc	tggctaatgt	tatttttgt	ttcactgttg	actcaatgtt	1440
tcaacttgtg	gaacttccaa	tagtatttct	tattgttccc	ttggagatat	aaaaagttcc	1500
cagtaaatag	atgtgtgctc	acatctttac	ttagagacca	tggaatactt	tatctccttt	1560
ctcatttcat	ggttggataa	actgaagtcc	acatgattat	gtctgaatat	tattcattct	1620
ttcgttctat	attctgatca	gcttcaggta	gctgaagtta	acgttttcca	ctttggagag	1680
tgagttgcct	tgggtttata	gtaagtgaca	aaaacaacaa	tctctctgtt	acataagaag	1740

gaaaactatt agcaaatttc ctaatccttg gtcagagaga taacctgttc ttcacattag	1800
agaaggcctc caaactggct atcagttatt cttttgcata ttttgcctaa ttcttctttt	1860
agcaggcatt ttaatggggg aatgaagaat tccatcaaat atctggaaat gcctgccacc	1920
tgcaaacttt gtgtgaaatt tcccgtacat ttccactctc ctttctggat cctggtttct	1980
acctetytee etgaetetee tttatagaag tyetetee atg gag caa yte aat aag Met Glu Gln Val Asn Lys 1 5	2036
act gtg gtg aga gag ttc gtc gtc ctc ggc ttc tca tcc ctg gcc agg Thr Val Val Arg Glu Phe Val Val Leu Gly Phe Ser Ser Leu Ala Arg 10 15 20	2084
ctg cag cag ctg ctc ttt gtt atc ttc ctg ctc ctc tac ctg ttc act Leu Gln Gln Leu Leu Phe Val Ile Phe Leu Leu Tyr Leu Phe Thr 25 30 35	2132
ctg ggc acc aat gca atc atc att tcc acc att gtg ctg gac aga gcc Leu Gly Thr Asn Ala Ile Ile Ile Ser Thr Ile Val Leu Asp Arg Ala 40 45 50	2180
ctt cat act ccc atg tac ttc ttc ctt gcc atc ctt tct tgc tct gag Leu His Thr Pro Met Tyr Phe Phe Leu Ala Ile Leu Ser Cys Ser Glu 55 65 70	2228
att tgc tat acc ttt gtc att gta ccc aag atg ctg gtt gac ctg ctg Ile Cys Tyr Thr Phe Val Ile Val Pro Lys Met Leu Val Asp Leu Leu 75 80 85	2276
tcc cag aag aag acc att tct ttc ctg ggc tgt gcc atc caa atg ttt Ser Gln Lys Lys Thr Ile Ser Phe Leu Gly Cys Ala Ile Gln Met Phe 90 95 100	2324
Ser Phe Leu Phe Phe Gly Ser Ser His Ser Phe Leu Leu Ala Ala Met 105	2372
ggc tat gat cgc tat atg gcc atc tgt aac cca ctg cgc tac tca gtg Gly Tyr Asp Arg Tyr Met Ala Ile Cys Asn Pro Leu Arg Tyr Ser Val 120 125 130	2420
ctc atg gga cat ggg gtg tgt atg gga cta atg gct gct gcc tgt gcc Leu Met Gly His Gly Val Cys Met Gly Leu Met Ala Ala Ala Cys Ala 135 140 145 150	2468
tgt ggc ttc act gtc tcc ctg gtc acc acc tcc cta gta ttt cat ctg Cys Gly Phe Thr Val Ser Leu Val Thr Thr Ser Leu Val Phe His Leu 155 160 165	2516
CCC ttc cac tcc tcc aac cag ctc cat cac ttc ttc tgt gac atc tcc Pro Phe His Ser Ser Asn Gln Leu His His Phe Phe Cys Asp Ile Ser 170 175 180	2564
CCt gtc ctt aaa ctg gca tct cag cac tcc ggc ttc agt cag ctg gtc Pro Val Leu Lys Leu Ala Ser Gln His Ser Gly Phe Ser Gln Leu Val 185 190 195	2612
ata ttc atg ctt ggt gta ttt gcc ttg gtc att cct ctg cta ctt atc Ile Phe Met Leu Gly Val Phe Ala Leu Val Ile Pro Leu Leu Leu Ile 200 205 210	2660
cta gtc tcc tac atc cgc atc atc tct gcc att cta aaa atc cct tcc Leu Val Ser Tyr Ile Arg Ile Ile Ser Ala Ile Leu Lys Ile Pro Ser 215 220 225 230	2708
tcc gtt gga aga tac aag acc ttc tcc acc tgt gcc tcc cat ctc att Ser Val Gly Arg Tyr Lys Thr Phe Ser Thr Cys Ala Ser His Leu Ile 235 240 245	2756
Val Val Thr Val His Tyr Ser Cys Ala Ser Phe Ile Tyr Leu Arg Pro 250 255 260	2804
aag act aat tac act tca agc caa gac acc cta ata tct gtg tca tac Lys Thr Asn Tyr Thr Ser Ser Gln Asp Thr Leu Ile Ser Val Ser Tyr	2852

16U 200 PCT FINAL.ST25 265 270 275	
acc atc ctt acc cca ttg ttc aat cca atg att tat agt ctg aga aat Thr Ile Leu Thr Pro Leu Phe Asn Pro Met Ile Tyr Ser Leu Arg Asn 280 285 290	2900
aag gaa ttc aaa tca gcc cta cga aga aca atc ggc caa act ttc tat Lys Glu Phe Lys Ser Ala Leu Arg Arg Thr Ile Gly Gln Thr Phe Tyr 295 300 305 310	2948
cct ctt agt taa agagctattt tttaaactac taatgcctag tacatgccag Pro Leu Ser	3000
gcagaacgtg tgttttatac atttttttc atttaattgt ccagctccac tgtaacataa	3060
gaacatttta catatgagaa gaatgagget cacagaagtt aagacagtet ggetttetae	3120
tetecatgat aetttaacaa gaetaateag atatgggaac agageacaca gttecataac	3180
aaatttaatt atattttact gctttaaata ttgctaattt aaaaactaat atgagagcaa	3240
agatgcatct aaactgatga gagctgtgtc ttgaagtaga gagcttggat acatcaggaa	3300
agaaaagatg tatccaaaaa aaaaaaaaga aagaaaaaag aaaaaaaaaa	3360
aggaaatcca tctatccgta cttttctttt cctaaagaca acagaaaact ttggtcccac	3420
acattetget acaaatettg gtggteettt ttgteeccaa tteattteet taacetacat	3480
attgaaatat cttggccttt acttggggtt gttttgttct tcctttgttt gaggtggaac	3540
cactttatgg ttctcttcct gatgcacatg tatgtccttc acatactagt gtgtcttagc	3600
ccccacattt gttcctgaga caccatacta atttgctctc ttcaaggaag ctactagcat	3660
tgcctacttg ctgaaatatc tcaagtaatt ccaagcaaag ggcttgagtt aatattaata	3720
gaaggctaga ttcctagaat gaccagaaaa ctcatggaaa accctccagt gactcccttt	3780
gccctacaag ataatgccaa gggtccttca ttgtcatgaa tctatcatct agtttccacc	3840
tacctcttca gtattatcat ttctaatttt gttattctcc attttctata tgcctttgt	3900
acactetgaa getaaccaac tatttgettg ttttaaaaca aataaatgtg atgaacaaaa	3960
taaatgtggt ctctgccctc ataggcctta ttgcctggtt caagatagtc ccagtaaaca	4020
gaaaaatgag ggaaaatacc ttaccagttt aagttgattc tctgaagaaa aagtgcatgc	4080
aggcgataga ggagagaata ctaagataaa cctaatttag atcgaatggc atagggttgg	4140
tttcccagag aaactgagag ttaacctgca tgtaacctga agggtaatta aaagtcttca	4200
ggtaaagggg atatccttta ggacagaaga aacaatgtgt acaaaacccc tgaagcaaga	4260
actggatgag ttggagacaa gcaaagaagg cctgtataaa tgctgtttta aaaatgcttt	4320
tcaattgaca aaattatata tatttatggt gtaaaacatg atattttctc ccatcctgta	4380
ggttgcctgt tcactctgat ggtattttct tttgctgtgc agaagctctt tagtttaatt	4440
agatcccatt tgtcaatttt ggcttttgtt gccattgcct ttggtgttta gacatgaagg	4500
ccttgcccat gcctatgccc tgaatggtac tgcctaggtt ttcttctagg gtttttatgg	4560
ttttaggtct aacatgtaag tcttttatcc atctggaata aatttttgta taaggtgtaa	4620
ggaagggatc cagtttcagc tttctacata tggctagcca gttttcccag caccatttat	4680
taaataggga atcctttccc catttcttgt ttttgtcaga caaagggcta atatccagaa	4740
tctacaatga actcaaacaa atttacaaga aaaaaacaaa caaccccatc aaaaa gtggg	4800
caaaggatat gaacagacac ttctcaaaag aagacattta tgcagccaga aaacacatga	4860
aaaaatgctc atcactggcc atcagagaaa tgcaaatcaa aaccacaatg agataccatc	4920
tcacaccagt tagaatggcg atcattaaaa agtcaggaaa caacaggtgc gggagaagat	4980
-	

4982

at <210> <211> 313 <212> PRT <213> Homo sapiens <400> 13 Met Glu Gln Val Asn Lys Thr Val Val Arg Glu Phe Val Val Leu Gly Phe Ser Ser Leu Ala Arg Leu Gln Gln Leu Leu Phe Val Ile Phe Leu 20 25 30Leu Leu Tyr Leu Phe Thr Leu Gly Thr Asn Ala Ile Ile Ser Thr 35 40 45Ile Val Leu Asp Arg Ala Leu His Thr Pro Met Tyr Phe Phe Leu Ala 50 60Ile Leu Ser Cys Ser Glu Ile Cys Tyr Thr Phe Val Ile Val Pro Lys 65 70 75 80Met Leu Val Asp Leu Leu Ser Gln Lys Lys Thr Ile Ser Phe Leu Gly 85 90 95 Cys Ala Ile Gln Met Phe Ser Phe Leu Phe Phe Gly Ser Ser His Ser Phe Leu Leu Ala Ala Met Gly Tyr Asp Arg Tyr Met Ala Ile Cys Asn 115 120 125Pro Leu Arg Tyr Ser Val Leu Met Gly His Gly Val Cys Met Gly Leu 130 135 140 Met Ala Ala Cys Ala Cys Gly Phe Thr Val Ser Leu Val Thr Thr 145 150 155 160 Ser Leu Val Phe His Leu Pro Phe His Ser Ser Asn Gln Leu His His 165 170 175 Phe Phe Cys Asp Ile Ser Pro Val Leu Lys Leu Ala Ser Gln His Ser 180 185 190 Gly Phe Ser Gln Leu Val Ile Phe Met Leu Gly Val Phe Ala Leu Val Ile Pro Leu Leu Leu Ile Leu Val Ser Tyr Ile Arg Ile Ile Ser Ala 210 215 220 Ile Leu Lys Ile Pro Ser Ser Val Gly Arg Tyr Lys Thr Phe Ser Thr 225 230 235 240Cys Ala Ser His Leu Ile Val Val Thr Val His Tyr Ser Cys Ala Ser 245 250 255 ...

Phe Ile Tyr Leu Arg Pro Lys Thr Asn Tyr Thr Ser Ser Gln Asp Thr

```
16U 200 PCT FINAL.ST25
 Leu Ile Ser Val Ser Tyr Thr Ile Leu Thr Pro Leu Phe Asn Pro Met
 Ile Tyr Ser Leu Arg Asn Lys Glu Phe Lys Ser Ala Leu Arg Arg Thr
 Ile Gly Gln Thr Phe Tyr Pro Leu Ser
                      310
<210>
       14
<211>
        24
 <212>
        DNA
<213> Homo sapiens
<400> 14
cctgttcact ctgggcacca atgc
                                                                            24
<210> 15
<211>
<212> DNA
<213> Homo sapiens
<400> 15
ctggttggag gagtggaagg gcag
                                                                            24
<210> 16
<211>
        50
<212>
        DNA
<213> Homo sapiens
<400> 16
tacctttctg tatataaaaa catataacta atacacacac actcatacac
                                                                            50
<210>
<211>
        17
        50
<212> DNA
<213> Homo sapiens
<400> 17
cttcagaagt atataaatga agactggata ccagcaagac atactggatg
                                                                            50
<210> 18
<211> 50
<212> DNA
<213> Homo sapiens
<400> 18
cccttggaga tataaaaagt tcccagtaaa tagatgtgtg ctcacatctt
                                                                            50
<210> 19
<211>
       50
<212>
       DNA
<213>
      Homo sapiens
<400> 19
taatactatg taaaaatcca ctggactaga atcagctgtc ctcatgtgcc
                                                                            50
<210> 20
<211> 960
<212> DNA
<213> Homo sapiens
<220>
<221> CDS
<222>
       (1)..(960)
<223>
<400> 20
atg aca cag ttg acg gcc agt ggg aat cag aca atg gtg act gag ttc
Met Thr Gln Leu Thr Ala Ser Gly Asn Gln Thr Met Val Thr Glu Phe
                                                                           48
```

1				5					10	16U	200	PCT	FINA	L.ST	25	
ctc Leu	ttc Phe	tct Ser	atg Met 20	ttc Phe	ccg Pro	cat His	gcg Ala	cac His 25	aga Arg	ggt Gly	ggc Gly	ctc Leu	tta Leu 30	ttc Phe	ttt Phe	96
att Ile	ccc Pro	ttg Leu 35	ctt Leu	ctc Leu	atc Ile	tac Tyr	gga Gly 40	ttt Phe	atc Ile	cta Leu	act Thr	gga Gly 45	aac Asn	cta Leu	ata Ile	144
atg Met	ttc Phe 50	att Ile	gtc Val	atc Ile	cag Gln	gtg Val 55	ggc Gly	atg Met	gcc Ala	ctg Leu	cac His 60	acc Thr	cct Pro	ttg Leu	tat Tyr	192
								ctg Leu								240
acc Thr	atc Ile	ccc Pro	aag Lys	atg Met 85	ctg Leu	tcc Ser	tgc Cys	cta Leu	atc Ile 90	agt Ser	gag Glu	cag Gln	aag Lys	agc Ser 95	att Ile	288
								atg Met 105								336
atc Ile	aca Thr	gaa Glu 115	agc Ser	tgt Cys	gtc Val	ctg Leu	aca Thr 120	gca Ala	atg Me t	gcc Ala	att Ile	gac Asp 125	agg Arg	tac Tyr	ata Ile	384
								cca Pro								432
tgt Cys 145	atc Ile	cag Gln	ctg Leu	aca Thr	gtt Val 150	gga Gly	tcc Ser	tgc Cys	ttt Phe	tgt Cys 155	ggc Gly	ttc Phe	ctc Leu	ctt Leu	gtg Val 160	480
ctt Leu	cct Pro	gag Glu	att Ile	gca Ala 165	tgg Trp	att Ile	tcc Ser	acc Thr	ttg Leu 170	cct Pro	ttc Phe	tgt Cys	ggc Gly	ser 175	aac Asn	528
								ttc Phe 185								576
tgc Cys	aca Thr	gat Asp 195	aca Thr	ttc Phe	cta Leu	gtg Val	gtc Val 200	att Ile	gtg Val	gat Asp	gcc Ala	atc Ile 205	cat His	gca Ala	gcg Ala	624
								att Ile								672
								tca Ser								720
ttt Phe	tcc Ser	acc Thr	tgt Cys	gct Ala 245	gct Ala	cac His	ctt Leu	gct Ala	gtg Val 250	ttc Phe	ttg Leu	cta Leu	ttt Phe	ttt Phe 255	Gly Gly	768
								ttc Phe 265								816
tgg Trp	gac Asp	aca Thr 275	gca Ala	att Ile	gct Ala	gtc Val	act Thr 280	ttt Phe	gtt Val	atc Ile	ctt Leu	gct Ala 285	ccc	ttt Phe	ttc Phe	864
								aac Asn								912
								agg Arg							tag	960

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

<210> 21

<211> 319

<212> PRT

<213> Homo sapiens

<400> 21

Leu Phe Ser Met Phe Pro His Ala His Arg Gly Gly Leu Leu Phe Phe 20

Ile Pro Leu Leu Ile Tyr Gly Phe Ile Leu Thr Gly Asn Leu Ile 35 40

Met Phe Ile Val Ile Gln Val Gly Met Ala Leu His Thr Pro Leu Tyr 50 55 60

Phe Phe Ile Ser Val Leu Ser Phe Leu Glu Ile Cys Tyr Thr Thr Thr 65 70 80

Thr Ile Pro Lys Met Leu Ser Cys Leu Ile Ser Glu Gln Lys Ser Ile $85 \hspace{1cm} 90 \hspace{1cm} 95$

Ser Val Ala Gly Cys Leu Leu Gln Met Tyr Phe Phe His Ser Leu Gly 100 105 110

Ile Thr Glu Ser Cys Val Leu Thr Ala Met Ala Ile Asp Arg Tyr Ile 115 120 125

Ala Ile Cys Asn Pro Leu Arg Tyr Pro Thr Ile Met Ile Pro Lys Leu 130 $$ 135 $$ 140

Cys Ile Gln Leu Thr Val Gly Ser Cys Phe Cys Gly Phe Leu Leu Val 145 $$ 150 $$ 155 $$ 160

Leu Pro Glu Ile Ala Trp Ile Ser Thr Leu Pro Phe Cys Gly Ser Asn 165 170 175

Gln Ile His Gln Ile Phe Cys Asp Phe Thr Pro Val Leu Ser Leu Ala $_{\odot}$ 180 $^{\circ}$ 185 $^{\circ}$ 190 $^{\circ}$

Cys Thr Asp Thr Phe Leu Val Val Ile Val Asp Ala Ile His Ala Ala 195 200 205

Glu Ile Val Ala Ser Phe Leu Val Ile Ala Leu Ser Tyr Ile Arg Ile 210 215 220

Ile Ile Val Ile Leu Gly Met His Ser Ala Glu Gly His His Lys Ala 225 230 235 240

Phe Ser Thr Cys Ala Ala His Leu Ala Val Phe Leu Leu Phe Phe Gly 245 250 255

Ser Val Ala Val Met Tyr Leu Arg Phe Ser Ala Thr Tyr Ser Val Phe 260 265 270

Trp Asp Thr Ala lle Ala Val Thr Phe Val Ile Leu Ala Pro Phe Phe 275 280 285

290 295 300	
Gly Arg Leu Phe His Tyr Gln Lys Arg Ala Gly Trp Ala Gly Lys 305 310 315	
<210> 22 <211> 24 <212> DNA <213> Homo sapiens	
<400> 22 ctctatgttc ccgcatgcgc acag	24
<210> 23 <211> 27 <212> DNA <213> Homo sapiens	
<400> 23 gcaaggtgga aatccatgca atctcag	27
<210> 24 <211> 50 <212> DNA <213> Homo sapiens	
<400> 24 agacagacgt taaaaaatga ccaaacctac agaaaatatt tccagataat	50
<210> 25 <211> 900 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (1)(900) <223>	
<400> 25 atg atc acc gag ttc atc ctt ata ggc ttc tca aac ctg ggg gat ctg	
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1 5 10 15	48
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu	48 96
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1 10 15 cag atc ctt ctc ttc ttt atc ttc cta tta gtc tac ctg acc act ctg Gln Ile Leu Leu Phe Phe Ile Phe Leu Leu Val Tyr Leu Thr Thr Leu	
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1	96
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1	96 144
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1	96 144 192
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1	96 144 192 240
Met Ile Thr Glu Phe Ile Leu Ile Gly Phe Ser Asn Leu Gly Asp Leu 1	96 144 192 240 288

Leu		al 30	Leu	Ala	Ser	Ser	Phe 135	Cys	Gly	Phe	16U Leu	200 Thr 140	PCT Ser	FINA Val	L.ST Ile	25 Val	
aat Asr 145	1	tc le	ctg Leu	gtg Val	ttc Phe	agt Ser 150	gtg Val	ctc Leu	ctc Leu	tgt Cys	gcc Ala 155	tcc Ser	aat Asn	cgg Arg	atc Ile	aac Asn 160	480
cac His	t s P	tt	ttc Phe	tgt Cys	gac Asp 165	att Ile	tcc Ser	cct Pro	gtc Val	ata Ile 170	aaa Lys	ctg Leu	ggc Gly	tgc Cys	aca Thr 175	gac Asp	528
Thi	c A	sn	Leu	Lys 180	Glu	Met	Val	Ile	ttt Phe 185	Pne	гел	ser	116	190	vai	Бей	576
Lei	ı V	al	Pro 195	Leu	Val	Leu	lle	Phe 200	atc Ile	Ser	туг	116	205	116	val	361	624
Th	r I	1e 210	Leu	Lys	Ile	Ser	Ser 215	Val	gaa Glu	GIĀ	GIN	220	Lys	AIG	FILE	A10	672
Th 22	r (:ys	Ala	Ser	His	Leu 230	Thr	Val	gtc Val	Val	235	HIS	i Tyr	GIY	Cys	240	720
Se	r I	Phe	lle	Tyr	Leu 245	Arg	Pro	Thr	tcc Ser	Leu 250	Tyr	Ser	: Ser	ASP	255	нър	768
Ar	g 1	Leu	Val	Ala 260	Val	Thr	Tyr	Thr	gtg Val 265	116	Thr	Pro	o rec	270	MSII	rio	816
ct Le	t e	gtc Val	tat Tyr 275	Thr	ctg Leu	aga Arg	aat Asn	aaa Lys 280	Glu	gta Val	aag Lys	ato Met	g gct t Ala 285	ren	aga Arg	aag Lys	864
gt Va	1	ctg Leu 290	Gly	aga Arg	tgc Cys	tta Leu	aat Asn 295	Ser	aaa Lys	act Thr	gta Val	i tga	a				900
<2 <2	210 211 212 213	>	26 299 PRT Homo	sar	oiens	.											
< 4	00	>	26														
M∈ 1	2t	Ile	Thi	Glu	Phe 5	e Ile	e Lev	1 Ile	e Gly	7 Phe 10	e Se	r As	n Le	u Gly	/ Asp 15	o Leu	
G:	ln	Ile	Lei	20	u Phe	e Phe	e Ile	e Ph∘	e Lei 25	ı Lei	ı Va	1 T y	r Le	u Th: 30	r Th	r Leu	
Mo	et	Ala	35	n Th	r Th	r Ile	e Me	t Th 40	r Va	1 110	≥ Hi	s Le	u As 45	p Ar	g Al	a Leu	
н	is	Th: 50	Pr	o Me	t Ty	r Phe	9 Pho 55	e Le	u Ph	e Va	l Le	u Se 60	er Cy	s Se	r Gl	u Thr	
C 6		Туі	Th	r Le	u Va	1 Ile 70	e Va) Pr	o Ly	s Me	t Le 75	u Th	r As	n Le		u Ser 80	; ;
A	la	116	e Pr	o Th	r Il 85		r Ph	e Se	r Gl	у Су 90	s Va	1 Vā	1 G1	n Le	u Ту 95	r Lev	1

Tyr	Asp	Arg 115	Tyr	Val	Ala	Ile	Cys 120	Asn	Pro	16U Leu	200 Asn	PCT Tyr 125	FINA Thr	AL.ST Leu	r25 Ile	
Leu	Val 130	Leu	Ala	Ser	Ser	Phe 135	Cys	Gly	Phe	Leu	Thr 140	Ser	Val	Ile	Val	
Asn 145	Ile	Leu	Val	Phe	Ser 150	Val	Leu	Leu	Cys	Ala 155	Ser	Asn	Arg	Ile	Asn 160	
His	Phe	Phe	Cys	Asp 165	Ile	Ser	Pro	Val	Ile 170	L ys	Leu	G1 y	Cys	Thr 175	Asp	
Thr	Asn	Leu	Lys 180	Glu	Met	Val	Ile	Phe 185	Phe	Leu	Ser	lle	Leu 190	Val	Leu	
Leu	Val	Pro 195	Leu	Val	Leu	Ile	Phe 200	Ile	Ser	Туr	Ile	Phe 205	lle	Val	Ser	
Thr	11e 210	Leu	Lys	Ile	Ser	Ser 215	Val	Glu	Gly	Gln	Cys 220	Lys	Ala	Phe	Ala	
Thr 225	Cys	Ala	Ser	His	Leu 230	Thr	Val	Val	Val	Val 235	His	туг	Gly	Cys	Ala 240	
Ser	Phe	Ile	Туr	Leu 245	Arg	Pro	Thr	Ser	Leu 250	Tyr	Ser	Ser	Asp	Lys 255	Asp	
Arg	Leu	Val	Ala 260	Val	Thr	Tyr	Thr	Val 265	Ile	Thr	Pro	Leu	Leu 270	Asn	Pro	
Leu	Val	Tyr 275	Thr	Leu	Arg	Asn	Lys 280	Glu	Val	Lys	Met	Ala 285	Leu	Arg	Lys	
Val	Leu 290	Gly	Arg	Cys	Leu	Asn 295	Ser	Lys	Thr	Val						
<210		27														
<211 <212 <213	'> C	9 NA Iomo	sapi	ens												
<400 tate		:7 :tc c	taat	gtto	a at	atac	·t cc									29
			- 5 5 -	,	- 9-	9-9-										23
<210	> 3	8														
<212 <213		NA Omo	sapi	ens												
<400 catc		8 cag	aacc	tttc	t ca	gage	cato	:								30
<210		•														
<211 <211 <212	> 5	9 0 NA														
<213	_	omo	sapi	ens												
<400 gtca		9 tg t	ataa	gcac	g ca	gtgc	aaag	gaa	atat	taa	aact	agaa	cc	, :		50
<210	> 3	0														
<211 <212	> 5															
<213	> H	omo	sapi	ens												

<400> 30 tttcttcatt tataacatga gggggcttgg ctagatattt aacagcctgc	50
<210> 31 <211> 50 <212> DNA <213> Homo sapiens	
<400> 31 gctagatatt taacagcctg cctgtattga ccacttatgc atcaggaaat	50
<210> 32 <211> 50 <212> DNA <213> Homo sapiens	
<400> 32 atttgagtta tgtatatgag agactgggta catcactttt tacttgtttt	50
<210> 33 <211> 5086 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (2034)(2972) <223>	
<400> 33 tatccaaatg gtgaaagaga ttctagaaca aggaaagagc tacagcaaag gttttaaatg	60
atatgtcact gaacacattt gatcgatgga aacgcagaac ctaatttaga atttaacagg	120
atcactctgg tgtgttgaga tgaggctaca agtgaacaaa tgcaagtagg gagatctgtt	180
aggagtcaat tacagtaaga ggggagagat aaaagtgact tggaccgagg tggtcaaaca	240
tagtcagttc ctggatatat gagagaaaga tagaaacaag gatgactgca ggagtttagc	300
ttgtcagttg aaagattgca attgccatca tttgtgatgg ggaagactag gggtagagac	360
cccaggagtt cagtttgaga tygctcttcg actcccaaga ggagatgtga gtaggcagtg	420
aaatatatga gtctggagta gcagagaaaa atatcgcctg agatatggat ttagatgtct	480
tcaacacatt tatagtgttt aaagctctgg tattggatgg tatagagcag aggagttgag	540
tttatataga agagaaaaaa aaaagattaa acactgacca tgggcactgt gacattaaaa	600
ggatggggca tggaggagaa actaaagttg gagaatgaga aggaatgact aataagatag	660
aaagtaacca aaagtatagt accccgagaa tcaagtcaag	720
ataaatcaat actgtcaaga aacagatagt ccaagtaagc tgaggaatga gaaatgacca	780
ttggatccag gaaatcttag ataattaatg tctatgagaa aggaggtttt aatggagtgg	840
tggtagtata aatctaatta gagtgggttt aagaagaaac ttaaagagag gcattaaagg	900
caatgcgtat agccgactct tggaagagtt ttcttttagg gacatagaaa gaaatagagc	960
agtggctgtg ggatgagtaa agagaaagaa tttaaggctc ttgcttttt gtttgtttag	1020
tagatgagaa taatagcatg tttttacatt gatagagtat tccatgaaag agctgtataa	1080
tagttagttg tttctctata ctctgtatta caatattagt ttgttaacat caggtgccac	1140
attttatttg tttagtccct gttctaagta taatgcccag agtactgaaa ataatcaatt	1200
attgttacat tgacctcaac acagtagagc atgtatattt aatatctaca gaagcaataa	1260
accagaaaag agcatttgaa gttgatagag ggggaaatgg caggaagaac tgatgaagtg	1320
gccacagtct gaagttgaaa tgcagaaaga tagatttgcc tcctgtcttt ctttggcttt	1380

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

tttatttact ctaac	ccttct tatttt	ttgac tggagctctc	accagtgtcc aaaagaggtc	1440
taaattctga cctad	catgcc cctgaa	aagat gctagcagac	ctgagttctc ataaaggaat	1500
aggagggagc agaag	gggaaa acaatt	gatt ctttggtagc	cagaaagttg aagaagaaaa	1560
caaattaaaa tgaga	aaatta gaaaat	taata ttcaaattat	atatatttgg tccagtacgg	1620
tatcaatata ttato	cagtat aaatga	atgat ttttacctta	gatgaacaat atgtataaat	1680
gttaatatat acct	tggatt agaaat	tacct aaatttctaa	aatctatata gattctattg	1740
agaaagtcaa ctgg	gttaca ggatgg	gatta ggaaggccaa	aaatgagctg tgttaatcag	1800
ggaagactaa acata	aaaggt gaatag	stotg aaggaggotg	ttgacaggaa gggcagggag	1860
ggatggaatt gaaat	tgttga cctctc	caaag catttactta	gagggcttta ctctggaggt	1920
gagagaaggg agggd	caatag taattt	gagg gttgccttct	tgttagaacc ctatagttca	1980
actttettte etate	cttcc acactt	caca totagggaca	tgaatggtga gca atg Met 1	2036
			atc atc ttg ggc ttc Ile Ile Leu Gly Phe 15	2084
			ctc ttg ttg ctt ctc Leu Leu Leu Leu 30	2132
			ata ttc ctg gtg gtc Ile Phe Leu Val Val 45	2180
			cac ttt gtc agc att His Phe Val Ser Ile 65	2228
			acc atc cct aag atg Thr Ile Pro Lys Met 80	2276
			tca ttc tct ggg tgt Ser Phe Ser Gly Cys 95	2324
			gcg act gag tgc tat Ala Thr Glu Cys Tyr 110	2372
			gcc atc tgc cgg ccc Ala Ile Cys Arg Pro 125	2420
			tgt gca gag att gcc Cys Ala Glu Ile Ala 145	2468
			gta gtt gaa att tcc Val Val Glu Ile Ser 160	2516
			cgc att cag cac gtc Arg Ile Gln His Val 175	2564
	Pro Pro Val		tgc act gat acg tct Cys Thr Asp Thr Ser 190	2612
			tgc aag atc cta gcc Cys Lys Ile Leu Ala 205	2660
			atc atc tgc aca gtg Ile Ile Cys Thr Val	2708

16U 200 PCT FINAL.ST25 210 215 220 225	
ctc aga att ccc tca gct gcc ggc aag agg aag gcc atc tcc acg tgt Leu Arg Ile Pro Ser Ala Ala Gly Lys Arg Lys Ala Ile Ser Thr Cys 230 235 240	2756
gcc tcc cac ttc act gtg gtt ctc atc ttc tat ggg agc atc ctt tcc Ala Ser His Phe Thr Val Val Leu Ile Phe Tyr Gly Ser Ile Leu Ser 245 250 255	2804
atg tat gtg cag ctg aag aag agc tac tca ctg gac tat gac cag gcc Met Tyr Val Gln Leu Lys Lys Ser Tyr Ser Leu Asp Tyr Asp Gln Ala 260 265 270	2852
ctg gca gtg gtc tac tca gtg ctc aca ccc ttc ctc aac ccc ttc atc Leu Ala Val Val Tyr Ser Val Leu Thr Pro Phe Leu Asn Pro Phe Ile 275 280 285	2900
tac age ttg ege aac aag gag ate aag gag get gtg agg agg eag eta Tyr Ser Leu Arg Asn Lys Glu Ile Lys Glu Ala Val Arg Arg Gln Leu 290 295 300 305	2948
aag aga att ggg ata ttg gca tga gttggggctg agagtaggcc aaggccgggc Lys Arg Ile Gly Ile Leu Ala 310	3002
ctgaggatat ggtggcccca gggatcaaca gtggccagag acgagaaact aaaaattcag	3062
tgcttttcta tgtggggtgg tggagctgca gcaagtgctg actgacttcc agtgttatag	3122
cgaccttcat actgtctgct ggagccacat ttggcttgag accagagact agggaaagta	3182
cacatecett caacatgatg tagtgeagtg attiteaaaa eteagatgit tatgiateae	3242
acttaggttt tttttaaaat ctgtgtctta cctattatac gtttataggc atttttcaaa	3302
tttacttgac ttaatataaa tatagtcagg catgtcctaa acaaaatgtg attcatgatg	3362
ttttttgtac cacttgcaat catttcatgt ggagaagact ggtacagtag aaaaaagcat	3422
gttttttgaa ctcatatata tctggattta aatcatgttt tattcagtca cttgctaatt	3482
acttaatett tagaaagtaa ettageatet etgagtetta attteattat ttgataatgg	3542
tattttcttg aagagtgttt tgaatattaa cgttaagatt tgtaaaccac agtgcacagt	3602
gtctgacatg taggtgatag taaataaata aggacttgtt tttatttatt ttattctgcg	3662
aagacttcac atcattactc tgggtcttag aacaatatct agtaaaacat aaataaacaa	3722
aaatactttc caagtatttt ctccaaagga aaggagcaaa ccagccagaa ggaatacttg	3782
tatagtatac aagtatacta tacttgaaaa gtatagtttg tcacagttct gttctgacaa	3842
gtttcatgta cctgtcttag tggtcctaat atctatggcc agtataatgt atgaaagtat	3902
aggagttgag tcagtggaaa gaaataggat tactttttac atcgaaccat ttctttattg	3962
aattgtaagc taattatttc ctgaaacgtg tgaaaaataa ttctaaaatg tagcatatga	4022
gagatctggg gattcaatta atagctaata ttatgtattc tttatgtatc cttccatgaa	4082
tggaggatca aatattaact acaagaaatc tttgaattct atagaacttc ctaagaagat	4142
tacaaaatat ttttaatacc acacttttaa aggtattcat ccatccatgc attcaaatta	4202
acacgtttat ttagctctta ctatatatca gatgcagtgt caactctaca aaagcaatga	4262
acaagacata tatatgtcca ggtcctacct ttagggtgtt ttaaaagagt tgagaatata	4322
aatattaaaa ttataattaa tttataatta gttataatta attataattg tgggaagtag	4'382
tattaagata aacatgcatt ctccttttt ttcacttgtc tttgaagttt attgagaatt	4442
ttaagcagat aaatgttttt acattaaata atcaccagga attcaaaata ttatactcta	4502
tcaaatggga acttgaattg ttctatttat atatgtagca ttctatttat aaatatattt	4562
cattlagtgt ttcatctaga ataaaaatga caagaaataa aattattaaa aacaagttgt	4622

gtttgacttt	tggtaaaatt	ttttgtcctg	gacatttttg	atgactaagt	atcactaaat	4682
ctatgctagg	taaatttgcc	cctattattt	tcttttttat	tttattttat	tttatttcat	4742
tattattta	tttagggtac	atgtgcacaa	cgtgcaagtt	ttttacatat	gtatacatgt	4802
gccatgttgg	tgtgctgcac	ccattaactc	atcatttagc	attaggagta	tctcctaatg	4862
ctatccctcc	cccatccccc	aaccccacaa	cagtccccag	tgtgtgatgt	tccccttctc	4922
aatatcatac	tgaatgggca	aaaactggaa	gcattccctt	tgaaaacggg	cacaagacag	4982
ggatgccctc	tctcaccact	cctattcaac	atagtgtttg	atgttctggc	cagggcaatc	5042
aggtaggaga	aggaaattaa	gggtgttcaa	ttaggaaaag	agga		5086

<210> 34

<211> 312

<212> PRT

<213> Homo sapiens

<400> 34

Met Asp Thr Gly Asn Trp Ser Gln Val Ala Glu Phe Ile Ile Leu Gly 1 $$ 5 $$ 10 $$ 15

Phe Pro His Leu Gln Gly Val Gln Ile Tyr Leu Phe Leu Leu Leu 20 \$25\$

Leu Ile Tyr Leu Met Thr Val Leu Gly Asn Leu Leu Ile Phe Leu Val 35 40 45

Val Cys Leu Asp Ser Arg Leu His Thr Pro Met Tyr His Phe Val Ser 50 55 60

Ile Leu Ser Phe Ser Glu Leu Gly Tyr Thr Ala Ala Thr Ile Pro Lys 65 70 75 80

Met Leu Ala Asn Leu Leu Ser Glu Lys Lys Thr Ile Ser Phe Ser Gly $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$

Cys Leu Leu Gl
n Ile Tyr Phe Phe His Ser Leu Gly Ala Thr Glu Cys 100
 $105\,$ 110

Tyr Leu Leu Thr Ala Met Ala Tyr Asp Arg Tyr Leu Ala Ile Cys Arg 115 120 125

Pro Leu His Tyr Pro Thr Leu Met Thr Pro Thr Leu Cys Ala Glu Ile 130 135 140

Ala Ile Gly Cys Trp Leu Gly Gly Leu Ala Gly Pro Val Val Glu Ile 145 150 150 160

Ser Leu Ile Ser Arg Leu Pro Phe Cys Gly Pro Asn Arg Ile Gln His 165 170 175

Val Phe Cys Asp Phe Pro Pro Val Leu Ser Leu Ala Cys Thr Asp Thr 180 185 190

Ser Ile Asn Val Leu Val Asp Phe Val Ile Asn Ser Cys Lys Ile Leu 195 200 205

Ala Thr Phe Leu Leu Ile Leu Cys Ser Tyr Val Gln Ile Ile Cys Thr 210 215 220

225	u Arg	116	710	230	Ald	AIG	GIY	цуз	235	Lys	AIG	116	361	240	
Cys Ala	a Ser	His	Phe 245	Thr	Val	Val	Leu	11e 250	Phe	Tyr	Gly	Ser	11e 255	Leu	
Ser Me	t Tyr	Val 260	Gln	Leu	Lys	Lys	Ser 265	Туr	Ser	Leu	Asp	Туг 270	Asp	Gln	
Ala Le	u Ala 275	Val	Val	Туr	Ser	Val 280	Leu	Thr	Pro	Phe	Leu 285	Asn	Pro	Phe	
11e Ty: 29		Leu	Arg	Asn	Lys 295	Glu	Ile	Lys	Glu	Ala 300	Val	Arg	Arg	Gln	
Leu Ly: 305	s Arg	Ile	Gly	11e 310	Leu	Ala									
<210><211><211><212><213>	35 29 DNA Homo	sapi	iens												
<400> ggaact	35 ggag o	ccago	gtago	ca ga	atto	catc									29
<210> <211> <212> <213>	36 25 DNA Homo	sapi	iens												
<400> ggagca	36 gagg a	atcaç	gcago	ga aç	ggtg										25
<210><211><211><212><213>	37 50 DNA Homo	sapi	i ens												
<400> acactg	37 cagt t	atat	tagge	gt g	gecea	aggta	agt!	tgago	etgg	tgaa	atti	ga			50
<210><211><211><212><213>	38 50 DNA Homo	sapi	iens												
<400> gcactg		attaa	aaag	ga tọ	3ggg	atg	g ago	gagaa	aact	aaaq	gttg	gag			50
<210><211><211><212><213>	39 50 DNA Homo	sapi	iens												
<400> attcaa	39 atta t	atat	tatti	tg gt	ccaç	tac	g gta	atca	atat	atta	itca	gta			50
<210><211><211><212><213>	40 848 DNA Homo	sapi	iens												
<220> <221> <222>	CDS (6)	. (848	3)												

Page 20

<	22	3>

<40 gaa	tc a	et A									ln S				tt gct eu Ala 15	50
				act Thr 20												98
				ctc Leu												146
				atg Met												194
				gtt Val												242
				tcc Ser												290
				agc Ser 100												338
				aga Arg												386
				tca Ser												434
				ttg Leu												482
				ttg Leu												530
His	Gly	Thr	Val	Ile 180	Gly	Val	Ala	Thr	Gly 185	Cys	Asn	Phe	Val	Leu 190	Ile	578
Leu	·G1 y	Leu	Туг 195	gga Gly	Gly	Ile	Leu	Asn 200	Ala	Val	Leu	Lys	Leu 205	Pro	Ser	626
Ala	Ala	Ser 210	Ser	gcc Ala	Lys	Ala	Phe 215	Ser	Thr	Cys	Ser	Ser 220	His	Йаl	Thr	674
Val	Val 225	Ala	Leu	ttc Phe	Tyr	Ala 230	Ser	Ala	Phe	Thr	Val 235	Tyr	Val	Gly	Ser	722
Pro 240	Gly	Ser	Arg	ect Pro	Glu 245	Ser	Thr	Asp	Lys	Leu 250	Val	Ala	Leu	Val	Tyr 255	770
Ala	Leu	Ile	Thr	Pro 260	Phe	Leu	Asn	Pro	11e 265						Asn	818
				tat Tyr					tga							848

<210> 41 <211> 280 <212> PRT

<213> Homo sapiens

<400> 41

Met Asp His Val Ser His Asn Trp Thr Gln Ser Phe Ile Leu Ala Gly 1 5 10 15

Phe Thr Thr Gly Thr Leu Gln Pro Leu Ala Phe Leu Gly Thr Leu $20 \hspace{1cm} 25 \hspace{1cm} 30 \hspace{1cm}$

Cys Ile Tyr Leu Leu Thr Leu Ala Gly Asn Ile Leu Ile Ile Val Leu $35 \hspace{1cm} 40 \hspace{1cm} 45$

Arg Cys Gly Met Ser Ala Pro Gln Cys Pro Cys Cys Cys Thr Pro Cys 50 55 60

Ser Lys Gly Val His Pro Ser His Gln Leu Tyr Ala Leu Phe Ser Tyr 65 70 75 80

Val Phe His Ser Leu Gly Met Thr Glu Cys Tyr Leu Leu Gly Val Met 85 90 95

Ala Leu Asp Ser Tyr Leu Ile Ile Cys His Pro Leu His Tyr His Ala $100 \hspace{1.5cm} 105 \hspace{1.5cm} 110$

Leu Met Ser Arg Gln Val Gln Leu Arg Leu Ala Gly Ala Ser Trp Val 115 120 125

Ala Gly Phe Ser Ala Ala Leu Val Pro Ala Thr Leu Thr Ala Thr Leu 130 135 140

Pro Phe Cys Leu Lys Glu Val Ala His Tyr Phe Cys Asp Leu Ala Pro 145 150 155 160

Leu Met Arg Leu Ala Cys Val Asp Thr Ser Trp His Ala Arg Ala His 165 170 175

Gly Thr Val Ile Gly Val Ala Thr Gly Cys Asn Phe Val Leu Ile Leu 180 185 190

Gly Leu Tyr Gly Gly Ile Leu Asn Ala Val Leu Lys Leu Pro Ser Ala 195 200 205

Ala Ser Ser Ala Lys Ala Phe Ser Thr Cys Ser Ser His Val Thr Val 210 215 220

Val Ala Leu Phe Tyr Ala Ser Ala Phe Thr Val Tyr Val Gly Ser Pro 225 230 230 235 240

Gly Ser Arg Pro Glu Ser Thr Asp Lys Leu Val Ala Leu Val Tyr Ala 245 250 255

Glu Leu Leu Tyr Cys Phe Leu Cys 275 280

<210> 42

<211> 26

<212> DNA

<213> Homo sapiens	
<400> 42 teaceaceac tgggacecta caacet	26
<210> 43 <211> 23 <212> DNA <213> Homo sapiens	
<400> 43 ggccacacca atcactgtgc cat	23
<210> 44 <211> 50 <212> DNA <213> Homo sapiens	
<400> 44 caatctgtta tttatacggc ctctacatcc atccagtacc tgcttatgta	50
<210> 45 <211> 50 <212> DNA <213> Homo sapiens	
<400> 45 gttctctttt tataaaaggc tatgtgggac ttgcaaaact tctagtggcc	50
<210> 46 <211> 50 <212> DNA <213> Homo sapiens	
<400> 46 gaacatgaaa tataagtagg ggagtatett ggggtagaaa ggatgeegag	50
<210> 47 <211> 1476 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (1)(1476) <223>	
<400> 47 atg gtc acc gaa ttc ctg ttg ctg ggt ttt tcc agc ctt ggt gaa att Met Val Thr Glu Phe Leu Leu Gly Phe Ser Ser Leu Gly Glu Ile 1 5 10 15	48
cag ctg gcc ctc ttt gta gtt ttt ctt ttt ctg tat cta gtc att ctt Gln Leu Ala Leu Phe Val Val Phe Leu Phe Leu Tyr Leu Val Ile Leu 20 25 30	96
agt ggc aat gtc acc att atc agt gtc atc cac ctg gat aaa agc ctc Ser Gly Asn Val Thr Ile Ile Ser Val Ile His Leu Asp Lys Ser Leu 35 40 45	144
35 40 45	
cac aca cca atg tac ttc ttc ctt ggc att ctc tca aca tct gag acc His Thr Pro Met Tyr Phe Phe Leu Gly Ile Leu Ser Thr Ser Glu Thr 50 55 60	192
cac aca cca atg tac ttc ttc ctt ggc att ctc tca aca tct gag acc His Thr Pro Met Tyr Phe Phe Leu Gly Ile Leu Ser Thr Ser Glu Thr	240
cac aca cca atg tac ttc ttc ctt ggc att ctc tca aca tct gag acc His Thr Pro Met Tyr Phe Phe Leu Gly Ile Leu Ser Thr Ser Glu Thr 50 60 ttc tac acc ttt gtc att cta ccc aag atg ctc atc aat cta ctt tct Phe Tyr Thr Phe Val Ile Leu Pro Lys Met Leu Ile Asn Leu Leu Ser	

			100					105		1 6 U	200	PCT	FINA 110	AL.ST	25	
tat Tyr	gat Asp	cgc Arg 115	tat Tyr	gct Ala	gcc Ala	att Ile	tgt Cys 120	cac His	cct Pro	ctg Leu	cat His	tac Tyr 125	ccc Pro	act Thr	ctt Leu	384
										gct Ala						432
										tta Leu 155						480
										ttc Phe						528
										gtt Val						576
										ccc Pro						624
										ctg Leu						672
										gcc Ala 235						720
										atc Ile						768
										gtg Val						816
										tat Tyr						864
										ggc Gly						912
										att Ile 315						960
										tcc Ser						1008
										gtc Val						1056
										gcc Ala						1104
										tgt Cys						1152
										ctg Leu 395						1200
										aag Lys						1248
tgt	gat	att	tca	cca	gtt	atc	cgt	ctc	gcc	tgt	-	gac Page	_	tac	atc	1296

Cys Asp Ile	Ser Pro Val	Ile Arg Leu Ala 425	
	Val Ile Phe	atc ttc ggg gtc Ile Phe Gly Val 440	
	Ile Cys Ile	tcc tat ggc ttc Ser Tyr Gly Phe 455	
		ggc aaa caa aaa Gly Lys Gln Lys	
		att gtc cat tat Ile Val His Tyr 490	1476
<210> 48 <211> 491 <212> PRT <213> Homo	sapiens		

<400> 48

Met Val Thr Glu Phe Leu Leu Gly Phe Ser Ser Leu Gly Glu Ile 1 5 10 15

Gln Leu Ala Leu Phe Val Val Phe Leu Phe Leu Tyr Leu Val Ile Leu 20 25 30

Ser Gly Asn Val Thr Ile Ile Ser Val Ile His Leu Asp Lys Ser Leu 35 40 45

His Thr Pro Met Tyr Phe Phe Leu Gly Ile Leu Ser Thr Ser Glu Thr 50 60

Phe Tyr Thr Phe Val Ile Leu Pro Lys Met Leu Ile Asn Leu Leu Ser 65 70 70 75 80

Val Ala Arg Thr Ile Ser Phe Asn Cys Cys Ala Leu Gln Met Phe Phe 85 90 95

Phe Leu Gly Phe Ala Ile Thr Asn Cys Leu Leu Leu Gly Val Met Gly 100 105 110

Tyr Asp Arg Tyr Ala Ala Ile Cys His Pro Leu His Tyr Pro Thr Leu 115 120 125

Met Ser Trp Gln Val Cys Gly Lys Leu Ala Ala Ala Cys Ala Ile Gly 130 135 140

Gly Phe Leu Ala Ser Leu Thr Val Val Asn Leu Val Phe Ser Leu Pro 145 150 155 160

Phe Cys Ser Ala Asn Lys Val Asn His Tyr Phe Cys Asp Ile Ser Ala 165 170 175

Val Ile Leu Leu Ala Cys Thr Asn Thr Asp Val Asn Glu Phe Val Ile 180 185 190

Phe Ile Cys Gly Val Leu Val Leu Val Val Pro Phe Leu Phe Ile Cys 195 200 205

Val Ser Tyr Leu Cys Ile Leu Arg Thr Ile Leu Lys Ile Pro Ser Ala 210 215 220

Glu Gly Arg Arg.Lys Ala Phe Ser Thr Cys Ala Ser His Leu Ser Val 225 230 235 240

Val Ile Val His Tyr Gly Cys Ala Ser Phe Ile Tyr Leu Arg Pro Thr 245 250 255

Ala Asn Tyr Val Ser Asn Lys Asp Arg Leu Val Thr Val Thr Tyr Thr 260 265 270

Ile Val Thr Pro Leu Leu Asn Pro Met Val Tyr Ser Leu Arg Asn Lys 275 280 285

Asp Val Gln Leu Ala Ile Arg Lys Val Leu Gly Lys Lys Gly Ile Leu 290 295 300

Ser Ile Ser Glu Ile Phe Tyr Thr Thr Val Ile Leu Pro Lys Met Leu 305 310 315 320

Ile Asn Leu Phe Ser Val Phe Arg Thr Leu Ser Phe Val Ser Cys Ala 325 330 335

Thr Gln Met Phe Phe Phe Leu Gly Phe Ala Val Thr Asn Cys Leu Leu 340 345 350

Leu Gly Val Met Gly Tyr Asp Arg Tyr Ala Ala Ile Cys Gln Pro Leu 355 360 365

Gln Tyr Ala Val Leu Met Ser Trp Arg Val Cys Gly Gln Leu Ile Ala 370 375 380

Thr Cys Ile Ile Ser Gly Phe Leu Ile Ser Leu Val Gly Thr Thr Phe 385 390 400

Val Phe Ser Leu Pro Phe Cys Gly Ser Asn Lys Val Asn His Tyr Phe 405 410 415

Cys Asp 11e Ser Pro Val 11e Arg Leu Ala Cys Ala Asp Ser Tyr 11e 420 425 430

Ser Glu Leu Val Ile Phe Ile Phe Gly Val Leu Val Leu Val Val Pro 435 440 445

Lys Ile Pro Ser Ala Glu Gly Lys Gln Lys Ala Phe Ser Thr Cys Ala 465 470 475 480

Ser His Leu Ile Val Val Ile Val His Tyr Gly 485 490

<210> 49

<211> 35 <212> DNA

<213> Homo sapiens

<400> 49

ctctgaaatc ttctacacaa ctgttattct gccca

35

<210 <211 <212 <213	> ?>	50 27 DNA Homo	sap:	iens											
<400 atga		50 ggg (aagc	acago	gt g	gagaa	ag								27
<210 <211 <212 <213	!> !>	51 50 DNA Homo	sap:	iens											
<400 atca		51 ttg	ttaa	aatg	gc c	gtaci	tgtc	a aaa	agcaa	ett	acaq	gatto	caa		50
<210 <211 <212 <213	:> ?>	52 50 DNA Homo	sap	iens											
<400 atat		52 acc (aaaa	aagco	cc to	caaal	tage	c caa	agtaa	accc	taaa	agaa	aaa		50
<210 <211 <212 <213	!> ?>	53 50 DNA Homo	sap	iens											
< 4 0 0)>	53 ttc			gt g	t ggga	aata	g cto	ggcta	agcc	atcı	gcad	gaa		50
<210 <211 <212 <213	!> !>	54 50 DNA Homo	sap.	iens											
<400		54	a++>	- -	-~ ~				.~~~	ata	202				50
Cata	iagg	gtt (ctta	aaac	Lg g	yaya	yaya	d LLa	iyaa	igic	ayaç	yaaa	gay		30
<210 <211 <212 <213	! > ? >	55 276 DNA Homo	sap:	iens											
<220 <221 <222 <223	!> !>	CDS (1).	. (27	6)											
<400		55		~ ~ ~										 • • •	4.0
		gtt Val													48
		ctt Leu													96
		tgt Cys 35													144
		att Ile													- 192
		att Ile													240
gca	tct	cct	atg	gct	tca	ttg	ttc	aca	сса	tcc	tga				276

```
16U 200 PCT FINAL.ST25
Ala Ser Pro Met Ala Ser Leu Phe Thr Pro Ser
<210> 56
<211> 91
<212> PRT
<213> Homo sapiens
<400> 56
Met Thr Val Tyr Asp Ser Tyr Val Ala Ile Cys His Pro Leu His Tyr
Pro Val Leu Thr Ser Trp Gln Ile Cys Ser Phe Leu Asp Phe Gln Leu 20 25 30
Leu Phe Cys Gly Pro Asn Lys Ile Asn His Tyr Phe Cys Asp Ile Ser 35 40
Ile Phe Ile Gly Gly Ile Leu Ala Leu Thr Val Pro Leu Ile Leu Phe
Ala Ser Pro Met Ala Ser Leu Phe Thr Pro Ser
               85
<210> 57
<211> 33
<212> DNA
<213> Homo sapiens
<400> 57
atgacagttt atgattccta tgttgccatc tgc
                                                                   33
<210> 58
<211> 29
<212> DNA
<213> Homo sapiens
<400> 58
tcaggatggt gtgaacaatg aagccatag
                                                                   29
<210>
      59
<211>
      50
<212>
      DNA
<213> Homo sapiens
<400> 59
ttccctattt aataaatggt gctgggaaaa ctggctagcc atatgtagaa
                                                                   50
<210>
      60
<211>
      50
<212>
      DNA
<213> Homo sapiens
aacaacccca tcaaaaagtg ggccaaagat atgaacagac acttctcaaa
                                                                   50
<210>
      61
<211>
      50
<212>
      DNA
<213> Homo sapiens
<400> 61
aatggcgatc attaaaaagt caggaaacaa caggtgctgg agaggatgtg
                                                                  50
```

```
<210>
<211>
       50
<212>
       DNA
<213>
       Homo sapiens
<400>
       62
cccagaggat tataaatcat gctgctgtaa agacacatgc ccacgtatgt
                                                                        50
<210>
       63
<211>
       5269
<212>
       DNA
<213>
       Homo sapiens
<220>
<221>
       CDS
       (2211)..(3152)
<222>
<223>
<400> 63
taaaagcaaa aataacaact aaattcaaaa ggagataaac tataggaaag aagatttcat
                                                                       60
ctgtcatatt tcggggattc aaatatttaa agcattatta cctattttat agttacgttt
                                                                      120
tggacacaaa ggccattatg taaaatgtaa cattagttta aaataaaatt taaatgcctt
                                                                      180
agataaataa aatgcagtgt taagaaaaaa atgtgctgtc caggcatttt ggctcatgcc
                                                                      240
tgtaatctca gctactcagg aggctgaggc aggagaatct cttgaaccca ggaggcggga
                                                                      300
ggcggaggtt acagtgagcc ataatcacgc cactgcactc cagtctgggc gacagagcaa
                                                                      360
gattctgtct ccaaaaaaaa aaaaggaaag aaaagaaagag aaaagaaaaa atatgctaat
                                                                      420
taggatatct gggtttgtga tggattgtct tttgaggttg tctattttt tttgagacgg
                                                                      480
agtictogoto tgtogocoag gotggagtgo agtggogogg ggtototgot tactggaago
                                                                      540
teegecteet gggtteactg ggtteacgee atteteetge eteagectee tgagtagetg
                                                                      600
ggactacagg cgcctgccac tacgcccggg taattttttg tattttttt ttagtagaga
                                                                      660
cggggtttca ccgtgttagc caggatggtc tcaatctctt gacctcgtga tccacccgcc
                                                                      720
tctgtctccc aaagtgctgg gattacagtc gtgagccacc gcgcccggcc ttgaggttgt
                                                                      780
ctttaataca caaattcatg agtataggaa gagagggccc ttgaatatgt tggtcttgca
                                                                      840
tgtaaattaa catctttctt gataggccgt ctaaaaattt gggtgggtta tgtgaataga
                                                                      900
tataatgtct attatgatag agaaagagat tacagatatg atagcatctg agaggtgttg
                                                                      960
gacactaatt agagcaatat aatgattett teetattatt gttttetgtt tteteatgaa
                                                                     1020
atgtattcat gttgctgtac tatctcaagt ttttagttct tctccttcat aggtaataga
                                                                     1080
tggacacaat gaatatataa tgtgtcttga agggagagaa aagaaataga catggagaca
                                                                     1140
gggatagaca gagaggacct agaagaaaag ggaagtttgc aagtcagact cttacactag
                                                                     1200
ttattctgg gtaaaaagat tttcctcaat cccattctca tgtgttttat cttgatgctg
                                                                     1260
ctttctaata tatctttgtg gcagtaactg tcactggact atgtagattc tctagtctgc
                                                                     1320
ttattaattg aaagtatggt tattaatgaa gggaatgtgt tagtatctcg acctagataa
                                                                     1380
tggagcagag tttggtgcgg gtaaagggtt acatgtctag gagttcaagg atcaaaaccc
                                                                     1440
tagtcacaga tgtgtagatt ggccttcctg ggcatatcga taggaaattc aaagcttctc
                                                                     1500
tggcttctac tttgtcacct atagaataaa gaataaacaa gggtatctgt attgactatt
                                                                     1560
cgatatttat tatttctcaa gcaatgagga aggattgata attagtacag cctgattttg
                                                                     1620
gagcatacgc tctgaaacaa ttagttcagc tgtattttga agtcaaattt tctgggtcag
                                                                     1680
acaaacatta aactgctata tggaattaac aataaaggca caaatgttaa gcgttagggc
                                                                     1740
```

t 0 t			•										FIN			
										_				_	attttg	1800
													-		agatca	1860
															aataat	1920
								Ī						_	ctaact	1980
															gtttt	2040
				_											atattt	2100
						_	_								tctcct	2160
Cala	catt	ccc	gete	tcag	aa c	taac	tttc	a ga	ttcg	aaga	aac	agaa		atg Met 1	_	2216
										ttc Phe						2264
										ttc Phe						2312
										999 Gly 45						2360
aaa Lys	atc Ile	aac Asn	ccc Pro	aaa Lys 55	ctg Leu	cat His	acc Thr	ccc Pro	atg Met 60	tac Tyr	ttt Phe	ttc Phe	ctc Leu	agc Ser 65	caa Gln	2408
										atc Ile						2456
										att Ile						2504
gta Val	gta Val 100	caa Gln	ttc Phe	ttt Phe	ttc Phe	ttc Phe 105	tgt Cys	acc Thr	ttt Phe	gtg Val	gtc Val 110	act Thr	gaa Glu	tcc Ser	ttt Phe	2552
tta Leu 115	tta Leu	gct Ala	gtg Val	atg Met	gcc Ala 120	tat Tyr	gac Asp	cgc Arg	ttc Phe	gtg Val 125	gcc Ala	att Ile	tgc Cys	aac Asn	cct Pro 130	2600
ctg Leu	ctc Leu	tac Tyr	aca Thr	gtt Val 135	aac Asn	atg Met	tcc Ser	cag Gln	aaa Lys 140	ctc Leu	tgc Cys	gtg Val	ctg Leu	ctg Leu 145	gtt Val	2648
										tcc Ser						2696
tct Ser	gct Ala	tta Leu 165	aag Lys	tta Leu	tgt Cys	ttt Phe	cat His 170	ggt Gly	ttc Phe	aac Asn	aca Thr	atc Ile 175	aat Asn	cac His	ttc Phe	2744
										tct Ser						2792
										acc Thr 205						2840
aca Thr	cta Leu	ctc Leu	atc Ile	gtt Val 215	ctc Leu	aca Thr	tct Ser	tat Tyr	gcg Ala 220	ttc Phe	att Ile	gtt Val	gta Val	acc Thr 225	atc Ile	2888
ctc Leu	aag Lys	atg Met	cgt Arg 230	tca Ser	gtc Val	agt Ser	ggg Gly	cgc Arg 235	cgc Arg	aaa Lys	gcc Ala	ttc Phe	tcc Ser 240	acc Thr	tgt Cys	2936
gcc	tcc	cac	ctg	act	gcc	atc	acc	atc	ttc	cat	ggc	acc	atc	ctc	ttc	2984

16U 200 PCT FINAL.ST25 Ala Ser His Leu Thr Ala Ile Thr Ile Phe His Gly Thr Ile Leu Phe 245 250 255	
ctt tac tgt gtg ccc aac tcc aaa aac tcc agg cac aca gtc aaa gtg Leu Tyr Cys Val Pro Asn Ser Lys Asn Ser Arg His Thr Val Lys Val 260 265 270	032
Ala Ser Val Phe Tyr Thr Val Val Ile Pro Met Leu Asn Pro Leu Ile 275 280 285 290	080
tac agt ctg aga aat aaa gat gtc aag gat aca gtc acc gag ata ctg Tyr Ser Leu Arg Asn Lys Asp Val Lys Asp Thr Val Thr Glu Ile Leu 295 300 305	128
gac acc aaa gtc ttc tct tac tga gcctgttact ttcatggagt ttgtcacaca 33 Asp Thr Lys Val Phe Ser Tyr 310	182
tataaataaa ttetgteeat aaatattgat ettaaagata tetttacaaa taaacaaagt 33	242
tagggttgta caactcaacg aaatggattt tottttcaac agactaaact tagctotgto 33	302
tottacttto tgggaagcat cagtaatooc totaatottt aatatttoat ttatgaaatt 33	362
agtatagtat gggttagatc atagtctgat tgtgaagatt aaaatataat ggacacctta 3	422
cgtaagtcaa tggatattaa tttcatgtcc tttccttata agataccggg aatagactaa 30	482
gtgcttagga aacatatgaa tttcttttat aaatgtgcaa aataagttaa aagaagaaat 3	542
agtcctcatc ttcaaggatg aaaactgtgt tgataatagg acaatgaaga agtggccatt 30	602
gtgtaaggca gaaattaata tgtaccaaag agagtttgag agaagagaaa gttcaaatct 36	662
acttagggat tttagaagga tgtcttaatg aaataggtat tgtttgaaac cggcttttga 3°	722
aaaggaaatg ggcggagttt atgagataca ttccagggag aaaggagttt tcttctggag 3°	782
aaaacaatgt gaataaaacc aattaggtaa gaatgtaata cctagtcaaa gatctaatac 38	842
ttgttttatt gagctaacat aatataatgt gtgtttgtgt gtgtgtgtt gtgtgtgt	902
atgtatacgg gttacttgac atgaactgaa attttaatat gatatgggct acatcctgaa 39	962
tgtgttttca aaggagetee agtgtgaeea tetgataaae cataatagae tteaceagat 40	022
gctgatgaat aatggatcag atcttagaaa atcctatgta ccaaattagg gatgatgaac 40	082
acctgcccaa cctgtatgtc atactgttag acatcacaaa ctttatcaat ccattatgat 41	142
ttttttatga gcatggaaat aatctctgaa tccttctcaa cagaattccc aacaaccttt 42	202
ataaaaaggt atttggagta gtcttaagtg ttgaaagctc tttggctgca taaacttatt 42	262
caaaataaat aaaaatcagg taatcattaa tatcaagacc tetttaacac agcaaattaa 43	322
aaatgctagc tctttcttac cttaataact cactttcatt cgaataaatt gtataccctt 43	382
ctccttttca atgtgtctag atacagttcc aaacaaatca tcaatatagt ggaagaagta 44	142
aatttccagg tgttttgtta agggagaaaa aataaactgg ggaacaattt tatataaact 45	502
tcttaaattt atttagaatg ttcatattat tttgacctta tgatgattat taaagttatg 45	62
ataattatta aagtgattca tcttacatat attatttgat aaagaatcca ctaaataatc 46	522
cttgtaatag aaaaattttt caaaatgtaa ggaacagtgt tttagatatt aaatgcctga 46	582
ggagggaata cttttctct tgatatctgt atctccaggt attcaaacat ttatcctttg 47	742
tacacatctg gtacttatac aatttttaat tttctcagaa gttgggacat tgttitaata 48	302
ttaaatcgaa tactgaattt caccatcttt tgaaatcctg aaaagctgcc atgggaacaa 48	362
gcataaaata ggatatttga taatgaggaa aattagccca tatccccatc acaagggctt 49	22
ttctctggca acctaccaga cttgagtgtg aagccctgtg agatgatctg acctgccage 49	82

50425102516252225269

160 200 per EINAL ST25	
16U 200 PCT FINAL.ST25 tgacagtata cacacaagca agacaagcca agatcagcca tgccagggac atgtgagcag	1
aaccacctgg atgatccatg caatagagtc acaggcaata agaggttgtt gtgtttagcc	:
aataggtttt ggagtggttt gttatacagt cattagagtc attcactttc tcaaattctg	ţ
ggaatgctac caaaaaaact tcacgatgtt ttattatgta ataattcacc atcttctatt	•
attccacatt gaggaaacat tttaaaataa taaaatgtgt taaattt	
<210> 64 <211> 313 <212> PRT <213> Homo sapiens	
<400> 64	
Met Leu Leu Thr Asp Arg Asn Thr Ser Gly Thr Thr Phe Thr Leu Leu 1 5 10 15	
Gly Phe Ser Asp Tyr Pro Glu Leu Gln Val Pro Leu Phe Leu Val Phe 20 25 30	
Leu Ala Ile Tyr Asn Val Thr Val Leu Gly Asn Ile Gly Leu Ile Val 35 40 45	
Ile Ile Lys Ile Asn Pro Lys Leu His Thr Pro Met Tyr Phe Phe Leu 50 60	
Ser Gln Leu Ser Phe Val Asp Phe Cys Tyr Ser Ser Ile Ile Ala Pro 65 70 75 80	
Lys Met Leu Val Asn Leu Val Val Lys Asp Arg Thr Ile Ser Phe Leu 85 90 95	
Gly Cys Val Val Gln Phe Phe Phe Phe Cys Thr Phe Val Val Thr Glu 100 105 110	
Ser Phe Leu Leu Ala Val Met Ala Tyr Asp Arg Phe Val Ala Ile Cys 115 120 125	
Asn Pro Leu Leu Tyr Thr Val Asn Met Ser Gln Lys Leu Cys Val Leu 130 135 140	
Leu Val Val Gly Ser Tyr Ala Trp Gly Val Ser Cys Ser Leu Glu Leu 145 150 150 160	
Thr Cys Ser Ala Leu Lys Leu Cys Phe His Gly Phe Asn Thr Ile Asn 165 170 175	
His Phe Phe Cys Glu Phe Ser Ser Leu Leu Ser Leu Ser Cys Ser Asp 180 185 190	
Thr Tyr Ile Asn Gln Trp Leu Leu Phe Phe Leu Ala Thr Phe Asn Glu 195 200 205	
Ile Ser Thr Leu Leu Ile Val Leu Thr Ser Tyr Ala Phe Ile Val Val 210 220	
Thr Ile Leu Lys Met Arg Ser Val Ser Gly Arg Arg Lys Ala Phe Ser 225 235 230 240	
Thr Cys Ala Ser His Leu Thr Ala Ile Thr Ile Phe His Gly Thr Ile	

50

16U 200 PCT FINAL.ST25 245 Leu Phe Leu Tyr Cys Val Pro Asn Ser Lys Asn Ser Arg His Thr Val 265

Lys Val Ala Ser Val Phe Tyr Thr Val Val Ile Pro Met Leu Asn Pro

Leu Ile Tyr Ser Leu Arg Asn Lys Asp Val Lys Asp Thr Val Thr Glu

Ile Leu Asp Thr Lys Val Phe Ser Tyr 310

<210> 65 <211> 50 <212> DNA

<213> Homo sapiens

<400> 65

ataggccgtc taaaaatttg ggtgggttat gtgaatagat ataatgtcta 50

<210> 66 <211> 50 <212> DNA

<213> Homo sapiens

<400> 66 acacaatgaa tatataatgt gtcttgaagg gagagaaaag aaatagacat

<210> 67 <211> 32 <212> DNA

Homo sapiens

<400> 67

atgctgctga ctgatagaaa tacaagtggg ac 32

<210> 68 <211> 29 <212> DNA

<213> Homo sapiens

<400> 68

gactttggtg tccagtatct cggtgactg 29

<210> 69 <211> 4558

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

(1822)..(2766) <222>

<223>

<400> 69

gggaactcag agtagaagct agacagttga gcattccctq aatattgatt tctcttqqca 60 tttttaccac ctaagagagc acctagagaa gtatgagaat aaaagggcaa caaaaagagg 120 agagaaaaga agagagaga aggggaatac acaagcaatg ataagatcat ataaggggag 180 tagaggagat gggagctaga ttggaaggaa ttaaaaatta agagatgagg aattagataa 240 tgaattactc aactttttca acaaacattg ttgtgaagga aaacagttgc gggtgttaga 300 tggagggaga tataaggcca aggaatgcgg gattgtgctg atgtcaagaa aaacatttta 360 aaaaaggggg cagatttgtg tgggaggggg aatattgtaa tgacaaataa cagttttaca 420

atgctttaca gtttacaggg	actettetea tgattttate	tttcctcatc atgaagcagt	480
aaaattgtca tggaagatat	ggttatgcct tcatctatag	gtgaggaacc taaatcttta	540
aaaatttcag tgattaatac	tagtttctat tggtaggctg	atggctcaat cattacttaa	600
acccaagtct ttagattcta	atttttttc ctatatcatg	ataatgagtt tgagttatat	660
ttctgttaac ttgaatatgc	tggatttata cttcttagga	agcaatgagg tagaagcaga	720
agagtgtagt gatttaagac	attggattgg gaggcagcaa	accagagttc tcgaagctca	780
agtttcatct ttgacgtttt	ccaactgttc tgattgttag	tgacttagtc ttttgttttc	840
tcaattcata agtgtccacg	tttactgagc actgtttcaa	gatttgtgct aagtgtttta	900
aaagatotoa aaatooccaa	aagaaagttt ttaggcagga	gctgaaaaaa aaggtggcac	960
aggtcaaaaa tattgcaagg	aaatgtttaa acgttttcaa	gggaaatgac aacagaaggt	1020
ggaaaaagat agaatgataa	ggatcccaga tggaaaaata	gtgtgaaggg aataggtcag	1080
tcttgcaaaa aggtaagtgt	ggggcatctc ttctgaatgt	catgaagtcc aggaaggaag	1140
aagcacgtag agatgaaggt	taaactacag ttaggcaaaa	gagaacaaca aaagggcttc	1200
tcatgttctc aagttacctg	gaggtagggc tttagttgga	gggatttctg agtgtcaaac	1260
aaggacctac gaaaccctgc	tagaaaaaaa atctaaagaa	cttgtaggga agtgaattat	1320
tagaaagtgc taactacatt	tattttcat atgaccaaga	ttgacatttc agggcagaaa	1380
ctctttcata attgggagtg	tagtttgaat tggaaggcaa	taggaagaca gtgacggtaa	1440
atgtttgtgg gtatgatttg	aattaagcag caatctgtat	tatttacaaa gttgcttttg	1500
gccacatgca ggccacaaaa	ggctcttcct ccacttgatt	tctcaataag gctgctttgt	1560
aatactagct ttattggaat	taaatgtcct gagcacccag	tgtttttata aacagcttaa	1620
gggcaaggat catgcataat	atttcatgat acatatgatt	attttctcat ttcttttcat	1680
gtctaaaaat gggtctaaga	actaatotto toacaagaat	gatcagagtt tgaatgtgag	1740
cattgtaatt ctgctgatat	tgaatattct ctggaagggc	cctgtggaag cagataagga	1800
ggaagagaat teccaggage	c atg tca gcc tcc aat Met Ser Ala Ser Asn 1 5		1851
	g gtg ggg att cca ggc u Val Gly Ile Pro Gly 20		1899
71 6 71 6 71 7	t ttc tgc tta gca tat o Phe Cys Leu Ala Tyr 35		1947
gga aac tgc act ctc ct Gly Asn Cys Thr Leu Le 45	t ctc atc atc cag gct u Leu Ile Ile Gln Ala 50		1995
gaa ccc atg tac ctc tt Glu Pro Met Tyr Leu Ph 60			2043
	g ccc aaa atg ctt gcc u Pro Lys Met Leu Ala 85		2091
gat cgg gag ata aac tt Asp Arg Glu Ile Asn Ph 95			2139
cac tcc ttc tcc atc at His Ser Phe Ser Ile Me 110			2187

Page 34

				160	200 PCT	FINAL.ST25	
gac cgc ta Asp Arg Ty 12	r Val Ala	Ile Cys I	aag cca Lys Pro 130	ctg cac	tac acc	aag gtc ctg Lys Val Leu	2235
						gcc cgg gct Ala Arg Ala	2283
						ttc cac tac Phe His Tyr 170	2331
						atg gct gtg Met Ala Val 185	2379
						tat ggc atc Tyr Gly Ile 200	2427
gct gtg gcd Ala Val Ala 20	a Met Phe	Ile Val V	gtg ttg Val Leu 210	gac ctg Asp Leu	ctc ctt Leu Leu 215	gtt atc ctg Val Ile Leu	2475
						tct cag gag Ser Gln Glu	2523
	r Lys Ala					ggt gcc atc Gly Ala Ile 250	2571
						cac cgt gta His Arg Val 265	2619
						aat ttc tat Asn Phe Tyr 280	2667
	e Pro Pro	Met Val A				gtc aag acc Val Lys Thr	2715
						aag gat atg Lys Asp Met	2763
tag agggtga	aggt ggaga	aagaa tgg	gttggct	tgtctgc	tgg agtt	ggagac	2816
aggctatggt	agaatgtgc	a cggctgc	cag gat	cttcatg	tttagttt	tt tcttggaaaa	2876
aaaaaaaatg	atgtcctga	a actcaga	gcc acc	agtctgt	tcaggact	ca tgggtctgtg	2936
tcctctggta	gcctgtgga	t tgaatgt	gct gac	tgtgctg	tcttctca	ca gtgccctcac	2996
ccctatcagt	aacttgaca	g agacttg	acc cat	gggtctc	caggtgac	tt caccgaaaga	3056
cacaaagatg	cttccaact	t catttgc	tga aga	gaagact	ttgaaaat	ct gagtttcttt	3116
tcttagtcat	tgggaattt	g gtgaact	atc tac	tcaggac	ctgggtga	gg gccaacagta	3176
tatctgacat	aggaatcct	t cattcat	tct gac	tggtggt	gtccagct	tc tgatgaaaca	3236
ctcagtgtta	ggaagtttg	a aacattc	cag ggc	tgcaggt	tctgagta	ag acacctatgc	3296
ttgctagaaa	atcattttt	t cacctaa	gcc agt	atgtgta	ttcttttg	ct tatatttacc	3356
aatccatcct	tatgtccaa	t tcctttt	att aag	tactttg	aataagac	at gtcctctggc	3416
tttatgtttc	atgcaactc	t ttctttg	cac ata	gatgtat	cttatgtt	tt caagaatgag	3476
aatggctcat	ttatttacta	a attccac	caa atc	tgtgata	ggggatgg	gg acacatatac	3536
taaattaggg	gtgtcagac	t tgtgtat	ttg tcc	taagaca	gagaagga	aa tgataattat	3596
gatagattct	gttctctgaa	a atttcca	tcc caa	ggcccag	cataataa	aa gaagagcaag	3656
accaagcaga	taggaggca	a gaatatt	atg ttt	ctctttt	cctgtctc	at gtgaacttac	3716

ctactatata	attctctatt	aatcctgaca	acacagctaa	gcttttcaca	caagccctgt	3776
ataaatacat	tgttctgctg	ttatcttctg	acccacttgt	ttcctcagat	attattgctt	3836
agaaattata	tatctctttt	gctatcactg	tatctttctc	tatttaccta	tctatattat	3896
ttagccttga	aagataattt	ccaagcctat	ttcaggtggg	gtgtagaagg	ttggaagctg	3956
tccaggaggg	aagagtatag	caagaaccta	gagttttact	tccccttata	ttccacctct	4016
gctcttataa	ttccctttga	cacaaaaaca	aataccccag	agaaataatg	tattacataa	4076
aaaattgcta	catgctagat	atatatattt	ttggagtata	tgtgatattc	tgatatattc	4136
atataataga	taatgatcaa	atcaggataa	ttggaatatc	catgacctta	aatgtttctt	4196
ttatgctagg	aacattaaaa	ttattctctt	ctagctattt	tgatatatac	agtagattgt	4256
tttctatagt	ccctactgat	ttcttgaaca	ctacatcttg	ttatttttta	tatctagctg	4316
tatttttata	ctcaattaat	ctcttatcct	ccctgcctcc	cttcccagcc	cccaataacc	4376
accaatctgc	tctctatttt	catgagctgt	acttagcatc	cacatgagtg	agaaatacaa	4436
taattgtctt	tctgtacctg	gcttgtttca	cttaacttaa	tgacctacag	tttcatccac	4496
gttgctgcaa	gtgacaggat	ttcattcttt	cttatgacta	atattccatg	tgtatcatat	4556
tt						4558

<210> 70 <211> 314 <212> PRT

<213> Homo sapiens

<400> 70

Met Ser Ala Ser Asn Ile Thr Leu Thr His Pro Thr Ala Phe Leu Leu 1 5 5 15 15 10 10 15 15

Val Gly Ile Pro Gly Leu Glu His Leu His Ile Trp Ile Ser Ile Pro $20 \hspace{1cm} 25 \hspace{1cm} 30$

Phe Cys Leu Ala Tyr Thr Leu Ala Leu Leu Gly Asn Cys Thr Leu Leu 40 $$45\$

Leu Ile Ile Gln Ala Asp Ala Ala Leu His Glu Pro Met Tyr Leu Phe 50 60

Leu Ala Met Leu Ala Ala Ile Asp Leu Val Leu Ser Ser Ser Ala Leu 65 70 75 80

Pro Lys Met Leu Ala Ile Phe Trp Phe Arg Asp Arg Glu Ile Asn Phe 85 90 95

Phe Ala Cys Leu Ala Gln Met Phe Phe Leu His Ser Phe Ser Ile Met 100 105 110

Glu Ser Ala Val Leu Leu Ala Met Ala Phe Asp Arg Tyr Val Ala Ile 115 120 125

Cys Lys Pro Leu His Tyr Thr Lys Val Leu Thr Gly Ser Leu Ile Thr 130 $^{\circ}$ 140 $^{\circ}$

Lys Ile Gly Met Ala Ala Val Ala Arg Ala Val Thr Leu Met Thr Pro 145 150 155 160

Leu Pro	Phe 1	Leu	Leu 165	Arg _.	Cys	Phe	His					Pro				
Ala His		Tyr 180	Cys	Glu	His	Met	Ala 185	Val	Val	Arg	Leu	Ala 190	Cys	Gly		
Asp Thr	Ser 1 195	Phe	Asn	Asn	Ile	T yr 200	Gly	Ile	Ala	Val	Ala 205	Met	Phe	Ile		
Val Val 210	Leu /	Asp	Leu	Leu	Leu 215	Val	Ile	Leu	Ser	Туг 220	Ile	Phe	Ile	Leu		
Gln Ala 225	Val 1	Leu	Leu	Leu 230	Ala	Ser	Gln	Glu	Ala 235	His	Туr	Lys	Ala	Phe 240		
Gly Thr	Cys '		Ser 245	His	Ile	Gly	Ala	Ile 250	Leu	Ala	Phe	Tyr	Thr 255	Thr		
Val Val		Ser 260	Ser	Val	Met	His	Arg 265	Val	Ala	Arg	His	Ala 270	Ala	Pro		
His Val	His :	Ile	Leu	Leu	Ala	Asn 280	Phe	Tyr	Leu	Leu	Phe 285	Pro	Pro	Met		
Val Asn 290	Pro :	Ile	Ile	Tyr	Gly 295	Val	Lys	Thr	Lys	Gln 300	Ile	Arg	Glu	Ser		
Ile Leu 305	Gly v	Val	Phe	Pro 310	Arg	Lys	Asp	Met								
<211> 5	71 50 ONA Homo :	sapi	ens													
<400> 7	71 att ti	taaa	iaaaç	gg gg	gcaç	gatti	t g t ç	jtggç	gagg	ggga	aatat	tg				50
<211> 5 12	72 50 DNA domo s	sapi	ens													
<400> 7 gctaagtg	72 gtt ti	taaa	agat	c to	aaaa	atcco	c caa	aaga	aaag	ttti	tago	jca				50
<211> 5 12	73 50 ONA Homo s	sapi	ens													
<400> 7 gcaggago	13 ctg aa	aaaa	aaaç	gg tg	gcac	aggt	caa	aaat	att	gcaa	aggaa	aat				50
<212> [14 50 NA Iomo s	sapi	ens										ı		.*	
<400> 7 cccagtgt	74 :tt tt	tata	aaca	ıg ct	taag	gggca	a ago	jatca	tgc	ataa	atatt	tc				50
	75 50															

16U 200 PCT FINAL.ST25	
<400> 75 ttttcatgtc taaaaatggg tctaagaact aatcttctca caagaatgat	50
<210> 76	
<211> 29	
<212> DNA <213> Homo sapiens	
<400> 76	
cagootocaa tatoacotta acacatoca	29
<210> 77	
<211> 25	
<212> DNA <213> Homo sapiens	
<400> 77 caccacaata aacatggcca cagcg	25
caccaca addatygeed cageg	23
<210> 78	
<211> 2520 <212> DNA	
<213> Homo sapiens	
<220> <221> CDS	
<222> (727)(1722)	
<223>	
<400> 78 attaaagtet teagteteea catteeetae ttteeaaatt eagettteee gggaggtetg	60
gagcagctgc ctctctgggg agatgctgga ggtctcggaa tcacctcacg cggcctcagg 1	20
gcccagttgg agccacccca agtgacacca gcaggcagat gaccagagag cctgagcctc 1	80
cggccccgag tctgtgaagc ctagccgctg ggctggagaa gccactgtgg gcaccaccgt 2	40
gggggaaaca ggcccgttgc cctggcctct ttgccctggg ccagcctttg tgaagtgggc 3	00
ccctcttctg ggccccttga agcatgctgg agaacttctc ggccgccgtg cccagccacc 3	60
gctgctgggc accectectg gacaacagca eggeteagge eageatecta gggagettga 4	20
gtcctgaggc cctcctggct atttccatcc cgccgggccc caaccagagg ccccaccagt 4	80
gccgccgctt ccgccagcca cagtggcagc tettggaccc caatgccacg gccaccagct 5	40
ggagcgaggc cgacacggag ccgtgtgtgg atggctgggt ctatgaccgc agcatcttca 6	00
cctccacaat cgtggccaag tggaacctcg tgtgtgactc tcatgctctg aagcccatgg 6	60
cccagtccat ctacctggct gggattctgg tgggagctgc tgcgtgcggc cctgcctcag $-\%$	20
	68
Met Glu Trp Thr Ala Ala Arg Ala Arg Pro Leu Val Met Thr 1 5 10	
	16
Leu Asn Ser Leu Gly Phe Ser Phe Gly His Gly Leu Thr Ala Ala Val 15 20 25 30	
gcc tac ggt gtg cgg gac tgg aca ctg ctg cag ctg gtg gtc tcg gtc 8	64
Ala Tyr Gly Val Arg Asp Trp Thr Leu Leu Gln Leu Val Val Ser Val 35 40 45	
	12
Pro Phe Phe Leu Cys Phe Leu Tyr Ser Trp Trp Leu Ala Glu Ser Ala	
50 55 60	
Cga tgg ctc ctc acc aca ggc agg ctg gat tgg ggc ctg cag gag ctg Arg Trp Leu Leu Thr Thr Gly Arg Leu Asp Trp Gly Leu Gln Glu Leu	60
65 70 75	

								aag Lys		gca	gtg	cag		acc	ctg	1008
								atg Met								1056
								ctg Leu								1104
								tgc Cys 135								1152
								cag Gln								1200
								gtg Val								1248
								ggc Gly								1296
								att Ile								1344
								gcc Ala 215								1392
								atc Ile								1440
								gca Ala								1488
								cct Pro								1536
								gtg Val								1584
								ccc Pro 295								1632
								aac Asn								1680
cat His	ggc Gly 320	acg Thr	ctg Leu	Gl y ggg	aac Asn	tct Ser 325	gtc Val	cta Leu	aaa Lys	tcc Ser	aca Thr 330	cag Gln	ttt Phe			1722
tago	ctcc	tg ç	ggaa	cctg	c ga	tggg	acgg	, tca	gagg	aag	agac	ttct	tc t	gtto	tctgg	1782
agaa	ggca	igg a	ggaa	agca	a ag	acct	ccat	ttc	caga	ggc	ccag	aggo	tg c	ccto	tgagg	1842
tccc	cact	ct c	cccc	aggg	c tg	cccc	tcca	ggt	gago	cct	gccc	ctct	ca c	agto	caagg	1902
ggcc	ccct	tc a	atac	tgaa	g gg	gaaa	agga	cag	tttg	att	ggca	ggag	gt g	accç	agtgc	1962
acca	tcac	cc t	gccc	tgcc	c tc	gtgg	cttc	gga	gago	aga	9 999	tcag	gc c	cagg	ggaac	2022
gago	tggc	ct t	gcca	accc	t ct	gctt	gact	ccg	cact	gcc	actt	gtcc	cc c	caca	cccgt	2082
ccac	ctgc	сс а	gago	tcag	a gc	taac	cacc	atc	catg	gtc	aaga	cctc	tc c	tago	tccac	2142
acaa	gcag	ta g	agtc	tcag	c tc	caca	gctt	tac	ccag	aag	ccct	gtaa	gc c	tggc	ccctg	2202

gcccctcccc	atgtccctcc	aggcctcagc	cacctgcccg	ccacatcctc	tgcctgctgt	2262
ccccttccca	ccctcatccc	tgaccgactc	cacttaaccc	ccaaacccag	cccccttcc	2322
aggggtccag	ggccagcctg	agatgcccgt	gaaactccta	cccacagtta	cagccacaag	2382
cctgcctcct	cccaccctgc	cagcctatga	gttcccagag	ggttggggca	gtcccatgac	2442
cccatgtccc	agctccccac	acagcgctgg	gccagagagg	cattggtgcg	agggattgaa	2502
taaagaaaca	aatgaatg					2520

<210> 79 <211> 332

<211> 332 <212> PRT

<213> Homo sapiens

<400> 79

Met Glu Trp Thr Ala Ala Arg Ala Arg Pro Leu Val Met Thr Leu Asn 1 5 10 15

Ser Leu Gly Phe Ser Phe Gly His Gly Leu Thr Ala Ala Val Ala Tyr $20 \hspace{1cm} 25 \hspace{1cm} 30$

Gly Val Arg Asp Trp Thr Leu Leu Gln Leu Val Val Ser Val Pro Phe 35 40 45

Phe Leu Cys Phe Leu Tyr Ser Trp Trp Leu Ala Glu Ser Ala Arg Trp 50 60

Leu Leu Thr Thr Gly Arg Leu Asp Trp Gly Leu Gln Glu Leu Trp Arg 65 70 75 80

Val Ala Ala Ile Asn Gly Lys Gly Ala Val Gln Asp Thr Leu Thr Pro 85 90 95

Glu Val Leu Ser Ala Met Arg Glu Glu Leu Ser Met Gly Gln Pro $100 \,$ $105 \,$ $110 \,$

Pro Ala Ser Leu Gly Thr Leu Leu Arg Met Pro Gly Leu Arg Phe Arg 115 120 125

Thr Cys Ile Ser Thr Leu Cys Trp Phe Ala Phe Gly Phe Thr Phe Phe 130 $$135\$

Gly Leu Ala Leu Asp Leu Gln Ala Leu Gly Ser Asn Ile Phe Leu Leu 145 150 155 160

Gln Met Phe Ile Gly Val Val Asp Ile Pro Ala Lys Met Gly Ala Leu 165 170 175

Leu Leu Ser His Leu Gly Arg Arg Pro Thr Leu Ala Ala Ser Leu 180 $$195\$

Leu Leu Ala Gly Leu Cys Ile Leu Ala Asn Thr Leu Val Pro His Glu 195 200 205

Met Gly Ala Leu Arg Ser Ala Leu Ala Val Leu Gly Leu Gly Gly Val 210 215 220

Gly Ala Ala Phe Thr Cys Ile Thr Ile Tyr Ser Ser Glu Leu Phe Pro 225 230 235 240

Thr	Val	Leu	Arg	Met 245	Thr	Ala	Val	Gly	Leu 250	Gly	Gln	Met	Ala	Ala 255	Arg	
Gly	Gly	Ala	11e 260	Leu	Gly	Pro	Leu	Val 265	Arg	Leu	Leu	Gly	Val 270	His	Gly	
Pro	Trp	Leu 275	Pro	Leu	Leu	Val	Tyr 280	Gly	Thr	Val	Pro	Val 285	Leu	Ser	Gly	
Leu	Ala 290	Ala	Leu	Leu	Leu	Pro 295	Glu	Thr	Gln	Ser	Leu 300	Pro	Leu	Pro	Asp	
Thr 305	Ile	Gln	Asp	Val	Gln 310	Asn	Gln	Ala	Val	Lys 315	Lys	Ala	Thr	His	Gly 320	
Thr	Leu	Gly	Asn	Ser 325	Val	Leu	Lys	Ser	Thr 330	Gln	Phe					
<210 <211 <212 <213	l > = 1 2 > = 1	80 2250 DNA Homo	sap:	iens												
<220 <221 <222 <223	l> (2>	CDS (10)	(7	38)												
<400 caaç												n Le			ca gtg er Val	51
						gcc Ala										99
Gly 15 ccc	Ser	Gln	Asp	Ala	Leu 20 tgg		Pro	Leu	Pro tgt	Pro 25 gga	Pro tct	Ala	Pro	Gln tgg	Asn 30 ggc	99 147
Gly 15 ccc Pro	Ser tcc Ser	Gln acc Thr	Asp cac His	tct Ser 35	Leu 20 tgg Trp	Ala	Pro cct Pro	ttg Leu cat	tgt Cys 40	Pro 25 gga Gly ttg	Pro tct Ser	Ala ctg Leu	Pro cct Pro	Gln tgg Trp 45	Asn 30 ggc Gly ctg	
Gly 15 ccc Pro ctc Leu	tcc Ser agc Ser	Gln acc Thr tgt Cys	cac His ctt Leu 50	tct ser 35 ctg Leu	Leu 20 tgg Trp gct Ala	Ala gac Asp	Pro cct Pro cag Gln ctg	ttg Leu cat His 55	tgt Cys 40 gtc Val	Pro 25 gga Gly ttg Leu	tct Ser gtc Val	Ala ctg Leu atg Met	Pro cct Pro gct Ala 60 cca	tgg Trp 45 tct ser	Asn 30 ggc Gly ctg Leu	147
Gly 15 ccc Pro ctc Leu ctc Leu	tcc Ser agc Ser tgt Cys	acc Thr tgt Cys gtc Val 65	cac His ctt Leu 50 tcc Ser	tct Ser 35 ctg Leu cac His	Leu 20 tgg Trp gct Ala ctg Leu	Ala gac Asp ctg Leu	CCt Pro cag Gln ctg Leu 70	ttg Leu cat His 55 ctt Leu	tgt Cys 40 gtc Val tgc Cys	Pro 25 gga Gly ttg Leu agt Ser	tct Ser gtc Val ctc Leu	Ala ctg Leu atg Met tcc Ser 75	Pro cct Pro gct Ala 60 cca Pro	tgg Trp 45 tct Ser gga Gly	Asn 30 ggc Gly ctg Leu gga Gly	147 195
Gly 15 ccc Pro ctc Leu ctc Leu	tcc ser agc ser tgt Cys tct ser atg	Gln acc Thr tgt Cys gtc Val 65 tac Tyr	Cac His Ctt Leu 50 tcc Ser tcc	tct Ser 35 ctg Leu cac His	Leu 20 tgg Trp gct Ala ctg Leu tct Ser	Ala gac Asp ctg Leu ctc Leu	Pro cct Pro cag Gln ctg Leu 70 ctc Leu act	ttg Leu cat His 55 ctt Leu ctg Leu	tgt Cys 40 gtc Val tgc Cys gcc Ala	Pro 25 gga Gly ttg Leu agt ser tcc ser	tct Ser gtc Val ctc Leu agc ser 90	Ala ctg Leu atg Met tcc Ser 75 ttc Phe agg	Pro cct Pro gct Ala 60 cca Pro ttt Phe	tgg Trp 45 tct Ser gga Gly tca Ser	Asn 30 ggc Gly ctg Leu gga Gly tgt Cys	147 195 243
Gly 15 ccc Pro ctc Leu ctc Leu ggt Gly 95	tcc Ser agc Ser tgt Cys tct Ser 80 atg Met cag	Gln acc Thr tgt Cys gtc Val 65 tac Tyr tct Ser	cac His ctt Leu 50 tcc Ser tcc Ser acc Thr	tct Ser 35 ctg Leu cac His cct Pro	tgg Trp gct Ala ctg Leu tct Ser	gac Asp ctg Leu ctc Leu cag Gln 85 caa Gln	Pro cct Pro cag Gln ctg Leu 70 ctc Leu act Thr	ttg Leu cat His 55 ctt Leu ctg Trp	tgt Cys 40 gtc Val tgc Cys	Pro 25 gga Gly ttg Leu agt Ser tcc Ser ggc Gly 105 cct	tct Ser gtc Val ctc Leu agc Ser 90 agc Ser	Ala ctg Leu atg Met tcc Ser 75 ttc Phe agg Arg	Pro cct Pro gct Ala 60 cca Pro ttt Phe ctg Leu	tgg Trp 45 tct Ser gga Gly tca Ser	Asn 30 ggc Gly ctg Leu gga Gly tgt Cys ctt Leu 110 acc	147 195 243
Gly 15 ccc Pro ctc Leu ctc Leu ggt Gly 95 gtc Val	tcc Ser agc Ser tgt Cys tct Ser 80 atg Met cag Gln cag	Gln acc Thr tgt Cys gtc Val 65 tac Tyr tct Ser gct Ala aag	Cac His ctt Leu 50 tcc Ser ccc Thr cca Pro cta	tct Ser 35 ctg Leu cac His cct Pro atc Ile tcc Ser 115 ccc	tgg Trp gct Ala ctg Leu tct Ser ctg Leu 100 tta Leu	gac Asp ctg Leu ctc Leu cag Gln 85 caa Gln	Pro cct Pro cag Gln ctg Leu 70 ctc Leu act Thr	ttg Leu cat His 55 ctt Leu ctg Trp	tgt Cys 40 gtc Val tgc Cys gcc Ala atg Met	Pro 25 gga Gly ttg Leu agt Ser tcc Ser ggc Gly 105 cct Pro	tct Ser gtc Val ctc Leu agc Ser gct Ala	Ala ctg Leu atg Met tcc Ser 75 ttc Phe agg Arg ctg Leu aac	Pro cct Pro gct Ala 60 cca Pro ttt Phe ctg Leu gtg Val	tgg Trp 45 tct Ser gga Gly tca Ser cct Pro	Asn 30 ggc Gly ctg Leu gga Gly tgt Cys ctt Leu 110 acc Thr	147 195 243 291
Ctc Pro ctc Leu ctc Leu ggt Gly 95 gtc Val	tcc Ser agc Ser tgt Cys tct Ser 80 atg Met cag Gln ctg	Gln acc Thr tgt Cys gtc Val 65 tac Tyr tct Ser gct Ala aag Lys cac	cac His ctt Leu 50 tcc Ser tcc Thr cca Pro cta Leu 130 ctt	tct Ser 35 ctg Leu cac His cct Pro atc 11e tcc Ser 115 ccc Pro tgt	tgg Trp gct Ala ctg Leu tct Ser ctg Leu tct ctg Arg	Ala gac Asp ctg Leu ctc Leu cag Gln gag Glu gcc Ala gga	Pro cct Pro cag Gln ctg Leu 70 ctc Leu act Thr ttc Phe atc Ile	ttg Leu cat His 55 ctt Leu ctg Trp ctt Leu	tgt Cys 40 gtc Val tgc Cys gcc Ala atg Met	Pro 25 gga Gly ttg Leu agt Ser tcc Ser ggc Gly 105 cct Pro cct Pro cat	tct Ser gtc Val ctc Leu agc Ser gct Ala gga Gly	Ala ctg Leu atg Met tcc Ser 75 ttc Phe agg Arg ctg Leu aac Asn	Pro cct Pro gct Ala 60 cca Pro ttt Phe ctg Leu gtg Val tcc Ser 140	tgg Trp 45 tct Ser gga Gly tca Ser cct Pro	Asn 30 ggc Gly ctg Leu gga Gly tgt Cys ctt Leu 110 acc Thr ctc Leu	147 195 243 291 339

160	165	16U 200 PCT FINAL.ST25 170	
	y Leu Leu Gly Ser	c ccc ggc cac gtg ttc ccc Pro Gly His Val Phe Pro 185 190	579
		c ctg gtt gtg gca ggg ctc Leu Val Val Ala Gly Leu) 205	627
		ttc aca cac tgg ggg ttg phe Thr His Trp Gly Leu 220	675
		ggg atg gtg ccc agt ggg Gly Met Val Pro Ser Gly 235	723
ggt gta tgg ggg gac ta Gly Val Trp Gly Asp 240	ggggaggg cagaactg	gct ggtcctatca gattcagcag	778
cgactggaat agggacatat	tttatatttg gaatco	caaga cttttccttg attcatctgg	838
totoottgaa tttoacactg	ttttctgctg tcccc	aagg tcacttccta ttccttccat	898
gggagtttcc ttctctggta	tcaccccccg ctctta	itgat attotgocca otoccacoto	958
ctttcccatc cctcaggata (cccactgcct cttgct	ccta aagcettetg teteetaggg	1018
ttatcctgct catggtggtc	tgttctcagc acctgg	gete etgecagttt catgtgtgee	1078
cctggaggcg agcttcaacg (catcaactc acactc	ctct ccctgtcttc cggctccttt	1138
		gaag ggctggcctg gagtgctcac	1198
		aagg ctggctcaga aggttctggg	1258
		tgat cccagtggcc tgtgtgtgga	1318
		agga actgtctgcc cccaccaagg	1378
		attg gcctttgctg acgcccagag	1438
		ctcc acatgcctgc agtcgagggc	1558
		tgct gggaagccc atgggcactg	1618
		tcca ggtacgtgga cctgggatgg	1678
		gaca ggaacttaca ccgattgatt	1738
		cttg gggaggtgcc taagattaga	1798
		cacg gaactcctgt gcatgaggca	1858
		agtt ctgctctctt ggggacttca	1918
		cttt ttaaaaaagt gggccaggca	1978
tggtggctca cacctgtaat o	ctagcacgt ggggag	gctg aggcaggcag atcacttagg	2038
ctaggaattc aagaccagcc t	ggccaacat ggtgaa	accc tgtctctact aaaaatacaa	2098
aaattagctg ggcatggtgt t	gtgcaccta taattc	cage tacteaggag getgaggeag	2158
gagaatcgct tgagcctggg a	iggcagaggt tgctgt	gage egagacegea ceactgeact	2218
ccagcctggg cgacagagcg a	gactccatc tc	•	2250
<210> 01		.1	

<210> 81 <211> 243 <212> PRT <213> Homo sapiens

<400> 81

Page 42

Met Ser Arg Ser Pro Leu Asn Pro Ser Gln Leu Arg Ser Val Gly Ser 1 5 10 15

Gln Asp Ala Leu Ala Pro Leu Pro Pro Pro Ala Pro Gln Asn Pro Ser 20 25 30

Thr His Ser Trp Asp Pro Leu Cys Gly Ser Leu Pro Trp Gly Leu Ser 35 40 45

Cys Leu Leu Ala Leu Gln His Val Leu Val Met Ala Ser Leu Leu Cys 50 60

Val Ser His Leu Leu Leu Cys Ser Leu Ser Pro Gly Gly Leu Ser 65 70 75 80

Tyr Ser Pro Ser Gln Leu Leu Ala Ser Ser Phe Phe Ser Cys Gly Met 85 90 95

Ser Thr Ile Leu Gln Thr Trp Met Gly Ser Arg Leu Pro Leu Val Gln 100 105 110

Ala Pro Ser Leu Glu Phe Leu Ile Pro Ala Leu Val Leu Thr Ser Gln 115 120 125

Lys Leu Pro Arg Ala Ile Gln Thr Pro Gly Asn Ser Ser Leu Met Leu

His Leu Cys Arg Gly Pro Ser Cys His Gly Leu Gly His Trp Asn Thr 145 150 150 155 160

Ser Leu Gln Glu Val Ser Gly Ala Val Val Ser Gly Leu Leu Gln

Gly Met Met Gly Leu Leu Gly Ser Pro Gly His Val Phe Pro His Cys 180 185 190

Gly Pro Leu Val Leu Ala Pro Ser Leu Val Val Ala Gly Leu Ser Ala 195 200 205

His Arg Glu Val Ala Gln Phe Cys Phe Thr His Trp Gly Leu Ala Leu 210 215 220

Leu Tyr Val Ser Pro Glu Arg Arg Gly Met Val Pro Ser Gly Gly Val 225 230 235 240

Trp Gly Asp

<210> 82

<211> 1865

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (99)..(1508)

<223>

<400> 82

attttatttc aggaatccat caacatcctt tgcagctaca taggcaggaa aatctagaaa ttgtaattta tatagaattt taaaactctt caattaca atg gat aga ggg gag aaa

16U 200 PCT FINAL.ST25 Met Asp Arg Gly Glu Lys 1 5

cag Gln									164
att Ile									212
gca Ala 40									260
tgt Cys									308
ata Ile									356
ttt Phe									404
999 Gly									452
atc Ile 120									500
tgt Cys									548
ggt Gly									596
gtg Val									644
aga Arg									692
gaa Glu 200									740
ttt Phe								•	788
aag Lys								1	836
ctg Leu								1	884
ctg Leu									932
gct Ala 280									980
tct Ser								1	028

Page 44

16U 200 PCT FINAL.ST25 tcg aga cca ata tat ctt gca agc caa gag ggc cag ctg cct ttg cta Ser Arg Pro Ile Tyr Leu Ala Ser Gln Glu Gly Gln Leu Pro Leu Leu	1076
315 320 325	
ttt aat aca ctt aat agt cac tct tct cca ttt aca gct gtg cta cta Phe Asn Thr Leu Asn Ser His Ser Ser Pro Phe Thr Ala Val Leu Leu 330 335 340	1124
ctt gtc act ttg gga tcc ctt gca att atc tta aca agt cta att gat Leu Val Thr Leu Gly Ser Leu Ala Ile Ile Leu Thr Ser Leu Ile Asp 345 350 355	1172
ttg ata aac tat att ttt ttc acg ggt tca tta tgg tct ata tta tta Leu Ile Asn Tyr Ile Phe Phe Thr Gly Ser Leu Trp Ser Ile Leu Leu 360 365 370	1220
atg ata gga ata cta agg cgg aga tac cag gaa ccc aat cta tct ata Met Ile Gly Ile Leu Arg Arg Tyr Gln Glu Pro Asn Leu Ser Ile 375 380 385 390	1268
CCt tat aag gtg ttt ttg tca ttt cca tta gca aca ata gtc atc gac Pro Tyr Lys Val Phe Leu Ser Phe Pro Leu Ala Thr Ile Val Ile Asp 395 400 405	1316
gtg ggc ttg gtt gtg ata cca ttg gta aag tct cca aat gtg cat tat Val Gly Leu Val Val Ile Pro Leu Val Lys Ser Pro Asn Val His Tyr 410 415 420	1364
gtc tac gtg ctt ctg tta gtt ctc agc gga tta cta ttt tac ata cct Val Tyr Val Leu Leu Leu Val Leu Ser Gly Leu Leu Phe Tyr Ile Pro 425 430 435	1412
tta ata cat ttt aaa ata aga ttg gct tgg ttt gag aag atg act tgc Leu Ile His Phe Lys Ile Arg Leu Ala Trp Phe Glu Lys Met Thr Cys 440 445 450	1460
tat tta caa tta cta ttt aat att tgc ctc cct gat gtg tct gag gaa Tyr Leu Gln Leu Leu Phe Asn Ile Cys Leu Pro Asp Val Ser Glu Glu 455 460 465 470	1508
tagatgtcgg aagtgcaaac tcttaaaaaa ttggccttct aaaaaacata tatcagattc	1568
Caaatcaagg ttaaacatat gatagaacat tcatggtgaa attcctatgg taaatatttt	1628
tttctcaaat gaaataagta atgtatacaa aagtgcctaa gacagtacct ggcttcagag	1688
tcactaagaa attgctaaaa gctctgcttc gcatggtaaa aaacttaagt cctggtttgc	1748
gtagcttgat agagtgatta tacaacttcc attctctcac ttttttttc tgtatcccac	1808
cccttttcta ctgaatttgt ggggatccta taataaaagt gaatgactaa aaatttt	1865
<210> 83 <211> 470 <212> PRT <213> Homo sapiens	
<400> 83	
Met Asp Arg Gly Glu Lys Ile Gln Leu Lys Arg Val Phe Gly Tyr Trp 1 10 15	
Trp Gly Thr Ser Phe Leu Leu Ile Asn Ile Ile Gly Ala Gly Ile Phe 20 25 30	
Val Ser Pro Lys Gly Val Leu Ala Tyr Ser Cys Met Asn Val Gly Val 35 40 45	
Ser Leu Cys Val Trp Ala Gly Cys Ala Ile Leu Ala Met Thr Ser Thr 50 60	

Leu Cys Ser Ala Glu Ile Ser Ile Ser Phe Pro Cys Ser Gly Ala Gln 65 70 75 80 80

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Tyr Tyr Phe Leu Lys Arg Tyr Phe Gly Ser Thr Val Ala Phe Leu Asn 85 90 95

Leu Trp Thr Ser Leu Phe Leu Gly Ser Gly Val Val Ala Gly Gln Ala 100 105 110

Leu Leu Leu Ala Glu Tyr Ser Ile Gln Pro Phe Phe Pro Ser Cys Ser 115 120 125

Val Pro Lys Leu Pro Lys Lys Cys Leu Ala Leu Ala Met Leu Trp Ile 130 135 140

Val Gly Ile Leu Thr Ser Arg Gly Val Lys Glu Val Thr Trp Leu Gln 145 $\,$ 150 $\,$ 160

Ile Ala Ser Ser Val Leu Lys Val Ser Ile Leu Ser Phe Ile Ser Leu 165 170 175

Thr Gly Val Val Phe Leu Ile Arg Gly Lys Lys Glu Asn Val Glu Arg 180 185 190

Phe Gln Asn Ala Phe Asp Ala Glu Leu Pro Asp Ile Ser His Leu Ile 195 200 205

Gln Ala Ile Phe Gln Gly Tyr Phe Ala Tyr Ser Gly Gly Ala Cys Phe 210 215 220

Thr Leu Ile Ala Gly Glu Leu Lys Lys Pro Arg Thr Thr Ile Pro Lys 225 230 235 240

Cys Ile Phe Thr Ala Leu Pro Leu Val Thr Val Val Tyr Leu Leu Val 245 250 255

Asn Ile Ser Tyr Leu Thr Val Leu Thr Pro Arg Glu Ile Leu Ser Ser 260 265 270

Asp Ala Val Ala Ile Thr Trp Ala Asp Arg Ala Phe Pro Ser Leu Ala 275 280 285

Trp Ile Met Pro Phe Ala Ile Ser Thr Ser Leu Phe Ser Asn Leu Leu 290 295 300

Ile Ser Ile Phe Lys Ser Ser Arg Pro Ile Tyr Leu Ala Ser Gln Glu 305 310 315 320

Gly Gln Leu Pro Leu Leu Phe Asn Thr Leu Asn Ser His Ser Ser Pro 325 330 335

Phe Thr Ala Val Leu Leu Leu Val Thr Leu Gly Ser Leu Ala Ile Ile 340 345 350

Leu Thr Ser Leu Ile Asp Leu Ile Asn Tyr Ile Phe Phe Thr Gly Ser 355 360 365

Leu Trp Ser Ile Leu Leu Met Ile Gly Ile Leu Arg Arg Arg Tyr Gln 370 380

Glu Pro Asn Leu Ser Ile Pro Tyr Lys Val Phe Leu Ser Phe Pro Leu 385 390 395 400

Ala Thr Ile Val Ile Asp Val Gly Leu Val Val Ile Pro Leu Val Lys $405 \hspace{1.5cm} 410 \hspace{1.5cm} 415 \hspace{1.5cm}$

Ser Pro Asn Val His Tyr Val Tyr Val Leu Leu Leu Val Leu Ser Gly
420 425 430

Leu Leu Phe Tyr Ile Pro Leu Ile His Phe Lys Ile Arg Leu Ala Trp 435 440 445

Phe Glu Lys Met Thr Cys Tyr Leu Gln Leu Leu Phe Asn Ile Cys Leu 450 455 460

Pro Asp Val Ser Glu Glu 465 470

<210> 84

<211> 1046

<212> DNA <213> Homo sapiens

<220>

<221> CDS

<222> (319)..(852)

<223>

<400> 84

gaacacatct gaatteette tetgtggeat atgetttagg agaggageag acagetetta 60 gctagggtca gatttcaaat tctcatctct tggtqccaat accaccaca gattcttctt 120 tgaagtcaac ttttgagatc ttcactaagt acacgttggt gtctgaagat tcacacgagt 180 240 gcctctggta atcattttct tcagggaatc acagtctctc ctctcagcaa agcatccact gtactgaact ttgcttttgg aaacatcttc ttcctgagac ctcgttgaaa gaaactctct 300 ggtgtcatac tttccaat atg gag gtg aag aac ttt gca gtt tgg gat tat Met Glu Val Lys Asn Phe Ala Val Trp Asp Tyr l 5 $$ 351 gtt gta ttt gca gcc ctc ttt ttc att tcc tct gga att ggg gtg ttc Val Val Phe Ala Ala Leu Phe Phe Ile Ser Ser Gly Ile Gly Val Phe 15 20 25 399 ttt gcc att aag gag aga aaa aag gca act tcc cga gag ttc ctg gtt Phe Ala Ile Lys Glu Arg Lys Lys Ala Thr Ser Arg Glu Phe Leu Val 30 35 447 495

ggg gga agg caa atg agc ttt ggc cct gtc ggc ttg tct ctg aca gcc
Gly Gly Arg Gln Met Ser Phe Gly Pro Val Gly Leu Ser Leu Thr Ala
45 50 55

agc ttc atg tca gct gtc acg gtc ctg ggg acc cct tct gaa gtc tac

agc ttc atg tca gct gtc acg gtc ctg ggg acc cct tct gaa gtc tac 543 Ser Phe Met Ser Ala Val Thr Val Leu Gly Thr Pro Ser Glu Val Tyr 60 65 70 75

cgc ttt ggg gca tcc ttc cta gtc ttc ttc att gct tac cta ttt gtc 591 Arg Phe Gly Ala Ser Phe Leu Val Phe Phe Ile Ala Tyr Leu Phe Val 80 85 90

atc ctc tta aca tca gag ctc ttt ctc cct gtg ttc tac aga tct ggt

1le Leu Leu Thr Ser Glu Leu Phe Leu Pro Val Phe Tyr Arg Ser Gly
95 100 105

atc acc agc act tat gag tac tta caa cta cga ttc aac aaa cca gtt

Ile Thr Ser Thr Tyr Glu Tyr Leu Gln Leu Arg Phe Asn Lys Pro Val

110 115 120

cgc tat gct gcc acg gtc atc tac att gta cag acg att ctc tac aca
Arg Tyr Ala Ala Thr Val Ile Tyr Ile Val Gln Thr Ile Leu Tyr Thr
125
130
135

16U 200 PCT FINAL.ST25 gga gtg gtg gtg tat gct cct gcc ctg gca ctc aat caa gtg act ggg Gly Val Val Val Tyr Ala Pro Ala Leu Ala Leu Asn Gln Val Thr Gly 140 145 150 155	783
ttt gat ctc tgg ggc tct gtg ttt gca aca gga att gtt tgc aca ttc Phe Asp Leu Trp Gly Ser Val Phe Ala Thr Gly Ile Val Cys Thr Phe 160 165 170	831
tac tgt acc ctg gta tgt atc tagctgtgaa gaagtattta acactacctc Tyr Cys Thr Leu Val Cys Ile 175	882
ctaatatggg ataagggcaa atctccagca ataggcatct aattatagca gaattcgtta	942
ttccaaaatt aagcagaagt atgtcggctt atctgtcaca gtttcctgag gaaggtgctg	1002
ttgtttaaca ttcttttcat taccaacctt taggagaatt taat	1046
<210> 85 <211> 178 <212> PRT <213> Homo sapiens	
<400> 85	
Met Glu Val Lys Asn Phe Ala Val Trp Asp Tyr Val Val Phe Ala Ala 1 10 15	
Leu Phe Phe Ile Ser Ser Gly Ile Gly Val Phe Phe Ala Ile Lys Glu 20 25 30	
Arg Lys Lys Ala Thr Ser Arg Glu Phe Leu Val Gly Gly Arg Gln Met 35 40 45	
Ser Phe Gly Pro Val Gly Leu Ser Leu Thr Ala Ser Phe Met Ser Ala 50 60	
Val Thr Val Leu Gly Thr Pro Ser Glu Val Tyr Arg Phe Gly Ala Ser 65 70 75 80	
Phe Leu Val Phe Phe Ile Ala Tyr Leu Phe Val Ile Leu Leu Thr Ser 85 90 95	
Glu Leu Phe Leu Pro Val Phe Tyr Arg Ser Gly Ile Thr Ser Thr Tyr 100 105 110	
Glu Tyr Leu Gln Leu Arg Phe Asn Lys Pro Val Arg Tyr Ala Ala Thr 115 120 125	
Val Ile Tyr Ile Val Gln Thr Ile Leu Tyr Thr Gly Val Val Val Tyr 130 135 140	
Ala Pro Ala Leu Ala Leu Asn Gln Val Thr Gly Phe Asp Leu Trp Gly 145 150 155 160	
Ser Val Phe Ala Thr Gly Ile Val Cys Thr Phe Tyr Cys Thr Leu Val 165 170 175	
Cys Ile ,	÷
<210> 86 <211> 4751 <212> DNA <213> Homo sapiens	,

<220 <221	> (DS								160	200	PCT	FINA	AL.ST	25	
<222 <223		(382)) (:	1929)											
<400 cccg		36 gag (ctgg	eggg	ge te	gaate	gagc	c ggt	tgca	tttc	gaag	ggcc	gag (cacto	gggat	t 60
ttcc	gcgo	cta (egeti	tece	tc to	ccg	gcgti	t gc	cago	tagg	ccc	ccgg	ccc (cagco	ctcgc	c 120
ggcg	ccto	ccg (ccca	gtcc	gc to	ccc	gccc	c acc	cgaa	gcgc	ggat	togo	gca (gcct	gggc	c 180
cggg	aagg	, 99 (ccact	tgcg	ca go	ggac	gcgg	c tc	ggcg	ggtg	cgc	ccg	ggg (gcat	gtccg	c 240
gcgc	taco	gc (cagg	gctg	ca g	ggt	cccg	g cga	aggc	cctg	gcaa	acca	cca 1	ttcta	acttt	t 300
tgtg	tcta	atg i	agtti	tgac	ta co	cta	agga	cto	caca	tggc	gagt	taacı	cca 1	tggg	ccagg	t 360
agcg	ttct	at (gcca	acct	tg a					agt Ser 5						411
tct Ser																459
aaa Lys																507
cca Pro																555
acc Thr																603
gag Glu 75																651
cat His																699
gtt Val																747
ttg Leu																795
aag Lys		_				-										843
ctt Leu 155																891
cgg Arg																939
acc Thr																987
tgg Trp																1035
gta Val																1083
act	cca	gag	cag	aac	cat	gtg	tgt	aga	act	cag	aaa	gaa	gac	att	aag	1131

WO 03/089583 PCT/US03/11497

Thr 235	Pro	Glu	Gln	Asn	His 240	Val	Суѕ	Arg	Thr					L.SI Ile		
	gtg Val															1179
	tgt Cys															1227
	gtt Val															1275
	ctc Leu 300															1323
	cag Gln															1371
	Gl y															1419
	tac Tyr															1467
	tat Tyr															1515
	tct Ser 380															1563
	gta Val															1611
	ggt Gly															1659
	atc Ile															1707
	ctc Leu															1755
	gct Ala 460															1803
	aaa Lys															1851
	tat Tyr															1899
	aca Thr									tgad	ccaa	agg (cctt	gage	cc	1949
caga	aatc	ttc a	aggt '	ttga	ga ga	aggt	gggg	cac	ccaga	attc	ttc	atgt	ttc 1	tgaaa	cttt	2009
tat	tttg	gca (gaag	gatt	gc c	tcca	aagga	a aat	tati	tatt	att	gttt	tgt 1	taaca	atatta	2069
ata	ttta	taa (ggga	aaac	ag c	acata	aataa	a gga	aaago	etgg	acta	agcc	cag a	agcci	ttctca	2129
ttt	ggga	ttt (gtgc	tcat	aa c	tgaad	ctcg	tato	ettt	tggt	caa	tggg	cat a	agcto	ctgtaa	2189

			1 6 U	200 PCT FIR	NAL.ST25	
gaaatgtaag	gacacagctg	atataattag	ctgtaattag	ggataatttc	aaagcataac	2249
caaagcagat	gacactgggc	agcagctttg	ttccagtctc	aggcccttca	tgttccctcc	2309
tcagaaagaa	aatggaaaca	ttaacgtgta	gctttgctta	ccttgttctg	gttagagaag	2369
ggaggtcagc	ttgggtgtgg	tggtgaagag	tgaagatgcc	atactttttc	atggtggagt	2429
ttctcattag	ggttttactt	gggattgtta	aagaatactt	gagattcttc	aaaaagtggt	2489
gattaatata	gaaagaaact	cttattttt	ttttctctta	gtcttccagc	cagecettge	2549
ctctgcccaa	gggtagacac	cactatgaga	atccaaataa	tcatggaatg	ccatggttgg	2609
aatagatctt	aaagggcatc	tggtaagatc	catttgaaat	tgtccactgg	aaaccgaaag	2669
ctcttttcct	aagactgggt	tccaggctct	cacatttgtt	accatcacat	ataatactta	2729
ctctaaattt	agcagaacac	acttagtcac	aaggacaacc	tctcaatctt	acctgaaatg	2789
tcaacaacac	caaaacttcc	cgtcttttac	cttcagagaa	gaagctctta	cttagactgc	2849
agacgcattc	ctgttaggtt	ggaaaaatgt	tggcagtatt	ccaattgggc	aggaactgaa	2909
ttcttgaatc	agcaggtctc	tggtgagagt	tttctttgca	gatcagacat	ttagttttat	2969
cattacccaa	aagaggattg	gagggagtca	gttgtctgaa	aaatattatc	ctagagatat	3029
tctaaaggtg	agattccttt	ctccctgtgt	taattcttgt	tccactatcc	actgctcttc	3089
atctctttat	agataataat	tagaaatcta	ctcattggat	tataagttta	ttcattctca	3149
aatactccac	ttttctatgg	tttgggataa	tttctgagtc	ttcagattga	agagggaagg	3209
catggaggga	agaaaaagtc	cagatecece	agcttgtttc	caaccatttt	aagtccaaag	3269
aattataatc	ctgaatctca	cagtgtgtca	cacctgtaat	aggagtaaat	tatgcaatca	3329
attttaatta	ccaggagttt	aaaatccaaa	tgtcaaggaa	ctgttttgac	cctgaaggct	3389
atttaatcca	ctgtccccta	caaggcctca	caagtgctgg	gggaaaaaaa	acagcaatga	3449
ggatgatcct	gagttaatgt	gtatgctccg	caagagagct	tgcctatacc	ttgattattt	3509
cataaaatca	catgttaata	cattgctttc	agaatgaaat	actgacttga	tctgatagga	3569
gaaaatggta	atatttcata	gttgttttcc	aaagacaaat	ttaaatgttg	tctgttatct	3629
ccttacttag	tttaagaatt	tagttttgaa	ccccattgac	tttgtcattt	gcaattttaa	3689
aaatatttgg	gactgggcat	ggtcgctcac	gcctgtaatc	ccagcacttt	gggaggctga	3749
ggcgggtgga	tcatgaggtc	aggagatcaa	gaccatcctg	gctaacatcg	tgaaactccg	3809
tctctactaa	aaatgcaaaa	gattagccag	gcgtggtggc	gggcgcctgt	agtcccagct	3869
actcatgagg	ctgaggccgg	acaatcgctt	gaacccagga	ggtggaggtt	gcagtgagcc	3929
aaaatcatgc	cactgcactc	caccctgggc	gacagagcaa	gactccatct	Саааааааа	3989
aaattggaag	gtatctgtaa	aatgtcaaag	ttaagatgaa	gttatatctg	tttggaatag	4049
cactttgccc	taaatatcat	ttcttgaatt	ttcaagccta	aagatgttta	aaaatatgaa	4109
tagttacaaa	tattcttata	catattttt	atcatgatca	caacaaaatt	ttgtttatgt	4169
ggttctgcaa	tataatttct	gtgaagtatt	acaagtattt	atgaaaaata	agcatagtga	4229
tcagaaattt	taaagatttt	gtataaaaac	atttgggaga	tttgacttta	tacatgcata	4289
gatttgcatt	ttactttccc	ttttgaggca	gcatttttag	aaaatcagta	agaaaaatgt	4349
	gtctactatt		_		, ,	4409
	tttatttcaa				•••	4469
agaacaaaat	tatagtggga	taattttaca	tcttaaatat	ttctttctac	tactgtaagc	4529
	aattatctga				•	4589
			-			

tgtatgattg	ggctttttgt	tcagattgta	atttcattaa	tagatgaaat	atttatgcta	4649
atattttctt	atttcaaaag	caaaataaaa	tgaatttatt	gtcctgtgta	aaaaaaaaa	4709
aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	aaaaaaaaa	aa		4751

<210> 87 <211> 516

<212> PRT

<213> Homo sapiens

<400> 8

Met Pro Ser Gly Ser His Trp Thr Ala Asn Ser Ser Lys Ile Ile Thr 1 5 10 15

Trp Leu Leu Glu Gln Pro Gly Lys Glu Glu Lys Arg Lys Thr Met Ala 20 25 30

Lys Val Asn Arg Ala Arg Ser Thr Ser Pro Pro Asp Gly Gly Trp Gly 40

Trp Met Ile Val Ala Gly Cys Phe Leu Val Thr Ile Cys Thr Arg Ala 50 55 60

Val Thr Arg Cys Ile Ser Ile Phe Phe Val Glu Phe Gln Thr Tyr Phe 65 70 75 80

Thr Gln Asp Tyr Ala Gln Thr Ala Trp Ile His Ser Ile Val Asp Cys $85 \hspace{1cm} 90 \hspace{1cm} 95$

Val Thr Met Leu Cys Ala Pro Leu Gly Ser Val Val Ser Asn His Leu 100 105 110

Ser Cys Gln Val Gly Ile Met Leu Gly Gly Leu Leu Ala Ser Thr Gly 115 120 125

Leu Ile Leu Ser Ser Phe Ala Thr Ser Leu Lys His Leu Tyr Leu Thr 130 135 140

Leu Gly Val Leu Thr Gly Leu Gly Phe Ala Leu Cys Tyr Ser Pro Ala 145 150 155 160

Ile Ala Met Val Gly Lys Tyr Phe Ser Arg Arg Lys Ala Leu Ala Tyr 165 170 175

Gly Ile Ala Met Ser Gly Ser Gly Ile Gly Thr Phe Ile Leu Ala Pro 180 190

Val Val Gln Leu Leu Ile Glu Gln Phe Ser Trp Arg Gly Ala Leu Leu 195 200 205

Ile Leu Gly Gly Phe Val Leu Asn Leu Cys Val Cys Gly Ala Leu Met 210 220

Arg Pro Ile Thr Leu Lys Glu Asp His Thr Thr Pro Glu Gln Asn His 225 230 235 240

Val Cys Arg Thr Gln Lys Glu Asp Ile Lys Arg Val Ser Pro Tyr Ser 245 250 255 Ser Leu Thr Lys Glu Trp Ala Gln Thr Cys Leu Cys Cys Leu Gln 260 265 270

Gln Glu Tyr Ser Phe Leu Leu Met Ser Asp Phe Val Val Leu Ala Val 275 280 285

Ser Val Leu Phe Met Ala Tyr Gly Cys Ser Pro Leu Phe Val Tyr Leu 290 295 300

Val Pro Tyr Ala Leu Ser Val Gly Val Ser His Gln Gln Ala Ala Phe 305 310315320

Leu Met Ser Ile Leu Gly Val Ile Asp Ile Ile Gly Asp Ile Thr Phe $325 \hspace{1.5cm} 330 \hspace{1.5cm} 335$

Gly Trp Leu Thr Asp Arg Arg Cys Leu Lys Asn Tyr Gln Tyr Val Cys 340 345 350

Tyr Leu Phe Ala Val Gly Met Asp Gly Leu Cys Tyr Leu Cys Leu Pro 355 360 365

Met Leu Gln Ser Leu Pro Leu Leu Val Pro Phe Ser Cys Thr Phe Gly 370 375 380

Tyr Phe Asp Gly Ala Tyr Val Thr Leu Ile Pro Val Val Thr Thr Glu 385 390 395 400

Ile Val Gly Thr Thr Ser Leu Ser Ser Ala Leu Gly Val Val Tyr Phe 405 410 415

Leu His Ala Val Pro Tyr Leu Val Ser Pro Pro Ile Ala Gly Arg Leu 420 425 430

Val Asp Thr Thr Gly Ser Tyr Thr Ala Ala Phe Leu Leu Cys Gly Phe 435 440 445

Ser Met Ile Phe Ser Ser Val Leu Leu Gly Phe Ala Arg Leu Ile Lys 450 455 460

Arg Met Arg Lys Thr Gln Leu Gln Phe Ile Ala Lys Glu Ser Asp Pro 465 470 470 480

Lys Leu Gln Leu Trp Thr Asn Gly Ser Val Ala Tyr Ser Val Ala Arg 485 490 495

Glu Leu Asp Gln Lys His Gly Glu Pro Val Ala Thr Ala Val Pro Gly 500 510

Tyr Ser Leu Thr

<210> 88

<211> 2150

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (63)..(1760)

<223>

<400> 88

16U 200 PCT FINAL.ST25 gctggacaaa gctggccgtg caggcgctca ggcgtgcagg gtagccagtg ccccggccag 6													60		
1				Leu					gcc Ala				Gly		107
									aga Arg 25						155
									ctg Leu						203
									gtg Val						251
									gcg Ala						299
									tct Ser						347
					Gly				aag Lys 105	Lys				Ile	395
									ctc						443
								Ile	ctg Leu						491
							Gln		tat Tyr			Ile			539
									ctg Leu		Ala				587
					Trp				ctc Leu 185					Asn	635
									ctg Leu						683
								Asn	atc Ile						731
									ctg Leu			Leu			779
									ctg Leu		Asp				827
									atg Met 265					Trp	875
									ggg Gly						923
								Glu	gtg Val						971
att	ccc	ttg	ggg	ctc	agc	atc	ggg	gtc	tgt	gtc		gtg	atg	gct	1019

Ile	Pro 3 0 5	Leu	Gly	Leu	Ser	Ile 310	Gly	Val	Cys	16U Val	200 Arg 315	PCT Val	FINA Gly	L.ST Met	25 Ala	
ctg Leu 320	ggg Gly	gct Ala	gcg Ala	gat Asp	act Thr 325	gtg Val	cag Gln	gcc Ala	aag Lys	cgc Arg 330	tcg Ser	gcc Ala	gtc Val	tcg Ser	ggc Gly 335	1067
gtg Val	ctc Leu	agc Ser	ata Ile	gtt Val 340	ggc Gly	att Ile	tcc Ser	ctg Leu	gtc Val 345	ctg Leu	ggc Gly	acc Thr	ctg Leu	ata Ile 350	agc Ser	1115
atc Ile	ctg Leu	aaa Lys	aat Asn 355	cag Gln	ctg Leu	ggg Gly	cat His	att Ile 360	ttt Phe	acc Thr	aat Asn	gat Asp	gaa Glu 365	gat Asp	gtc Val	1163
att Ile	gcc Ala	ctg Leu 370	Val	agc Ser	cag Gln	gtc Val	ttg Leu 375	ccg Pro	gtt Val	tat Tyr	agt Ser	gtc Val 380	ttt Phe	cac His	gtg Val	1211
ttt Phe	gag Glu 385	gcc Ala	atc Ile	tgt Cys	tg t Cys	gtc Val 390	tat Tyr	ggc Gly	gga Gly	gtt Val	ctg Leu 395	Arg	gga Gly	act Thr	ggg Gly	1259
aag Lys 400	Gln	gcc Ala	ttt Phe	ggt Gly	gcc Ala 405	gct Ala	gtg Val	aat Asn	gcc Ala	atc Ile 410	aca Thr	tat Tyr	tac Tyr	atc Ile	atc Ile 415	1307
ggc Gly	cta Leu	cca Pro	ctg Leu	ggc Gly 420	Ile	ctt Leu	ctg Leu	acc Thr	ttt Phe 425	gtg Val	gto Val	aga Arg	atg Met	aga Arg 430	116	1355
atg Met	ggc Gly	cto Lev	tgg Trp 435	Leu	ggc Gly	atg Met	ctg Leu	gcc Ala 440	tgt Cys	gtc Val	tto Phe	ctg Leu	gca Ala 445	act Thr	gct Ala	1403
gcc Ala	ttt Phe	gtt Val 450	L Ala	tat Tyr	act Thr	gcc Ala	cgg Arg 455	Leu	gac Asp	tgg Trp	aaq Lys	g ctt Leu 460	ı wra	gca Ala	gag Glu	1451
gaç Glu	gct Ala 465	Lys	g aaa S Lys	cat His	tca Ser	ggc G1y 470	Arc	cag Glr	g cag Gln	cag Gln	Gl:	n Arg	gca J Ala	gag Glu	agc Ser	1499
act Thi 480	: Ala	a acc	c aga r Arg	cct Pro	ggg Gly 485	Pro	gaç Gli	g aaa i Lys	a gca s Ala	gtc Val 490	. ье	a tct u Ser	tca Ser	gto Val	gct Ala 495	1547
aca Th:	a ggo c Gly	ag y Se	t tco r Sei	c cct Pro	o G13	att / Ile	aco Thi	tto Lei	aca Thi	Thi	ta Ty	t tca r Sei	a agg	tct Ser 510	gag Glu	1595
tg:	c cad s His	c gt s Va	g gad 1 Asp 515	o Phe	c tto e Phe	ago Aro	act Thi	520	o GI	g gaç ı Glu	g gc ı Al	c cad a His	c gcc s Ala 525	, re	tca Ser	1643
gc Al	t cc	t ac o Th 53	r Se	c ag	acta g Lei	tca Sei	gto Val	l Ly:	a caq s Gli	g cto	g gt u Va	c ato 1 110 540	5 WIG	cgi g Ar	t ggg g Gly	1691
gc Al	t gc a Al 54	a Le	g gg u Gl	g gc	g gco a Alo	g toa ser 550	r Al	c aca	a cto	g ato	g gt t Va 55	1 61	g cto y Lev	ace Th	g gtc r Val	1739
ag Ar 56	g 11	c ct e Le	a gc	c ac a Th	c age r Are 56	g Hi	c ta s	gcaa	agaa	gct	tgga	aat	agaa	agcc	ag	1790
ga	gtgg	ctgt	ccc	cagt	atg	caaa	caca	сс а	cggt	ctgc	c ct	gcaa	aaac	acc	aatgggg	1850
															ttccttg	
															tggagtt	
															agagtcc	
ag	ggat	tgto	act	atta	tta	ataa	tgta	aa t	ggct	tcaa	a to	ggac	actg	cag	ataaaat	2090
са	caaa	aaco	act	gtta	tat	taaa	gatt	ac a	catt	tcct	g gg	gaaaa	aaaa	aaa	aaaaaa	2150

<210>	89
<211>	566
Z2125	TGG

<213> Homo sapiens

<400> 89

Met Asp Ser Leu Gln Asp Thr Val Ala Leu Asp His Gly Gly Cys Cys 1 $$ 5 $$ 10 $$ 15

Pro Ala Leu Ser Arg Leu Val Pro Arg Gly Phe Gly Thr Glu Met Trp $20 \hspace{1cm} 25 \hspace{1cm} 30$

Thr Leu Phe Ala Leu Ser Gly Pro Leu Phe Leu Phe Gln Val Leu Thr 35 40 45

Phe Met Ile Tyr Ile Val Ser Thr Val Phe Cys Gly His Leu Gly Lys 50 60

Val Glu Leu Ala Ser Val Thr Leu Ala Val Ala Phe Val Asn Val Cys 65 70 75 80

Gly Val Ser Val Gly Val Gly Leu Ser Ser Ala Cys Asp Thr Leu Met 85 90 95

Ser Gln Ser Phe Gly Ser Pro Asn Lys Lys His Val Gly Val Ile Leu 100 105 110

Gln Arg Gly Ala Leu Val Leu Leu Cys Cys Leu Pro Cys Trp Ala 115 . 120 . 125

Leu Phe Leu Asn Thr Gln His Ile Leu Leu Leu Phe Arg Gln Asp Pro 130 135 140

Asp Val Ser Arg Leu Thr Gln Asp Tyr Val Met Ile Phe Ile Pro Gly 145 150 150 160

Leu Pro Val Ile Phe Leu Tyr Asn Leu Leu Ala Lys Tyr Leu Gln Asn 165 170 175

Gln Lys Ile Thr Trp Pro Gln Val Leu Ser Gly Val Val Gly Asn Cys 180 185 190

Val Asn Gly Val Ala Asn Tyr Ala Leu Val Ser Val Leu Asn Leu Gly 195 200 205

Val Phe Leu Leu Leu Tyr Ile Val Leu Lys Lys Leu His Leu Glu Thr 225 230 235 240

Trp Ala Gly Trp Ser Ser Gln Cys Leu Gln Asp Trp Gly Pro Phe Phe 245 250255

Ser Leu Ala Val Pro Ser Met Leu Met Ile Cys Val Glu Trp Trp Ala 260 265 270

Tyr Glu Ile Gly Ser Phe Leu Met Gly Leu Leu Ser Val Val Asp Leu 275 280 285

Ser Ala Gln Ala Val Ile Tyr Glu Val Ala Thr Val Thr Tyr Met Ile 290 295 300

Pro Leu Gly Leu Ser Ile Gly Val Cys Val Arg Val Gly Met Ala Leu 305 310 315 320

Gly Ala Ala Asp Thr Val Gln Ala Lys Arg Ser Ala Val Ser Gly Val $325 \hspace{1cm} 330 \hspace{1cm} 335$

Leu Ser Ile Val Gly Ile Ser Leu Val Leu Gly Thr Leu Ile Ser Ile $340 \hspace{1.5cm} 345 \hspace{1.5cm} 350 \hspace{1.5cm}$

Ala Leu Val Ser Gln Val Leu Pro Val Tyr Ser Val Phe His Val Phe 370 375 380

Glu Ala Ile Cys Cys Val Tyr Gly Gly Val Leu Arg Gly Thr Gly Lys 385 390 395 400

Gln Ala Phe Gly Ala Ala Val Asn Ala Ile Thr Tyr Tyr Ile Ile Gly 405 410 415

Leu Pro Leu Gly Ile Leu Leu Thr Phe Val Val Arg Met Arg Ile Met 420 . 430

Gly Leu Trp Leu Gly Met Leu Ala Cys Val Phe Leu Ala Thr Ala Ala 435 440 445

Ala Lys Lys His Ser Gly Arg Gln Gln Gln Gln Arg Ala Glu Ser Thr 465 470 475 480

Ala Thr Arg Pro Gly Pro Glu Lys Ala Val Leu Ser Ser Val Ala Thr 485 490 495

Gly Ser Ser Pro Gly Ile Thr Leu Thr Thr Tyr Ser Arg Ser Glu Cys 500 505 510

His Val Asp Phe Phe Arg Thr Pro Glu Glu Ala His Ala Leu Ser Ala 515 520 525

Pro Thr Ser Arg Leu Ser Val Lys Gln Leu Val Ile Arg Arg Gly Ala 530 535 540

Ala Leu Gly Ala Ala Ser Ala Thr Leu Met Val Gly Leu Thr Val Arg 545 550 560

Ile Leu Ala Thr Arg His 565

<210> 90

<211> 3067

<212> DNA

<213> Homo sapiens

<220>

<22 <22	_	CDS (36)	(2	984)						160	200	PCT	FIN	AL.S	Т25	
<22																
<40 gga		90 act	ggtt	ctga	ga t	tctg	tgca	a gc						ctg Leu 5		53
ggc Gly	cag Gln	gaa Glu	999 Gly 10	ttt Phe	gaa Glu	gcc Ala	tcc Ser	agt Ser 15	gct Ala	cct Pro	aga Arg	aat Asn	att Ile 20	cct Pro	tca Ser	101
															gat Asp	149
ggc Gly	ccc Pro 40	tca Ser	gac Asp	aca Thr	gag Glu	agc Ser 45	aag Lys	gaa Glu	ctg Leu	gga Gly	gta Val 50	ccc Pro	aaa Lys	gac Asp	cct Pro	197
ctg Leu 55	ctc Leu	ttc Phe	att Ile	cag Gln	ctg Leu 60	aat Asn	gag Glu	ctg Leu	ctg Leu	ggc Gly 65	tgg Trp	ccc Pro	cag Gln	gcg Ala	ctg Leu 70	245
			gag Glu													293
gga Gly	gaa Glu	atg Met	ccc Pro 90	tca Ser	ata Ile	aca Thr	ctg Leu	tct Ser 95	acc Thr	cac His	ctt Leu	cat His	cac His 100	Arg	tgg Trp	341
gta Val	ctg Leu	ttt Phe 105	gag Glu	gag Glu	aag Lys	ttg Leu	gag Glu 110	gtg Val	gct Ala	gca Ala	ggc Gly	cgg Arg 115	tgg Trp	agt Ser	gcc Ala	389
		Val	ccc Pro													437
ctg Leu 135	ctg Leu	gcc Ala	gag Glu	ggc Gly	ctt Leu 140	gta Val	ctg Leu	ctg Leu	gac Asp	tgc Cys 145	cca Pro	gct Ala	cag Gln	agc Ser	ctc Leu 150	485
ctg Leu	gag Glu	ctc Leu	gtg Val	gag Glu 155	cag Gln	gtg Val	acc Thr	agg Arg	gtg Val 160	gag Glu	tcg Ser	ctg Leu	agc Ser	cca Pro 165	gag Glu	533
ctg Leu	aga Arg	ggg Gly	cag Gln 170	ttg Leu	cag Gln	gcc Ala	ttg Leu	ctg Leu 175	ctg Leu	cag Gln	aga Arg	ccc Pro	cag Gln 180	cat His	tac Tyr	581
aac Asn	cag Gln	acc Thr 185	aca Thr	ggc Gly	acc Thr	agg Arg	ccc Pro 190	tgc Cys	tgg Trp	ggc Gly	tct Ser	act Thr 195	cat His	cca Pro	aga Arg	629
aag Lys	gct Ala 200	tct Ser	gac Asp	aat Asn	gag Glu	gaa Glu 205	gcc Ala	ccc Pro	ctg Leu	agg Arg	gaa Glu 210	cag Gln	tgt Cys	cag Gln	aac Asn	677
ccc Pro 215	ctg Leu	aga Arg	cag Gln	aag Lys	cta Leu 220	cct Pro	cca Pro	gga Gly	gct Ala	gag Glu 225	gca Ala	G1 y ggg	act Thr	gtg Val	ctg Leu 230	725
			ctg Leu													773
ctg Leu	cgg Arg	aac Asn	cct Pro 250	gtg Val	gta Val	ctg Leu	ggg Gly	tcc Ser 255	ctt Leu	act Thr	gag Glu	gtg Val	tcc Ser 260	ctc Leu	cca Pro	821
agc Ser	agg Arg	ttt Phe 265	ttc Phe	tgc Cys	ctt Leu	ctc Leu	ctg Leu 270	ggc Gly	ccc Pro	tgt Cys	atg Met	ctg Leu 275	gga Gly	aag Lys	ggc Gly	869
tac Tyr	cat His	gag Glu	atg Met	gga Gly	cgg Arg	gca Ala	gca Ala	gct Ala	gtc Val	ctc Leu	ctc Leu	agt Ser	gac Asp	ccg Pro	caa Gln	917

	280					285				16U	200 290	PCT	FINA	AL.SI	r25	
ttc		taa	tca	att	cat		acc	age	aac	ctt		asc	ctt	ctg	aca.	965
														Leu		303
														ggt Gly 325		1013
														tct Ser		1061
														gcc Ala		1109
														cac His		1157
														ctt Leu		1205
														ttg Leu 405		1253
														ctg Leu		1301
														gcc Ala		1349
														gtg Val		1397
														ctg Leu		1445
														ttc Phe 485		1493
aga Arg	gat Asp	tac Tyr	agc Ser 490	ctg Leu	gac Asp	tac Tyr	ctg Leu	ccc Pro 495	ttc Phe	cgc Arg	cta Leu	tgg Trp	gtg Val 500	ggc Gly	atc Ile	1541
														agt Ser		1589
														gcc Ala		1637
														aac Asn		1685
														ggg Gly 565		1733
														ata Ile		1781
														ggc Gly		1829
atc	aat	gca	tcc	ttg	ctg	ccg	сса	cct	gag	tgc		cgg		gga	ggc	1877

Ile	Asn 600	Ala	Ser	Leu	Leu'	Pro 605	Pro	Pro	Glu				AL.ST Gly		
					ggc G1 y 620										1925
					ctt Leu					-	-	-		-	1973
					cgc Arg										2021
					gtc Val										2069
					gcc Ala										2117
					999 Gly 700										2165
					agt Ser										2213
					atg Met										2261
					ctg Leu										2309
					ctg Leu										2357
					act Thr 780										2405
					gcc Ala										2453
					agg Arg										2501
					ctg Leu										2549
					ttc Phe										2597
					agg Arg 860										2645
					ctc Leu										2693
					cag Gln										2741
					gcc Ala										2789

Page 60

											200						
		Val			gcc Ala					ttc	tca	cca Pro	cag		ctc	2	837
	Trp				ctg Leu 940											2	885
					gaa Glu											2	933
				Tyr	cag Gln											2	981
aat Asn		ctgg	agt	agga	gtct	gg ga	agtg	gaga	c cc	cagg	aaac	agc	atga	ggt		3	034
gag	ggtg	tga	ggga	agtg	ct c	ctgai	tgtt	g age	g							3	067
<21 <21 <21 <21	1 > 2 >	91 983 PRT Homo	sap	iens													
< 40	0>	91															
Met 1	Glu	Met	Lys	Leu 5	Pro	Gly	Gln	Glu	Gly 10	Phe	Glu	Ala	Ser	Ser 15	Ala		
Pro	Arg	Asn	11e 20	Pro	Ser	Gly	Glu	Leu 25	Asp	Ser	Asn	Pro	Asp 30	Pro	Gly		
Thr	Gly	Pro 35	Ser	Pro	Asp	Gly	Pro 40	Ser	Asp	Thr	Glu	Ser 45	Lys	Glu	Leu		
Gly	Val 50	Pro	Lys	Asp	Pro	Leu 55	Leu	Phe	Ile	Gln	Leu 60	Asn	Glu	Leu	Leu		
Gly 65	Trp	Pro	Gln	Ala	Leu 70	Glu	Trp	Arg	Glu	Thr 75	Gly	Ser	Ser	Ser	Ala 80		
Ser	Leu	Leu	Leu	Asp 85	Met	Gly	Glu	Met	Pro 90	Ser	Ile	Thr	Leu	Ser 95	Thr		
His	Leu	His	His 100	Arg	Trp	Val	Leu	Phe 105	Glu	Glu	Lys	Leu	Glu 110	Val	Ala		
Ala	Gly	Arg 115	Trp	Ser	Ala	Pro	His 120	Val	Pro	Thr	Leu	Ala 125	Leu	Pro	Ser		
Leu	Gln 130	Lys	Leu	Arg	Ser	Leu 135	Leu	Ala	Glu	Gly	Leu 140	Val	Leu	Leu	Asp		
Cys 145	Pro	Ala	Gln	Ser	Leu 150	Leu	Glu	Leu	Val	Glu 155	Gln	Val	Thr	Arg	Val 160		
Glu	Ser	Leu	Ser	Pro 165	Glu	Leu	Arg	Gly	Gln 170	Leu	Gln	Ala	Leu	175	Leu :		
Gln	Arg	Pro	Gln 180	His	Tyr	Asn	Gln	Thr 185	Thr	Gly	Thr	Arg	Pro 190				
Gly	Ser	Thr	His	Pro	Arg	Lys	Ala	Ser	Asp	Asn		Glu Page		Pro	Leu		

16U 200 PCT FINAL.ST25 95 200 205

Arg Glu Gln Cys Gln Asn Pro Leu Arg Gln Lys Leu Pro Pro Gly Ala 210 215 220

Glu Ala Gly Thr Val Leu Ala Gly Glu Leu Gly Phe Leu Ala Gln Pro 225 230 235 240

Leu Gly Ala Phe Val Arg Leu Arg Asn Pro Val Val Leu Gly Ser Leu 245 250 255

Thr Glu Val Ser Leu Pro Ser Arg Phe Phe Cys Leu Leu Gly Pro 260 265 270

Cys Met Leu Gly Lys Gly Tyr His Glu Met Gly Arg Ala Ala Val 275 280 285

Leu Leu Ser Asp Pro Gln Phe Gln Trp Ser Val Arg Arg Ala Ser Asn 290 295 300

Leu His Asp Leu Leu Ala Ala Leu Asp Ala Phe Leu Glu Glu Val Thr 305 310 315 320

Val Leu Pro Pro Gly Arg Trp Asp Pro Thr Ala Arg Ile Pro Pro Pro 325 330 335

Ile Arg Gly Pro Ala Val Pro Arg Leu Thr Ser Ala Glu Asp Arg His 355 360 365

Arg His Gly Pro His Ala His Ser Pro Glu Leu Gln Arg Thr Gly Arg 370 375 380

Leu Phe Gly Gly Leu Ile Gln Asp Val Arg Arg Lys Val Pro Trp Tyr 385 390 395 400

Pro Ser Asp Phe Leu Asp Ala Leu His Leu Gln Cys Phe Ser Ala Val405 $$ 410 $$ $$ $$ $$

Leu Tyr Ile Tyr Leu Ala Thr Val Thr Asn Ala Ile Thr Phe Gly Gly 420 425 430

Leu Leu Gly Asp Ala Thr Asp Gly Ala Gln Gly Val Leu Glu Ser Phe 435 440 445

Leu Gly Thr Ala Val Ala Gly Ala Ala Phe Cys Leu Met Ala Gly Gln 450 460

Pro Leu Thr Ile Leu Ser Ser Thr Gly Pro Val Leu Val Phe Glu Arg 465 470 475 480

Leu Leu Phe Ser Phe Ser Arg Asp Tyr Ser Leu Asp Tyr Leu Pro Phe 485 490 495 ...

Arg Leu Trp Val Gly Ile Trp Val Ala Thr Phe Cys Leu Val Leu Val
500 510

Ala	Thr	Glu 515	Ala	Ser	Val	Leu.	Val 520	Arg	Туг					Thr	
Glu	Gly 530	Phe	Cys	Ala	Leu	11e 535	Ser	Leu	Ile	Phe	Ile 540	Туг	Asp	Ala	Val
Gly 545	Lys	Met	Leu	Asn	Leu 550	Thr	His	Thr	Tyr	Pro 555	Ile	Gln	Lys	Pro	Gly 560
Ser	Ser	Ala	Туr	Gly 565	Cys	Leu	Cys	Gln	Tyr 570	Pro	Gly	Pro	Gly	Gly 575	Asn
Glu	Ser	Gln	Trp 580	Ile	Arg	Thr	Arg	Pro 585	Lys	Asp	Arg	Asp	Asp 590	Ile	Val
Ser	Met	Asp 595	Leu	Gly	Leu	lle	Asn 600	Ala	Ser	Leu	Leu	Pro 605	Pro	Pro	Glu
Cys	Thr 610	Arg	Gln	Gly	Gly	His 615	Pro	Arg	Gly	Pro	Gly 620	Cys	His	Thr	Val
Pro 625	Asp	Ile	Ala	Phe	Phe 630	Ser	Leu	Leu	Leu	Phe 635	Leu	Thr	Ser	Phe	Phe 640
Phe	Ala	Met	Ala	Leu 645	Lys	Cys	Val	Lys	Thr 650	Ser	Arg	Phe	Phe	Pro 655	Ser
Val	Val	Arg	Lys 660	Gly	Leu	Ser	Asp	Phe 665	Ser	Ser	Val	Leu	Ala 670	Ile	Leu
Leu	Gly	Cys 675	Gly	Leu	Asp	Ala	Phe 680	Leu	Gly	Leu	Ala	Thr 685	Pro	Lys	Leu
Met	Val 690	Pro	Arg	Glu	Phe	Lys 695	Pro	Thr	Leu	Pro	Gly 700	Arg	Gly	Trp	Leu
Val 705	Ser	Pro	Phe	Gly	Ala 710	Asn	Pro	Trp	Trp	Trp 715	Ser	Val	Ala	Ala	Ala 720
Leu	Pro	Ala	Leu	Leu 725	Leu	Ser	lle	Leu	11e 730	Phe	Met	Asp	Gln	Gln 735	Ile
Thr	Ala	Val	11e 740	Leu	Asn	Arg	Met	Glu 745	Туr	Arg	Leu	Gln	Lys 750	Gly	Ala
Gly	Phe	His 755	Leu	Asp	Leu	Phe	Cys 760	Val	Ala	Val	Leu	Met 765	Leu	Leu	Thr
Ser	Ala 770	Leu	Gly	Leu	Pro	Trp 775	Tyr	Val	Ser	Ala	Thr 780	Val	Ile	Ser	Leu
Ala 785	His	Met	Asp	Ser	Leu 790	Arg	Arg	Glu	Ser	Arg 795	Ala	Cys	Ala	Pro	61y 800
Glu	Arg	Pro	Asn	Phe 805	Leu	Gly	Ile	Arg	Glu 810	Gln	Arg	Leu	Thr	Gly 815	
Val	Val	Phe	11e 820	Leu	Thr	Gly	Ala	Ser 825	Ile	Phe	Leu	Ala	Pro 830	Val	Leu

WO 03/089583 16U 200 PCT FINAL.ST25 Lys Phe Ile Pro Met Pro Val Leu Tyr Gly Ile Phe Leu Tyr Met Gly Val Ala Ala Leu Ser Ser Ile Gln Phe Thr Asn Arg Val Lys Leu Leu Leu Met Pro Ala Lys His Gln Pro Asp Leu Leu Leu Arg His Val Pro Leu Thr Arg Val His Leu Phe Thr Ala Ile Gln Leu Ala Cys Leu Gly Leu Leu Trp Ile Ile Lys Ser Thr Pro Ala Ala Ile Ile Phe Pro Leu Met Leu Leu Gly Leu Val Gly Val Arg Lys Ala Leu Glu Arg Val 915 920 925 Phe Ser Pro Gln Glu Leu Leu Trp Leu Asp Glu Leu Met Pro Glu Glu Glu Arg Ser Ile Pro Glu Lys Gly Leu Glu Pro Glu His Ser Phe Ser Gly Ser Asp Ser Glu Asp Ser Glu Leu Met Tyr Gln Pro Lys Ala Pro Glu Ile Asn Ile Ser Val Asn <210> 700 <211> <212> DNA <213> Homo sapiens <220> <221> CDS <222> (60)..(497) <223> <400> 92 gaaagaagga aataaacaca ggcaccaaac cactatccta agttgactgt cctttaaat atg tca aga tcc aga ctt ttc agt gtc acc tca gcg atc tca acg ata Met Ser Arg Ser Arg Leu Phe Ser Val Thr Ser Ala Ile Ser Thr Ile

59

107

ggg atc ttg tgt ttg ccg cta ttc cag ttg gtg ctc tcg gac cta cca Gly Ile Leu Cys Leu Pro Leu Phe Gln Leu Val Leu Ser Asp Leu Pro 155

tgc gaa gaa gat gaa atg tgt gta aat tat aat gac caa cac cct aat Cys Glu Glu Asp Glu Met Cys Val Asn Tyr Asn Asp Gln His Pro Asn

ggc tgg tat atc tgg atc ctc ctg ctg ctg gtt ttg gtg gca gct ctt Gly Trp Tyr Ile Trp Ile Leu Leu Leu Leu Val Leu Val Ala Ala Leu 50 60251

ctc tgt gga gct gtg gtc ctc tgc ctc cag tgc tgg ctg agg aga ccc Leu Cys Gly Ala Val Val Leu Cys Leu Gln Cys Trp Leu Arg Arg Pro 65 70 75 80299

cga att gat tot cac agg cgc acc atg gca gtt ttt gct gtt gga gac Arg Ile Asp Ser His Arg Arg Thr Met Ala Val Phe Ala Val Gly Asp 347 85 90

ttg gac tct att tat ggg aca gaa gca gct gtg agt cca act gtt gga 395

WO 03/089583 PCT/US03/11497

Leu	Asp	Ser	Ile 100	Tyr	Gly	Thr	Glu	Ala 105	Ala			PCT Pro				
		ctt Leu 115														443
		ggc Gly														491
aca Thr 145		tgat	ttta	agg t	igtg	gatta	at ca	atti	taaaq	g tai	taad	cgac	atci	tgtaa	att	547
ccaa	aaca	atc a	aatt	tage	ga at	agtt	attt	caç	gttgt	tgg	aaat	gtco	ag a	agato	ctattc	607
atat	agto	ctg a	aggaa	aggad	a at	tcga	acaaa	a aga	aatg	gatg	ttg	gaaaa	aa t	tttç	ggtcat	667
ggaç	gatgi	ttt a	aata	igtaa	aa gt	agca	aggct	tt	t.							700
<210 <211 <212 <213	.> : !> !	93 146 PRT Homo	sapi	ens												
< 400)> !	93														
Met 1	Ser	Arg	Ser	Arg 5	Leu	Phe	Ser	Val	Thr 10	Ser	Ala	Ile	Ser	Thr 15	Ile	
Gly	Ile	Leu	Cys 20	Leu	Pro	Leu	Phe	Gln 25	Leu	Val	Leu	Ser	Asp 30	Leu	Pro	
Cys	Glu	Glu 35	Asp	Glu	Met	Cys	Val 40	Asn	Tyr	Asn	Asp	Gln 45	His	Pro	Asn	
Gly	Trp 50	Tyr	Ile	Trp	Ile	Leu 55	Leu	Leu	Leu	Val	Leu 60	Val	Ala	Ala	Leu	
Leu 65	Cys	Gly	Ala	Val	Val 70	Leu	Cys	Leu	Gln	Cys 75	Trp	Leu	Arg	Arg	Pro 80	
Arg	Ile	Asp	Ser	His 85	Arg	Arg	Thr	Met	Ala 90	Val	Phe	Ala	Val	Gly 95	Asp	
Leu	Asp	Ser	Ile 100	Tyr	Gly	Thr	Glu	Ala 105	Ala	Val	Ser	Pro	Thr 110	Val	Gly	
lle	His	Leu 115	Gln	Thr	Gln	Thr	Pro 120	Asp	Leu	Tyr	Pro	Val 125	Pro	Ala	Pro	
	Phe 130	Gly	Pro	Leu		Ser 135	Pro	Pro	Pro	туг	Glu 140	Glu	Ile	Val	Lys	
Thr 145	Thr															
<210 <211 <212 <213	> 1 > 0	324 0NA tomo	sapi	ens										; *		<i>:</i>
<220 <221 <222 <223	> C	DS (44).	. (77	2)												

< 400	0 > -	94		,						160	200	PCT	FINA	AL.ST	25	
			gatg	ccct	t g g∉	cagg	gtga	g cc	caca	agga	gca			cag Gln		55
														act Thr		103
														aag Lys 35		151
														G] À		199
														cac His		247
														gtg Val		295
														cag Gln		343
														cca Pro 115		391
														ttc Phe		439
														cca Pro		487
														ccc Pro		535
														cct Pro		583
														gtt Val 195		631
														aat Asn		679
														tgt Cys		727
								gaa Glu								772
taga	ggct	at t	ctga	tata	a ta	acac	aatç	cto	agct	cag	ggaç	caaç	gtg t	ttco	gtcat	832
tgtt	acct	ga c	aaco	gtgg	jt gt	tcta	tgtt	gta	acct	tca	gaag	ttac	ag o	cagcg	cccag	892
gcag	jcct <u>c</u>	jac a	gaga	tcat	t ca	aggg	ggga	aaç	ggga	agt	ggga	ggtç	jca a	attto	tcaga	952
ttgg	,taa <i>a</i>	aa t	tago	ctgg	g ct	gggg	jaaat	tct	cctc	cgg	aaca	gttt	ca a	atto	cctcg	1012
ggta	agaa	at c	tcct	gtat	a ag	gttc	agga	gca	iggaa	ttt	cact	tttt	ca t	ccac	caccc	1072
tccc	cctt	ct c	tgta	iggaa	ig go	attg	gtgç	ctc	aatt	tta	acco	cago	ag o	caat	ggaaa	1132
aato	acga	ct t	ctga	gact	t tç	ggaç	tttc	cac	agaç	ıgtg	agag	tcgg	ıgt ç	gggaa	ggaag	1192

Page 66

			160	200 PCT FIN	IAI., ST25	
cagggaagag	aaagcaggcc	cagctggaga				1252
agactcacta	atcccaaaca	actcagctgc	catctggcct	ctctgaggac	tctgggtacc	1312
ttaaagacta	ta					1324
<210> 05						

243

<212> PRT

<213> Homo sapiens

Met Glu Gln Gly Ser Gly Arg Leu Glu Asp Phe Pro Val Asn Val Phe 1 5 10 15

Ser Val Thr Pro Tyr Thr Pro Ser Thr Ala Asp Ile Gln Val Ser Asp 20 25 30

Asp Asp Lys Ala Gly Ala Thr Leu Leu Phe Ser Gly Ile Phe Leu Gly 35 40

Leu Val Gly Ile Thr Phe Thr Val Met Gly Trp Ile Lys Tyr Gln Gly 50 60

Val Ser His Phe Glu Trp Thr Gln Leu Leu Gly Pro Val Leu Leu Ser 65 70 75 80

Val Gly Val Thr Phe Ile Leu Ile Ala Val Cys Lys Phe Lys Met Leu 85 90 95

Ser Cys Gln Leu Cys Lys Glu Ser Glu Glu Arg Val Pro Asp Ser Glu

Gln Thr Pro Gly Gly Pro Ser Phe Val Phe Thr Gly Ile Asn Gln Pro 115 120 125

Ile Thr Phe His Gly Ala Thr Val Val Gln Tyr Ile Pro Pro Pro Tyr

Gly Ser Pro Glu Pro Met Gly Ile Asn Thr Ser Tyr Leu Gln Ser Val 145 150 155 160

Val Ser Pro Cys Gly Leu Ile Thr Ser Gly Gly Ala Ala Ala Ala Met 165 170170175

Ser Ser Pro Pro Gln Tyr Tyr Thr Ile Tyr Pro Gln Asp Asn Ser Ala 180 185 190

Phe Val Val Asp Glu Gly Cys Leu Ser Phe Thr Asp Gly Gly Asn His 195 200 200 205

Arg Pro Asn Pro Asp Val Asp Gln Leu Glu Glu Thr Gln Leu Glu Glu 210 215 220

Glu Ala Cys Ala Cys Phe Ser Pro Pro Pro Tyr Glu Glu Ile Tyr Ser 225 230 235 240

Leu Pro Arg

<210> 96

<211> 5350	16U 200 PCT FINAL.ST25	
<212> DNA		
<213> Homo sapiens		
<220> <221> CDS		
<222> (2275)(3213)		
<223>		
<400> 96 gtccaggcgt accatgactc	tcacattttg cagttgtttt atttgacggg acagacattg	60
actgacagtg gctggagcag	ggctgatagt gaatttctga aacggtttac ctgattctct	120
gctttctgag ttcttggata	totgagagac agggootota tgotgtttca otgotggata	180
tatgcttcat tcttggacca	taatttttt ttcaaatttt tctagatgat gttgcttcat	240
	ataattccac tgaatttttg aagtttattg ggaattattt	300
	aaattacttt ctgctccagg aaaatatagt ttattagtct	360
agtaatttat taattgacta	aaatctacca tttgttatgg ccaatgacat gtttatttac	420
tgaaaataca ttaagtcccc	tttggtttta agtctcttaa cataagaaag caatttgtta	480
aaaactggca ttactttact	cttatgcttt ctgtgtcctt tgctaagtat ttctaaaaca	540
aaatgaaaac ccacgagttt	agtcttggcc agggcaagat atttgaaata aaaaaggaaa	600
taatatgacc aattgcaata	attcttattt ataaatttta agttaatgat aaaaaatata	660
aagtgtacat tacaatgtaa	aaggttacat aagaaaagct gcaatataaa aaggatgaat	720
atgtgtctga tttaaataaa	catttgacac gttattaata tattgaacat taatgatatc	780
taaaactatt cattttataa	aggatatgca ttttctttaa gtagagaata ataataatga	840
gcatccatat gtaaatcaca	gaattotgaa caagagaaag atagtgotat caacgggaaa	900
gggctgacca gcaccactga	cccccaaaa tagccaggta gaagaagagt cctacagcct	960
attacaaggt gattaattga	ctagatgctc tgagaagaaa ttggaacttg gatgatctga	1020
agatagttat ctcaattgat	tgttcacagc cagttacaga tagaattcct tgttctacat	1080
tttcctccct tctcactagt	gcacttgagt agtctttaaa aaaaattgca acttcagaga	1140
cccccatgct tgaaccactg	ggagaagaaa ccttaggatg acctacctgc atacaataaa	1200
tatgttggat gtcacgataa	gataagtata aattgaggca aactttetet caccaaaatt	1260
ctacaggcaa aatggggaga	ttggaagaaa agatgtgggc ttgtaaaatc caattacatt	1320
ttactttaat tttataaaga	aggitcacat caagaaatte caagigaggi teagaccaat	1380
cacctcagaa taaactgatt	ggatgataat gctgattcct aaagcatcat tgatctgaga	1440
tagccataat tttttttga	tatcttgaaa gattggcaga aacacaacgg attagaacat	1500
cttgatggaa attatgaaaa	tatgaataaa taactcacaa gattaatgtc tttgtaatag	1560
gttaagtgga agtataaaaa	tacattttat aaatcacata tgtgtaaaag taaatcattt	1620
	actagtgtct ttaatacatt taaagaaatt tgactaaatt	1680
tgtaacgtta tataagggtt	tggaatttta tgtttaaaat gtttacaatt actggtggct	1740
	ttgaaaaatt gtatgttcgt agatttgtaa cgagatttaa	1800
	ccttttttgc agacatgact cttgagggtc aaatatatag	1860
	agctctgtaa aatcccatgg gaatgggatt tgggcaatac	1920
	tactaacaca cacaaaatgt gaacatatat aagtaaaaat	1980
	tctataggaa ataatttaat ttcagttgta tggacctctt	2040
cattgagaat ataaatattt	cattcccatt ctagatgggg aatcagattc acaatctaat	2100

gtg	ctgt	ctc	tttt	tagt	gc a	aatt	caca	g tt	catg	ttgg	aaa	taca	ctc	tgat	tttcac	2160
att	gatt	ttt	aaaa	ggta	aa g	tgaa	gcaa	a ca	tact	ttta	cgt	ggta	cac	acat	gattat	2220
aaa	taaa	gtt	tact	tttg	tc c	tcca	ggta	a ag	ccac	ttca	gcc	gato	ata	cagc	atg Met 1	2277
cgg Arg	ctg Leu	gcc Ala	aac Asn 5	cag Gln	acc Thr	ctg Leu	ggt Gly	ggt Gly 10	gac Asp	ttt Phe	t t c Phe	ctg Leu	ttg Leu 15	gga Gly	atc Ile	2325
ttc Phe	agc Ser	cag Gln 20	atc	tca Ser	cac His	cct Pro	ggc Gly 25	cgc Arg	ctc Leu	tgc Cys	t t g Leu	ctt Leu 30	ato Ile	ttc Phe	agt Ser	2373
										aca Thr						2421
										tac Tyr 60						2469
										gtc Val						2517
ctg Leu	gtg Val	aac Asn	cag Gln 85	ctg Leu	gcc Ala	aaa Lys	gac Asp	aag Lys 90	acc Thr	atc Ile	tcg Ser	gtc Val	ctt Leu 95	ggg Gly	tgt Cys	2565
			Met							gga Gly						2613
										gtg Val						2661
ctc Leu 130	cgt Arg	tac Tyr	tct Ser	gtg Val	ctc Leu 135	atg Met	agc Ser	cat His	agg Arg	gta Val 140	tgt Cys	ctc Leu	ctc Leu	ctg Leu	gca Ala 145	2709
tca Ser	ggc Gly	tgc Cys	tgg Trp	ttt Phe 150	gtg Val	ggc Gly	tca Ser	gtg Val	gat Asp 155	ggc Gly	ttc Phe	atg Met	ctc Leu	act Thr 160	ccc Pro	2757
atc Ile	gcc Ala	atg Met	agc Ser 165	ttc Phe	ccc Pro	ttc Phe	tgc Cys	aga Arg 170	tcc Ser	cat His	gag Glu	att Ile	cag Gln 175	cac His	ttc Phe	2805
										tct Ser						2853
ctt Leu	tac Tyr 195	aag Lys	att Ile	ttc Phe	atg Met	tac Tyr 200	ttg Leu	tgc Cys	tgt Cys	gtc Val	atc Ile 205	atg Met	ctc Leu	ctg Leu	ata Ile	2901
cct Pro 210	gtg Val	acg Thr	gtc Val	att Ile	tca Ser 215	gtg Val	tct Ser	tac Tyr	tac Tyr	tat Tyr 220	atc Ile	atc Ile	ctc Leu	acc Thr	atc Ile 225	2949
cat His	aag Lys	atg Met	aac Asn	tca Ser 230	gtt Val	gag Glu	ggt Gly	cgg Arg	aaa Lys 235	aag Lys	gcc Ala	ttc Phe	acc Thr	acc Thr 240	tgc Cys	2997
tcc Ser	tcc Ser	cac His	att Ile 245	aca Thr	gtg Val	gtc Val	agc Ser	ctc Leu 250	ttc Phe	tat Tyr	gga Gly	gct Ala	gct Ala 255	att Ile	tac Tyr	3045
										cct Pro						3093
										gtc Val						3141

16U 200 PCT FINAL.ST25 275 280 285

tac agt ttc agg aat aag gat gtc aca agg gct g Tyr Ser Phe Arg Asn Lys Asp Val Thr Arg Ala 1 290 295 300		3189
agc gtg cag aaa cct cca tat taa agtgtgaaag a Ser Val Gln Lys Pro Pro Tyr 310	acttaagtt ggtcctctct	3243
tcttagagtc tctcttcact ttaggtgtcc ttccaccaaa	caatcagcat attgtggtag	3303
tgtctgactc cctgagttgt ccttcagggg gattcagccc a	agtgttcttc cctcctataa	3363
tcacacttga gatgatgttc acttatcccc cccttccctc (gtagcattga tctctagtcc	3423
agtccttcgg ggccaatggt cctttttta gattacagtg	gagaaatatg aaaataaatg	3483
tgtttatgac ccttgagcac cttccaccac agagaaaatt	tgttttgcta tcatgggccc	3543
attgatgagt atgaaataac accatattca gagtgttcct	cagcatccac tctgtgctaa	3603
acgetttteg tteaceacet cattegacet teaceetetg	tggctgaggc taaggtcacc	3663
cacatttcac aaatgacaaa acagcctttg aggcttcccc	tgacttgccc caagcagggg	3723
atcctcaggg acaagggggt tcattcatcc ataggcattt	ggagataaac acattcaaga	3783
cctcagagat gctaaatgta cagttgagat ttttcttcca	tcagaatttc tagaatgtgt	3843
tctcaatcaa attcttattt tctgtgagca tataagaagt o	caaacctccc aaaattagag	3903
cagagacatg ggctatccag tagacatggg ctacaacatg	tttggagtat aattggttta	3963
ttcatagact taaccagaga aatatggaag tttcgcacac (ttctccctgt tcaagccaat	4023
ggtgacacat acttagaata taatttcaaa tcacagtttt a	acgtatgtgc atggttgtat	4083
ttgtatttaa caaataacat aataattata acgtcttgtg	tgattattat gatctggcac	4143
catttttagt gcgtgacatg tatggaacac ttttattttc a	acagootatt attagootta	4203
ttctacagtt gatttaactg aaacccatgg gtttgagtaa	catgaacaaa agggtgtgca	4263
gcttataaag tgctcaacag ggatttaagc ccaggcaggc a	aggccggagt ccctgcccct	4323
gaccactgca tgtgccacgt cttgtggagt ctgtggcctt	ttccacactg cattgcctct	4383
ccctctggga gggccatact ccaaccttgg aaacactata	gttctttcca tacccaatgt	4443
tttcacgtgg ctttccctct cttcggaatg ttttttatc	tgtaagtaca aggatacgaa	4503
gataactttc catgactaca taatcttcct ttaggcccca a	agtcattcat tcattcaaca	4563
aataactact gagcccctat agtttgccag gccccgttct a	acaaactgag gatacatcag	4623
tgagcaaaac aaataaaaat cttcatcttt tttagcactt a	aaagggtgta tacagaaaat	4683
aaatttggta attgagaaga agacatggag tattatcaga a	agaaaagtgt tggaaaatct	4743
tgagcaggag aggggtcttg gagtgtgtag gggtcacgtt t	ttatgtaggg atttaggcta	4803
atcctcactg gttatagttg agcaaagatg tggagtttac a	aagttaatga gccacattga	4863
tattgggaga aatgctttca agacagagca tagggacatc t	taccageetg teaateaaga	4923
gtccagtagg accatgtctc agtaataggg atgaactaga t	tgtagattga gtctaactcc	4983
aattataaaa aatgatagta aaataaattt ttccaacaaa o	caaaagtgga taaaattctt	5043
cagccataga aaaattatct caaaagtaaa ctcagaaata t	taagcaaaaa tgacaacat	5103
caaccccaaa gagtaatatg taaatgagtc ataatcaata t	ttgacttagc aataatttta	5163
atgtaatata cagtttagat ttgtgcaaaa cttaaatgta t	tgaaaaatca tgttgaagat	5223
atatcaatat tgatgtagta gaggtagtaa attgtgttaa a	accttagtaa atcaagagta	5283
catgctgtaa tgtttatagt aaacgccaaa accagtttat a	aaaatgaaaa aatgatagat	5343
	Page 70	

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

5350

cccaca					_
<210> 97 <211> 312 <212> PRT <213> Homo	sapiens				
<400> 97					
Met Arg Leu 1	Ala Asn Gln 5	Thr Leu Gl	y Gly Asp Phe 10	e Phe Leu Leu 15	Gly
Ile Phe Ser	Gln Ile Ser 20	His Pro Gl 25	y Arg Leu Cys	Leu Leu Ile 30	Phe
Ser Ile Phe 35	Leu Met Ala	Val Ser Tr 40	p Asn Ile Thr	Leu Ile Leu 45	Leu
lle His Ile 50	: Asp Ser Ser	Leu His Th	r Pro Met Tyr 60	Phe Phe Ile	Asn
Gln Leu Ser 65	Leu Ile Asp 70	Leu Thr Ty	r Ile Ser Val 75	Thr Val Pro	Lys 80
Met Leu Val	Asn Gln Leu 85	Ala Lys As	p Lys Thr Ile 90	e Ser Val Leu 95	Gly
Cys Gly Thr	Gln Met Tyr 100	Phe Tyr Le	eu Gln Leu Gly 5	Gly Ala Glu 110	Cys
Cys Leu Leu 115		Ala Tyr As 120	p Arg Tyr Val	Ala Ile Cys 125	His
Pro Leu Arg 130	Tyr Ser Val	Leu Met Se 135	er His Arg Val 140		Leu
Ala Ser Gly 145	Cys Trp Phe 150		r Val Asp Gly 155	Phe Met Leu	Thr 160
Pro Ile Ala	Met Ser Phe 165	Pro Phe Cy	s Arg Ser His 170	Glu Ile Gln 175	
Phe Phe Cys	Glu Val Pro 180	Ala Val Le 18	u Lys Leu Ser 5	Cys Ser Asp 190	Thr
Ser Leu Tyr 195		Met Tyr Le 200	u Cys Cys Val	Ile Met Leu 205	Leu
Ile Pro Val 210	Thr Val Ile	Ser Val Se 215	r Tyr Tyr Tyr 220		Thr
Ile His Lys 225	Met Asn Ser 230		y Arg Lys Lys 235	Ala Phe Thr	Thr 240
Cys Ser Ser	His Ile Thr 245	Val Val Se	r Leu Phe Tyr 250	Gly Ala Ala 255	
Tyr Asn Tyr	Met Leu Pro 260	Ser Ser Ty 26	r Gln Thr Pro 5	Glu Lys Asp 270	Met

Page 71

ttttata

Met Ser Ser Phe Phe Tyr Thr Ile Leu Thr Pro Val Leu Asn Pro Ile 275 280 285

Ile Tyr Ser Phe Arg Asn Lys Asp Val Thr Arg Ala Leu Lys Lys Met 290 295 300

Leu Ser Val Gln Lys Pro Pro Tyr

<210> 98 <211> 3486 <212> DNA <213> Homo sapiens

<221> CDS <222> (1)..(3483)

<222> (1)..(348 <223>

<400> 98

atg ggg cca cct gaa ttc atg tat gaa cag cag gac aat tca acg cac 48

Met Gly Pro Pro Glu Phe Met Tyr Glu Gln Gln Asp Asn Ser Thr His

1 10 15

ctg Cag Cca ctt aag aca tgc Ccc gtg gca agg cag Cta atc cga ggg 96 Leu Gln Pro Leu Lys Thr Cys Pro Val Ala Arg Gln Leu Ile Arg Gly 20 25 30

gtg ctg cgg gca cct gat gga gcc aag cca gga gag gac agg ggc cag 144 Val Leu Arg Ala Pro Asp Gly Ala Lys Pro Gly Glu Asp Arg Gly Gln 35 40 45

gcc cgc tgc aat gga cgt gta tgt gga gag aaa tca aaa caa cct att 192 Ala Arg Cys Asn Gly Arg Val Cys Gly Glu Lys Ser Lys Gln Pro Ile 50 55

gag gct ttt aag ccc gtc tgc tac aaa ccc caa ttt atg tcc cac att
Glu Ala Phe Lys Pro Val Cys Tyr Lys Pro Gln Phe Met Ser His Ile
65 70 80

att ccc ctt tac tcc atc cat gca tcc cag agt tcc agc caa tcc aag

Ile Pro Leu Tyr Ser Ile His Ala Ser Gln Ser Ser Gln Ser Lys

85

90

95

ctg cct gca cat ctc cat ttg gac ccc tta ggc tgt gcc agt ctc agc
Leu Pro Ala His Leu His Leu Asp Pro Leu Gly Cys Ala Ser Leu Ser

ttc tcc tcc acc cag ccc tca cca cct tat tac cca ggg ttg gta cta

Phe Ser Ser Thr Gln Pro Ser Pro Pro Tyr Tyr Pro Gly Leu Val Leu

gga tgc agc aag cag aat act gga ggt gca aaa tgt cag aag cca ctc 432 Gly Cys Ser Lys Gln Asn Thr Gly Gly Ala Lys Cys Gln Lys Pro Leu 130 135 140

act cgc agg ttt gag cac ttg gga aca gca aag aag ccc aag aaa tca
Thr Arg Arg Phe Glu His Leu Gly Thr Ala Lys Lys Pro Lys Lys Ser

gtc tgg cca ctg cag agc ctg cct caa aga gat ttg aag ctg gtc aat
Val Trp Pro Leu Gln Ser Leu Pro Gln Arg Asp Leu Lys Leu Val Asn

gca agg agc cag gcc tgc tgg aat cca agg acc tgg ggt gca gca acc
Ala Arg Ser Gln Ala Cys Trp Asn Pro Arg Thr Trp Gly Ala Ala Thr
180
180

cca gat aca gac cct gaa gag gcc aac agc ggt cag cag aac ata aag
Pro Asp Thr Asp Pro Glu Glu Ala Asn Ser Gly Gln Gln Asn Ile Lys

gag Caa cag tac cgt gtc tct ctg ggg aac aac act ggt tct ccc ttg
Glu Gln Gln Tyr Arg Val Ser Leu Gly Asn Asn Thr Gly Ser Pro Leu
210 215 220

								agg Arg	cag	cag	aag	ctg	aat	720
								gcc Ala 250						768
								cag Gln						816
								gaa Glu						864
								ggt Gly						912
								cag Gln						960
								tgg Trp 330						1008
								gcc Ala						1056
								gcc Ala						1104
								gaa Glu						1152
								gtg Val						1200
								gca Ala 410						1248
								gtc Val						1296
		Pro	Gln		Lys	Ğİy	Ser	ggt Gly		Glu				1344
								tgc Cys						1392
_	-	-		-			-	cct Pro				-	-	1440
								gga Gly 490						1488
								cca Pro						1536
								gaa Glu						1584
								atg Met						1632

Page 73

			agc Ser					1680
			acc Thr					1728
			tcc Ser					1776
			ggg Gly					1824
			tct Ser 615					1872
			gcc Ala					1920
			aca Thr					1968
			gaa Glu					2016
			ttc Phe					2064
			cag Gln 695					2112
			ccg Pro					2160
			cag Gln					2208
			tca Ser					2256
			acc Thr					2304
			aag Lys 775					2352
			tac Tyr					2400
			ctc Leu					2448
			gac Asp					2496
			ccg Pro					2544
			tgc Cys					2592

	850					855				1 6 U		00 P 60	СŦ	FIN	AL.S	T25	5	
	ggc Gly										. S					e C		2640
	aac Asn															n G		2688
gcc Ala	cgg Arg	ccg Pro	cac His 900	tgc Cys	cgg Arg	ctg Leu	gtg Val	gcc Ala 905	ctg Leu	gac Asp	g g g	gg c	ac is	ctg Leu 910	Ту	t ge	CC la	2736
	ggc Gly											/r A						2784
	cgc Arg 930											зр Т						2832
	cac His										P					/ G		2880
	ctg Leu															T		2928
	gcc Ala														Met			2976
	gtc Val							Ph€	: gad : As _l			Asn .		S	-	_	ggc Gly	3024
	gcc Ala 1010	Val					Al	c aç .a. S∈					T		tac Tyr			3069
_	gcc Ala 1025	Thr	tac Tyr				ту	r Pr					G		tgc Cys			3114
	gtg Val 1040	Asp	aac Asn				· cy	ic gt 's Va					S		acc Thr			3159
Cys	ttc Phe 1055	Leu	Ala	Asp	Ser	Val 106	\$e 0	a co r Pr	o Ai	rg S	er	Val 106	A 5	la	gtc Val	Ph€	>	3204
Leu	tct Ser 1070	Gly	Ser	Trp	Gly	Asn 107	Hi 5	s Hi	s G	ln S	er	Ala 1080	D D	tt eu	cag Gln	ggt Gly	/	3249
	agc Ser 1085	Пlе					СУ						G		cta Leu			3294
	gtg Val 1100	Ser	acg Thr				G1						Ğ					3339
	aga Arg 1115	Ser					Ly						V		ggc Gly			3384
gac Asp	ata Ile 1130	Arg	gga Gly				Le				rg		P		cca Pro			3429
ctg Leu	-		gct Ala				G1				er		S					3474
gcc	aca	ggg	tga															3486

Page 75

Ala Thr Gly 1160

<210> 99

<211> 1161 <212> PRT

<213> Homo sapiens

<400> 99

Leu Gln Pro Leu Lys Thr Cys Pro Val Ala Arg Gln Leu Ile Arg Gly 20 25 30

Val Leu Arg Ala Pro Asp Gly Ala Lys Pro Gly Glu Asp Arg Gly Gln 35 40 45

Ala Arg Cys Asn Gly Arg Val Cys Gly Glu Lys Ser Lys Gln Pro Ile 50 55

Glu Ala Phe Lys Pro Val Cys Tyr Lys Pro Gln Phe Met Ser His Ile 65 70 80

Ile Pro Leu Tyr Ser Ile His Ala Ser Gln Ser Ser Ser Gln Ser Lys $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$

Leu Pro Ala His Leu His Leu Asp Pro Leu Gly Cys Ala Ser Leu Ser 100 105 110

Phe Ser Ser Thr Gln Pro Ser Pro Pro Tyr Tyr Pro Gly Leu Val Leu 115 120 125

Gly Cys Ser Lys Gln Asn Thr Gly Gly Ala Lys Cys Gln Lys Pro Leu 130 135 140

Thr Arg Arg Phe Glu His Leu Gly Thr Ala Lys Lys Pro Lys Lys Ser 145 150 155 160

Val Trp Pro Leu Gln Ser Leu Pro Gln Arg Asp Leu Lys Leu Val Asn 165 170 175

Ala Arg Ser Gin Ala Cys Trp Asn Pro Arg Thr Trp Gly Ala Ala Thr $180 \\ \hspace{1.5cm} 185 \\ \hspace{1.5cm} 190 \\ \hspace{1.5cm}$

Pro Asp Thr Asp Pro Glu Glu Ala Asn Ser Gly Gln Gln Asn Ile Lys $195 \hspace{1.5cm} 200 \hspace{1.5cm} 205 \hspace{1.5cm}$

Glu Gln Gln Tyr Arg Val Ser Leu Gly Asn Asn Thr Gly Ser Pro Leu 210 215 220

Cys Ser Thr Glu Val Asn Phe Gly Ser Arg Gln Gln Gly Lys Leu Asn 225 230 235 240

Arg Thr Thr Arg Glu Ala Trp Lys Glu Ala Ser Arg Trp Asp Leu Pro 245 250 250 255.

Ala Leu Gly Pro Ser Gly His Pro Leu Gln Leu Lys Val Thr Phe Ala 260 265 270

Pro Leu Leu Ser Ser Ala Gly Gln Pro Glu Pro Ala Gln Asn Ser Leu 275 280 285

Pro Ser Ala Gln Gln Asp Pro Gly Thr Gly Pro Tyr Trp Ala Ile Ile 290 295 300

Asn Gln Ile Leu Asp Ile Pro Gln Pro Gln Val Gly Trp Arg Ser Met 305 310 315 320

Phe Pro Arg Gly Ala Glu Ala Gln Asp Trp His Leu Asp Met Gln Leu 325 330 335

Thr Gly Lys Val Val Leu Ser Ala Ala Ala Leu Leu Leu Val Thr Val $340 \hspace{1.5cm} 345 \hspace{1.5cm} 350$

Ala Tyr Arg Leu Tyr Lys Ser Arg Pro Ala Pro Ala Gln Arg Trp Gly 355 360 365

Gly Asn Gly Gln Ala Glu Ala Lys Glu Glu Ala Glu Gly Ser Gly Gln 370 380

Pro Ala Val Gln Glu Ala Ser Pro Gly Val Leu Leu Arg Gly Pro Arg 385 390 395 400

Arg Arg Arg Ser Ser Lys Arg Ala Glu Ala Pro Gln Gly Cys Ser Cys 405 410 415

Glu Asn Pro Arg Gly Pro Tyr Val Leu Val Thr Gly Ala Thr Ser Thr 420 425 430

Asp Arg Lys Pro Gln Arg Lys Gly Ser Gly Glu Glu Arg Gly Gln Gln 435 . 445

Gly Ser Asp Ser Glu Gln Val Pro Pro Cys Cys Pro Ser Gln Glu Thr 450 460

Arg Thr Ala Val Gly Ser Asn Pro Asp Pro Pro His Phe Pro Arg Leu 465 470 470 480

Gly Ser Glu Pro Lys Ser Ser Pro Ala Gly Leu Ile Ala Ala Ala Asp $485 \hspace{1.5cm} 490 \hspace{1.5cm} 495$

Gly Ser Cys Ala Gly Gly Glu Pro Ser Pro Trp Gln Asp Ser Lys Pro 500 505 510

Arg Glu His Pro Gly Leu Gly Gln Leu Glu Pro Pro His Cys His Tyr 515 520 525

Val Ala Pro Leu Gln Gly Ser Ser Asp Met Asn Gln Ser Trp Val Phe 530 535 540

Thr Arg Val Ile Gly Val Ser Arg Glu Glu Ala Gly Ala Leu Glu Ala 545 550 560

Ala Ser Asp Val Asp Leu Thr Leu His Gln Gln Glu Gly Ala Pro Asn 565 570 575

Ser Ser Tyr Thr Phe Ser Ser Ile Ala Arg Val Arg Met Glu Glu His 580 590

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Phe Ile Gln Lys Ala Glu Gly Val Glu Pro Arg Leu Lys Gly Lys Val 595 600 605

Tyr Asp Tyr Tyr Val Glu Ser Thr Ser Gln Ala Ile Phe Gln Gly Arg 610 615 620

Leu Ala Pro Arg Thr Ala Ala Leu Thr Glu Val Pro Ser Pro Arg Pro 625 630 635 640

Pro Pro Gly Ser Leu Gly Thr Gly Ala Ala Ser Gly Gly Gln Ala Gly 645 650 655

Asp Thr Lys Gly Ala Ala Glu Arg Ala Ala Ser Pro Gln Thr Gly Pro $660 \hspace{1cm} 665 \hspace{1cm} 670 \hspace{1cm}$

Trp Pro Ser Thr Arg Gly Phe Ser Arg Lys Glu Ser Leu Leu Gln Ile 675 680 685

Ala Glu Asn Pro Glu Leu Gln Leu Gln Pro Asp Gly Phe Arg Leu Pro 690 695 700

Ala Pro Pro Cys Pro Asp Pro Gly Ala Leu Pro Gly Leu Gly Arg Ser 705 710 715 720

Ser Arg Glu Pro His Val Gln Pro Val Ala Gly Thr Asn Phe Phe His $725 \hspace{1.5cm} 730 \hspace{1.5cm} 735$

Ile Pro Leu Thr Pro Ala Ser Ala Pro Gln Val Arg Leu Asp Leu Gly 740 745 750

Asn Cys Tyr Glu Val Leu Thr Leu Ala Lys Arg Gln Asn Leu Glu Ala 755 760 765

Leu Lys Glu Ala Ala Tyr Lys Val Met Ser Glu Asn Tyr Leu Gln Val $770 \ \ 775 \ \ 780$

Leu Arg Ser Pro Asp Ile Tyr Gly Cys Leu Ser Gly Ala Glu Arg Glu 785 790 795 800

Leu Ile Leu Gin Arg Arg Leu Arg Gly Arg Gln Tyr Leu Val Val Ala 805 810 815

Asp Val Cys Pro Lys Glu Asp Ser Gly Gly Leu Cys Cys Tyr Asp Asp 820 825 830

Glu Gln Asp Val Trp Arg Pro Leu Ala Arg Met Pro Pro Glu Ala Val 835 840 845

Ser Arg Gly Cys Ala Ile Cys Ser Leu Phe Asn Tyr Leu Phe Val Val 850 855 860

Ser Gly Cys Gln Gly Pro Gly His Gln Pro Ser Ser Arg Val Phe Cys 865 870 875 880

Tyr Asn Pro Leu Thr Gly Ile Trp Ser Glu Val Cys Pro Leu Asn Gln 885 890 895

Ala Arg Pro His Cys Arg Leu Val Ala Leu Asp Gly His Leu Tyr Ala 900 910

Ile Gly Gly Glu Cys Leu Asn Ser Val Glu Arg Tyr Asp Pro Arg Leu 915 920 925

Asp Arg Trp Asp Phe Ala Pro Pro Leu Pro Ser Asp Thr Phe Ala Leu

Ala His Thr Ala Thr Val Arg Ala Lys Glu Ile Phe Val Thr Gly Gly 945 950 950 960

Ser Leu Arg Phe Leu Leu Phe Arg Phe Ser Ala Gln Glu Gln Arg Trp 965 970 975

Trp Ala Gly Pro Thr Gly Gly Ser Lys Asp Arg Thr Ala Glu Met Val $980 \hspace{1.5cm} 995 \hspace{1.5cm} 995$

Ala Val Asn Gly Phe Leu Tyr Arg Phe Asp Leu Asn Arg Ser Leu Gly

Ile Ala Val Tyr Arg Cys Ser Ala Ser Thr Arg Leu Trp Tyr Glu 1010 1020

Cys Ala Thr Tyr Arg Thr Pro Tyr Pro Asp Ala Phe Gln Cys Ala

Val Val Asp Asn Leu Ile Tyr Cys Val Gly Arg Arg Ser Thr Leu 1040 1045 1050

Cys Phe Leu Ala Asp Ser Val Ser Pro Arg Ser Val Ala Val Phe 1055 1060 1065

Leu Ser Gly Ser Trp Gly Asn His His Gln Ser Ala Leu Gln Gly 1070 1075 1080

Asp Ser Ile Ile Cys Pro Pro Cys Ala Arg Trp Ser Gln Leu Asp 1085 1090 1095

Pro Val Ser Thr Glu Ala Ala Gly Ala Gln Ala Val Gly Leu Val 1100 1105 1110

Gly Arg Ser Arg Thr Gly Thr Lys Asp Glu Lys Glu Val Gly Met 1115 1120 1125

Asp Ile Arg Gly Glu Leu Ala Leu Asp His Arg Arg Pro Pro Ser 1130 1140

Leu Val \mbox{Trp} Ala Leu Ala Pro \mbox{Gly} Ser Ala Ser \mbox{Gly} Ser Ser \mbox{Glu} 1145 $\mbox{1150}$ 1155

Ala Thr Gly 1160

<210> 100 <211> 2953

<212> DNA <213> Homo sapiens

<220>

<221> CDS

<222> (350)..(2536)

<223>

Page 79

160 200 PC1 FINAL.S1	25
<400> 100 ctcagetetg caaagagete etceeeetee aggaaaagea etteteeeca ggeag	gggcgg 60
tcagtcccct cagcgcacct gcatgcacac acaccacctc acaaccacac actgo	catgca 120
cacacataca ccacagccac acactgtgca tacacgcaca ccccacaacc cacat	actgc 180
atgcacacac acacacctac aagctgcatg ctgcatacac aagtcataca ggaga	ataaac 240
tcagagtccc agccccaaat agaccccatc tcttgctcag ttgctgtcat cctag	gacctg 300
tttctttcgc cacatttcta taatctgcca gtgtctgcaa ggagaagac atg gg	
Met GI	ty Gly
ttt cta cct aag gca gaa ggg ccc ggg agc caa ctc cag aaa ctt Phe Leu Pro Lys Ala Glu Gly Pro Gly Ser Gln Leu Gln Lys Leu 5 10	
ccc tcc ttt ctg gtc aga gaa caa gac tgg gac cag cac ctg gac Pro Ser Phe Leu Val Arg Glu Gln Asp Trp Asp Gln His Leu Asp 20 25 30	
ctt cat atg ctg cag cag aag agg att cta gag tct cca ctg ctt Leu His Met Leu Gln Gln Lys Arg Ile Leu Glu Ser Pro Leu Leu 40 45 50	
gca tcc aag gaa aat gac ctg tct gtt ctt agg caa ctt cta ctg Ala Ser Lys Glu Asn Asp Leu Ser Val Leu Arg Gln Leu Leu 55 60 65	
tgc acc tgt gac gtt cga caa aga gga gcc ctg ggg gag acg gcg Cys Thr Cys Asp Val Arg Gln Arg Gly Ala Leu Gly Glu Thr Ala 70 80	
cac ata gca gcc ctc tat gac aac ttg gag gcg gcc ttg gtg ctg His Ile Ala Ala Leu Tyr Asp Asn Leu Glu Ala Ala Leu Val Leu 85 90 95	
gag gct gcc cca gag ctg gtc ttt gag ccc acc aca tgt gag gct Glu Ala Ala Pro Glu Leu Val Phe Glu Pro Thr Thr Cys Glu Ala 100 105 110	
gca ggt cag act gca ctg cac atc gct gtt gtg aac cag aat gtg Ala Gly Gln Thr Ala Leu His Ile Ala Val Val Asn Gln Asn Val 120 125 130	
Ctg gtg cgt gcc ctg ctc acc cgc agg gcc agt gtc tct gcc aga Leu Val Arg Ala Leu Leu Thr Arg Arg Ala Ser Val Ser Ala Arg 135 140 145	
aca ggc act gcc ttc cgc cgt agt ccc cgc aac ctc atc tac ttt Thr Gly Thr Ala Phe Arg Arg Ser Pro Arg Asn Leu Ile Tyr Phe 150 155 160	
gag cac cct ttg tcc ttt gct gcc tgt gtg aac agc gag gag atc Glu His Pro Leu Ser Phe Ala Ala Cys Val Asn Ser Glu Glu Ile 165 170 175	
cgg ctg ctc att gag cat gga gct gac atc agg gcc cag gac tcc Arg Leu Leu Ile Glu His Gly Ala Asp Ile Arg Ala Gln Asp Ser 180 185 190	
gga aac aca gta tta cac atc ctc atc ctc cag ccc aac aaa acc Gly Asn Thr Val Leu His Ile Leu Ile Leu Gln Pro Asn Lys Thr 200 205 210	
gcc tgc cag atg tac aac ctg ctg ctg tcc tac gat gga cat ggg Ala Cys Gln Met Tyr Asn Leu Leu Leu Ser Tyr Asp Gly His Gly 215 220 225	
cac ctg cag ccc ctg gac ctt gtg ccc aat cac cag ggt ctc acc His Leu Gln Pro Leu Asp Leu Val Pro Asn His Gln Gly Leu Thr 230 235 240	
ttc aag ctg gct gga gtg gag ggt aac act gtg atg ttc cag cac Phe Lys Leu Ala Gly Val Glu Gly Asn Thr Val Met Phe Gln His 245 250 255	

atg Met 260	Gln	aag Lys	cgg Arg	agg Arg	cac His 265	atc Ile	cag Gln	tgg Trp	acg Thr	tat Tyr 270	gga Gly	ccc Pro	ctg Leu	acc Thr	tcc Ser 275	1174
att Ile	ctc Leu	tac Tyr	gac Asp	ctc Leu 280	aca Thr	gag Glu	atc Ile	gac Asp	tcc Ser 285	tgg Trp	gga Gly	gag Glu	gag Glu	ctg Leu 290	tcc Ser	1222
					gtc Val											1270
ctg Leu	gaa Glu	cag Gln 310	acc Thr	cca Pro	gtg Val	aag Lys	gag Glu 315	ctg Leu	gtg Val	agc Ser	ttc Phe	aag Lys 320	tgg Trp	aac Asn	aag Lys	1318
tat Tyr	ggc Gly 325	cgg Arg	ccg Pro	tac Tyr	ttc Phe	tgc Cys 330	atc Ile	ctg Leu	gct Ala	gcc Ala	ttg Leu 335	tac Tyr	ctg Leu	ctc Leu	tac Tyr	1366
	Ile				acg Thr 345											1414
					cat His											1462
					tat Tyr											1510
					atc Ile											1558
					agg Arg											1606
					ttc Phe 425											1654
gtg Val	ctg Leu	gtg Val	acc Thr	atg Met 440	gtg Val	atg Met	cgg Arg	ctc Leu	acc Thr 445	aac Asn	acc Thr	aat Asn	ggg Gly	gag Glu 450	gtg Val	1702
gtg Val	ccc Pro	atg Met	tcc Ser 455	ttt Phe	gcc Ala	ctg Leu	gtg Val	ctg Leu 460	ggc Gly	tgg Trp	tgc Cys	agt Ser	gtc Val 465	atg Met	tat Tyr	1750
					cag Gln											1798
					gac Asp											1846
					gcc Ala 505											1894
					61 y 999											1942
					ttt Phe											1990
					ttc Phe											2038
atc Ile	att Ile	gcc Ala	aca Thr	ctg Leu	ctc Leu	atg Met	ctc Leu	aac Asn	ttg Leu	ttc Phe	atc Ile	gcc Ala	atg Met	atg Met	ggc Gly	2086

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25 575 575	
gac acc cac tgg agg gtg gcc cag gag agg gat gag ctc tgg agg gcc Asp Thr His Trp Arg Val Ala Gln Glu Arg Asp Glu Leu Trp Arg Ala 580 585 590 595	2134
Cag gtc gtg gcc acc aca gtg atg ctg gag cgg aag ctg cct cgc tgc Gln Val Val Ala Thr Thr Val Met Leu Glu Arg Lys Leu Pro Arg Cys 600 605 610	2182
Ctg tgg cct cgc tcc ggg atc tgt ggg tgc gaa ttc ggg ctg ggg gac Leu Trp Pro Arg Ser Gly Ile Cys Gly Cys Glu Phe Gly Leu Gly Asp 615 620 625	2230
cgc tgg ttc ctg cgg gtt gag aac cac aat gat cag aat cct ctg cga Arg Trp Phe Leu Arg Val Glu Asn His Asn Asp Gln Asn Pro Leu Arg 630 635 640	2278
gtg ctt cgc tat gtg gaa gtg ttc aag aac tca gac aag gag gat gac Val Leu Arg Tyr Val Glu Val Phe Lys Asn Ser Asp Lys Glu Asp Asp 645 650 655	2326
Cag gag cat cca tct gag aaa cag ccc tct ggg gct gag agt ggg act Gln Glu His Pro Ser Glu Lys Gln Pro Ser Gly Ala Glu Ser Gly Thr 660 665 670 675	2374
Cta gcc aga gcc tct ttg gct ctt cca act tcc tcc ctg tcc cgg acc Leu Ala Arg Ala Ser Leu Ala Leu Pro Thr Ser Ser Leu Ser Arg Thr 680 685 690	2422
gcg tcc cag agc agc agt cac cga ggc tgg gag atc ctt cgt caa aac Ala Ser Gln Ser Ser His Arg Gly Trp Glu Ile Leu Arg Gln Asn 695 700 705	2470
acc ctg ggg cac ttg aat ctt gga ctg aac ctt agt gag ggg gat gga Thr Leu Gly His Leu Asn Leu Gly Leu Asn Leu Ser Glu Gly Asp Gly 710 715 720	2518
gag gag gtc tac cat ttt tgattaacat cgctatcact cttgacctta Glu Glu Val Tyr His Phe 725	2566
ctcccggttg gcctgggggc ggggacagag acggagacct ctgcctatgc aagtgtctaa	2626
cttctgtgcc tgttaatcat gggagggtga gacagaacaa tccctaaagg gtcatgcctc	2686
acacttcaca tcagaatttc tggcaatggg caatggtcat cgattgtctc acgtattttc	2746
tgggctcttg caagtcaccc atctcaggaa aaaggaggtt ggcaactaaa gacatgaggc	2806
agggatgcta gattaatgtc aggacccatt tetettetge eecacgcage eectagaaag	2866
tagtaagetg tgaggetatt etggeteece agggettaeg tgggaagage caggeatgge	2926
atagaggttg tggcccttct ttttttc	2953
<210> 101 <211> 729 <212> PRT <213> Homo sapiens	
<400> 101	
Met Gly Gly Phe Leu Pro Lys Ala Glu Gly Pro Gly Ser Gln Leu Gln 1 5 10 15	
Lys Leu Leu Pro Ser Phe Leu Val Arg Glu Gln Asp Trp Asp Gln His 20 25 30	
Leu Asp Lys Leu His Met Leu Gln Gln Lys Arg Ile Leu Glu Ser'Pro 35 40 45	
Leu Leu Arg Ala Ser Lys Glu Asn Asp Leu Ser Val Leu Arg Gln Leu 50 55 60	

Leu Leu Asp Cys Thr Cys Asp Val Arg Gln Arg Gly Ala Leu Gly Glu 65 70 75 80

Thr Ala Leu His Ile Ala Ala Leu Tyr Asp Asn Leu Glu Ala Ala Leu 85 90 95

Val Leu Met Glu Ala Ala Pro Glu Leu Val Phe Glu Pro Thr Thr Cys 100 105 110

Glu Ala Phe Ala Gly Gln Thr Ala Leu His Ile Ala Val Val Asn Gln 115 120 125

Asn Val Asn Leu Val Arg Ala Leu Leu Thr Arg Arg Ala Ser Val Ser 130 135 140

Ala Arg Ala Thr Gly Thr Ala Phe Arg Arg Ser Pro Arg Asn Leu Ile 145 150 150 155 160

Tyr Phe Gly Glu His Pro Leu Ser Phe Ala Ala Cys Val Asn Ser Glu 165 170 175

Glu Ile Val Arg Leu Leu Ile Glu His Gly Ala Asp Ile Arg Ala Gln 180 185 190

Asp Ser Leu Gly Asn Thr Val Leu His Ile Leu Ile Leu Gln Pro Asn 195 200 205

Lys Thr Phe Ala Cys Gln Met Tyr Asn Leu Leu Leu Ser Tyr Asp Gly 210 215 220

His Gly Asp His Leu Gln Pro Leu Asp Leu Val Pro Asn His Gln Gly 225 230 235 240

Leu Thr Pro Phe Lys Leu Ala Gly Val Glu Gly Asn Thr Val Met Phe 245 250 255

Gln His Leu Met Gln Lys Arg Arg His Ile Gln Trp Thr Tyr Gly Pro $260 \hspace{1cm} 265 \hspace{1cm} 270$

Leu Thr Ser Ile Leu Tyr Asp Leu Thr Glu Ile Asp Ser Trp Gly Glu 275 280 285

Glu Leu Ser Phe Leu Glu Leu Val Val Ser Ser Asp Lys Arg Glu Ala 290 295 300

Arg Gln Ile Leu Glu Gln Thr Pro Val Lys Glu Leu Val Ser Phe Lys 305 310 315 320

Trp Asn Lys Tyr Gly Arg Pro Tyr Phe Cys Ile Leu Ala Ala Leu Tyr 325 330 335

Leu Leu Tyr Met Ile Cys Phe Thr Thr Cys Cys Val Tyr Arg Pro Leu 340 345 350

Lys Phe Arg Gly Gly Asm Arg Thr His Ser Arg Asp Ile Thr Ile Leu 355 360 365

Gln Gln Lys Leu Leu Gln Glu Ala Tyr Glu Thr Arg Glu Asp Ile Ile 370 375 380

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Arg Leu Val Gly Glu Leu Val Ser Ile Val Gly Ala Val Ile Ile Leu Leu Leu Glu Ile Pro Asp Ile Phe Arg Val Gly Ala Ser Arg Tyr Phe 405 410 415Gly Lys Thr Ile Leu Gly Gly Pro Phe His Val Ile Ile Ile Thr Tyr 420 425 430Ala Ser Leu Val Leu Val Thr Met Val Met Arg Leu Thr Asn Thr Asn 435 440 445 Gly Glu Val Val Pro Met Ser Phe Ala Leu Val Leu Gly Trp Cys Ser 450 455 460 Val Met Tyr Phe Thr Arg Gly Phe Gln Met Leu Gly Pro Phe Thr Ile 465 470 480 Met Ile Gln Lys Met Ile Phe Gly Asp Leu Met Arg Phe Cys Trp Leu 485 490 495 Gln Thr Glu Asp Pro Thr Ser Leu Gly Gln Phe Tyr Asp Tyr Pro Met 515 520 525Ala Leu Phe Thr Thr Phe Glu Leu Phe Leu Thr Val Ile Asp Ala Pro 530 535 540 Ala Asn Tyr Asp Val Asp Leu Pro Phe Met Phe Ser Ile Val Asn Phe 545 550 560 Ala Phe Ala Ile Ile Ala Thr Leu Leu Met Leu Asn Leu Phe Ile Ala 565 570 575 Met Met Gly Asp Thr His Trp Arg Val Ala Gln Glu Arg Asp Glu Leu 580 590 Trp Arg Ala Gln Val Val Ala Thr Thr Val Met Leu Glu Arg Lys Leu 595 600 605Pro Arg Cys Leu Trp Pro Arg Ser Gly Ile Cys Gly Cys Glu Phe Gly 610 615 620 Leu Gly Asp Arg Trp Phe Leu Arg Val Glu Asn His Asn Asp Gln Asn Pro Leu Arg Val Leu Arg Tyr Val Glu Val Phe Lys Asn Ser Asp Lys

Ser Gly Thr Leu Ala Arg Ala Ser Leu Ala Leu Pro Thr Ser Ser Leu 675 680 685

Glu Asp Asp Gln Glu His Pro Ser Glu Lys Gln Pro Ser Gly Ala Glu

Ser Arg Thr Ala Ser Gln Ser Ser Ser His Arg Gly Trp Glu Ile Leu Page 84

60

120

16U 200 PCT FINAL.ST25 695 700

Arg Gln Asn Thr Leu Gly His Leu Asn Leu Gly Leu Asn Leu Ser Glu 705 710 715 720

Gly Asp Gly Glu Glu Val Tyr His Phe

<210> 102 <211> 1545 <212> DNA <213> Homo sapiens

690

<220> <221> CDS <222> (130)..(438)

<223>

<400> 102
atttgtgatg ggcactggct cctggctgag gaccgcctct tcgggctctg gcacttctgc
accaccacca accagacgat ctgcttcaga gacctgggcc aggcccatgt gcccgggctg

gccgtgggc atg ggc ctg gta cgc agc gtg ggc gcc ttg gcc gtg gtg gcc 171

Met Gly Leu Val Arg Ser Val Gly Ala Leu Ala Val Val Ala

1 5 10

gcc att ttt ggc ctg gag ttc ctc atg gtg tcc cag ttg tgc gag gac 219
Ala Ile Phe Gly Leu Glu Phe Leu Met Val Ser Gln Leu Cys Glu Asp
15 20 25 30

aaa cac tca cag tgc aag tgg gtc atg ggt tcc atc ctc ctc ctg gtg
Lys His Ser Gln Cys Lys Trp Val Met Gly Ser Ile Leu Leu Leu Val
35 40 45

tct ttc gtc ctc tcc tcc ggc ggg ctc ctg ggt ttt gtg atc ctc ctc 315 Ser Phe Val Leu Ser Ser Gly Gly Leu Leu Gly Phe Val Ile Leu Leu 50 60

agg aac caa gtc aca ctc atc ggc ttc acc cta atg ttt tgg tgc gaa 363 Arg Asn Gln Val Thr Leu Ile Gly Phe Thr Leu Met Phe Trp Cys Glu 65 70 75

ttc act gcc tcc ttc ctc ctc ttc ctg aac gcc atc agc ggc ctt cac

Phe Thr Ala Ser Phe Leu Leu Phe Leu Asn Ala Ile Ser Gly Leu His
80
85

atc aac agc atc acc cat ccc tgg gaa tgaccgtgga aattttaggc 458

Ile Asn Ser Ile Thr His Pro Trp Glu

cccctccagg gacatcagat tccacaagaa aatatggtca aaatgggact tttccagcat 518 gtggcctctg gtggggctgg gttggacaag ggccttgaaa cggctgcctg tttgccgata 578

acttgtgggt ggtcagccag aaatggcccg ggggcctctg cacctggtct gcagggccag 638 aggccaggag ggtgcctcag tgccaccaac tgcacaggct tagccagatg ttgattttag 698

aggaagaaaa aaacatttta aaacteette ttgaatttte tteeetggae tggaatacag 758

ttggaagcac aggggtaact ggtacctgag ctagctgcac agccaaggat agttcatgcc 818 tgtttcattg acacgtgctg ggataggggc tgcagaatcc ctggggctcc cagggttgtt 878

aagaatggat cattetteea getaagggte caateagtge etaggaettt ettecaceag 938

ctcaaagggc cttcgtatgt atgtccctgg cttcagcttt ggtcatgcca aagaggcaga 998

gttcaggatt ccctcagaat gccctgcaca cagtaggttt ccaaaccatt tgactcggtt 1058

tgcctccctg cccgttgttt aaaccttaca aaccctggat aaccccatct tctagcagct 1118

ggctgtgcct ctgggagctc tgcctatcag aaccctacct taaggtgggt ttccttccga 1178

gaagagttet tgageaaget eteccaggag ggeeeacetg actgetaata cacageeete 1238

16U 200 PCT FINAL.ST25	
cccaaggccc gtgtgtgcat gtgtctgtct tttgtgaggg ttagacagcc tcagggcacc	1298
atttttaatc ccagaacaca tttcaaagag cacgtatcta gacctgctgg actctgcagg	1358
gggtgagggg gaacagcgag agcttgggta atgattaaca cccatgctgg ggatgcatgg	1418
aggtgaaggg ggccaggaac cagtggagat ttccatcctt gccagcacgt ctgtacttct	1478
gttcattaaa gtgctccctt tctagtcgat gtgtcactgc tgtatcatac ttttatgcta	1538
cacaacc	1545
<210> 103 <211> 103 <212> PRT <213> Homo sapiens <400> 103 Met Gly Leu Val Arg Ser Val Gly Ala Leu Ala Val Val Ala Ala Ile	
1 5 10 15	
Phe Gly Leu Glu Phe Leu Met Val Ser Gln Leu Cys Glu Asp Lys His 20 25 30	
Ser Gln Cys Lys Trp Val Met Gly Ser Ile Leu Leu Val Ser Phe 35 40 45	
Val Leu Ser Ser Gly Gly Leu Leu Gly Phe Val Ile Leu Leu Arg Asn 50 55 60	
Gln Val Thr Leu Ile Gly Phe Thr Leu Met Phe Trp Cys Glu Phe Thr 65 70 75 80	
Ala Ser Phe Leu Leu Phe Leu Asn Ala Ile Ser Gly Leu His Ile Asn 85 90 95	
Ser Ile Thr His Pro Trp Glu 100	
<210> 104 <211> 24 <212> DNA <213> Homo sapiens	
<400> 104 gcgcttccgg acctgtatct ccac	24
<210> 105 <211> 24 <212> DNA <213> Homo sapiens	
<400> 105 caagetetgg gtetegggea gaag	24
<210> 106 <211> 50 <212> DNA <213> Homo sapiens	÷
<400> 106 aaagagcete taaagaaggg ttecagaeta eeaggagete aetggaaata	50
<210> 107 <211> 24 <212> DNA	

16U 200 PCT FINAL.ST25 <213> Homo sapiens <400> 107 accatectge aaacttggat ggge 24 <210> 108 <211> 24 <212> DNA <213> Homo sapiens

aaggagccgg aagacaggga gagg 24 <210> 109

<211> 50 <212> DNA <213> Homo sapiens <400> 109 gctttatgta tatgaaaacc ctgtttatct gagcctagaa ctgtctttgc 50

<210> 110 <211> 50 <212> DNA

<400> 110 agtgatagtt ttaaatggga gggaataaag tctgcaaaat ttccccatat 50

<210> 111 <211> 24 <212> DNA <213> Homo sapiens <400> 111 gagtctccct gtgcgtttgg gctg 24

<210> 112 <211> 24 <212> DNA

<213> Homo sapiens <400> 112 aagtgtaaag catgccccgc ctga 24

<210> 113 <211> 50 <212> DNA <213> Homo sapiens <400> 113

agtcccagct taaaaaagag acagacagac agagagagag agagacagag 50

<210> 114 <211> 50 <212> DNA <213> Homo sapiens <400> 114

ttagtgattt aaaaaaatgt gaagaagaga gagtcaaggc agtaaaagga 50

<210> 115 <211> 24 <212> DNA <213> Homo sapiens <400> 115 gttcgctatg ctgccacggt catc 24

<210> 116

<400> 108

<213> Homo sapiens

<211> <212> <213>	24 DNA Homo sapiens					
<400>	116					24
agtect	gca gtcctggcat	tgtg				24
<210> <211>	117 50					
<211>	DNA					
<213>	Homo sapiens					
<400>	117					
gatacaa	iata attaaaagcc	caggttaagg	taaatatatt	aaagaccaag		50
<210>	118					
<211>	50					
<212> <213>	DNA Homo sapiens					
<400>	118					
atctcad	gaa ttaaaaatgc	tgaggtggta	aattgttatc	aattctatgt		50
<210>	119					
<211>	24					
<212>	DNA					
<213>	Homo sapiens					
<400>	119					~ .
caggatt	acg cacaaacggc	atgg				24
<210>	120					
<211>	24					
<212>	DNA					
<213>	Homo sapiens					
<400>	120					
tgggagg	cag agatagcaga	gccc	• •			24
<210>	121					
<211>	50					
<212>	DNA					
<213>	Homo sapiens					
<400>	121					
ctagact	att taaaaaaacc	cctggcttgc	acagtggctc	aagcctgtaa		50
<210>	122					
<211>	24					
<212> <213>	DNA Homo sapiens					
<400×	122	•				
<400>	122	2200				24
crygree	tgg gcaccctgat	aagc				24
<210>	123					
<211>	24					
<212>	DNA					
<213>	Homo sapiens					
	123	_				٠.
cccaggt	ctg gttgcagtgc	tctc				24
<210>	124				r 1	
<211>	50				. :	
<212>	DNA					
<213>	Homo sapiens					
<400>	124					
agctgtc	ctc attaaaagtg	acctggagtg	agatggattc	ttctqcctat		50

<210>	125				
<211>	50				
<212>	DNA				
<213>	Homo sapiens				
<400>	125				
	cttc tgaaaaacgg gagtca	stat sassassast			
ccuarc	cere egandanegg gagica	cigi gggcaccard	. acyceegggt		50
<210>	126				
<211>	24				
<212>	DNA				
<213>	Homo sapiens				
<400>	126				
ctgagg	tgtc cctcccaage aggt				24
<010×	127				
<210>	127				
<211> <212>	24				
<213>	DNA Homo canions				
\213/	Homo sapiens				
<400>	127				
	cgag aagcactgga gatg				24
	-y-yyyya yaty				24
<210>	128				
<211>	50				
<212>	DNA				
<213>	Homo sapiens				
<400>	128				
taaaca	aata cataaatgag gcagtt	acta gtagtggtaa	ctgctaggaa		50
<210>	129				
<211>	50				
<211>	DNA				
<213>	Homo sapiens				
	nome suprems				
<400>	129				
	aata taaaaatcag ccaggo	ctoo toocacatot	ctgtaatctc		50
	333-				
<210>	130				
<211>	50				
<212>	DNA				
<213>	Homo sapiens				
<400>	130				
gggatg	catt ataaatgcaa ccagcc	agag ggcccctggc	ttcagaacct		50
<210>	131				
<211>	27				
<212>					
<213>	Homo sapiens				
	Tomo Daprens				
<400>	131				
gtcacci	cag cgatctcaac gatagg	1			27
		•			**
<210>	132				
<211>	28				
<212>	DNA				
<213>	Homo sapiens				
- 400-	120				
<400>	132				
cygagca	igga acaggatata ggtcago	ıg		r 1	28
<210>	133			. •	
<211>	50				
<212>	DNA				
<213>	Homo sapiens				
<400>	133				

atatacc	ttg tttaaaagag	16U 200 PCT FINAL.ST25 gggtattatc acaataaaac aaggaaagct	50
<211> <212>	134 50 DNA Homo sapiens		
<400>	134	tgacaaacag tgctaaagtt ctcaccttaa	50
<211> <212>	135 24 DNA Homo sapiens		
<400>	135 _{Jaag} gaagcaggga	agag	24
<210><211><211><212><212><213>	136 24 DNA Homo sapiens		
<400> ccagcta	136 agtt catgettgge	gcag	24
<210><211><211><212>	137 50 DNA		
<213> <400> ttattge	Homo sapiens 137 ggca taaaaatatg	aagagaggtc ccagagagtc cctaggttct	50
<210><211><211><212><213>	138 30 DNA Homo sapiens		
<400> ctgttg	138 ggaa tottoagooa	gateteacae	30
<210><211><211><212><213>	139 24 DNA Homo sapiens		
<400> atggag	139 gttt ctgcacgcto	c agca	24
<210> <211> <212> <213>	140 50 DNA Homo sapiens		
<400> aagcaa	140 tttg ttaaaaacto	g gcattacttt actcttatgc tttctgtgtc	50
<210> <211> <212> <213>	141 50 DNA Homo sapiens		
<400> acttta	141 aattt tataaagaa	g gttcacatca agaaattcca agtgaggttc	50
<210><211><211><212><213>	142 24 DNA Homo sapiens		

Page 90

<400> gggcca	142 cttc cacagacagg aagc	24
<210><211><211><212><213>		
<400> tggcct	143 gaga ggtagattcc acatagtagt cgt	33
<210><211><211><212>	144 50 DNA	
<213> <400> aaggct	•	50
<210><211><211><212>	145 31 DNA	
<213> <400>	Homo sapiens	2.
<210>	tctg gtcagagaac aagactggga c	31
<211> <212> <213>	27 DNA Homo sapiens	
<400> gtgatg	146 toto gagaatgagt goggttg	27
<210><211><211><212><213>	147 50 DNA Homo sapiens	
<400> cagcgag	147 ggca gaaaaatgte ceacaagttg ageeeteece acteecagtg	50
<210><211><211><212><212><213>	148 50 DNA Homo sapiens	
<400> taatata	148 aaaa tatataaaat agtgcaacat tacttattcc tcctggtgtt	50
<210> <211> <212>	149 27 DNA	
<213> <400> gcagato	Homo sapiens 149 gacc cgacctgact gttcttc	27
<210> <211>	150 27	
<212> <213> <400>	DNA Homo sapiens	
	150 gca gctagctcag gtaccag	27
<210>	151 50	

Page 91

<212		DNA								1 6 U	200	PCT	FINA	L.ST	25	
<213		Homo	sapı	ens												
<400 gcca		151 agt 1	taaa	atgaa	eg co	cctac	ttt	9 999	gcago	gagc	999	agga	aac			50
<210 <211 <212 <213	.> ?>	152 945 DNA Homo	sapi	iens												
<220 <221 <222 <223	.> !>	CDS (1).	. (945	5)												
	ggt	152 gta Val														48
		gaa Glu														96
		tac Tyr 35														144
		ctg Leu														192
		tct Ser														240
		tca Ser														288
		att Ile														336
		ctg Leu 115														384
cca Pro	ctg Leu 130	ctc Leu	tac Tyr	agg Arg	gtc Val	atc Ile 135	atg Met	t <i>cc</i> Ser	cct Pro	agg Arg	gtc Val 140	tgt Cys	tct Ser	ctg Leu	ctg Leu	432
gtg Val 145	gct Ala	gct Ala	gtc Val	ttc Phe	tca Ser 150	gta Val	ggt Gly	ttc Phe	act Thr	gat Asp 155	gct Ala	gtg Val	atc Ile	cat His	gga Gly 160	480
		ata Ile														528
		tgt Cys														576
		gat Asp 195														624
		agc Ser										Ile				672
atc Ile 225	ctg Leu	cgc Arg	atc Ile	cac His	tct Ser 230	aaa Lys	aag Lys	ggc Gly	agg Arg	tgc Cys 235	aaa Lys	gcg Ala	ttt Phe	agc Ser	acc Thr 240	720
tgt	ago	tcc	cac	ctg	aca	gct	gtt	ctt	atg	ttt				ctg	atg	768
												Page	92			

Cys Ser	Ser	His	Leu 245	Thr	Ala	Val	Leu	Met 250				FINA Ser				
tcc atg Ser Met	tat Tyr	ctc Leu 260	aaa Lys	cct Pro	gct Ala	tct Ser	agc Ser 265	agt Ser	tca Ser	ctc Leu	acc Thr	cag Gln 270	gag Glu	aaa Lys		816
gta tcc Val Ser	tca Ser 275	gta Val	ttt Phe	tat Tyr	acc Thr	act Thr 280	gtg Val	att Ile	ctc Leu	atg Met	ttg Leu 285	aat Asn	ccc Pro	ttg Leu		864
ata tat Ile Tyr 290	Ser	ctg Leu	agg Arg	aac Asn	aat Asn 295	gaa Glu	gta Val	aga Arg	aat Asn	gct Ala 300	ctg Leu	atg Met	aaa Lys	ctt Leu		912
tta aga Leu Arg 305	aga Arg	aaa Lys	ata Ile	tct Ser 310	tta Leu	tct Ser	cca Pro	gga Gly	taa							945
<210><211><211><212><213>	153 314 PRT Homo	sap:	iens													
<400>	153	_	_		.	m	**- 1	m>	63.	Dh o	Lou	Lou	Sor	Gly		
Met Gly	y Val	Lys	Asn 5	His	Ser	Thr	val	10	GIU	ьие	reu	ren	15	GIY		
Leu Th	Glu	Gln 20	Ala	Glu	Leu	Gln	Leu 25	Pro	Leu	Phe	Cys	Leu 30	Phe	Leu		
Gly Ile	Tyr 35	Thr	Val	Thr	Val	Val 40	Gly	Asn	Leu	Ser	Met 45	Ile	Ser	Ile		
Ile Are	g Leu	Asn	Arg	Gln	Leu 55	His	Thr	Pro	Met	Tyr 60	Tyr	Phe	Leu	Ser		
Ser Le	ı Ser	Phe	Leu	Asp 70	Phe	Cys	Tyr	Ser	Ser 75	Val	Ile	Thr	Pro	Lys 80	•	
Met Le	u Ser	Gly	Phe 85	Leu	Cys	Arg	Asp	Arg 90	Ser	Ile	Ser	Tyr	Ser 95	Gly		
Cys Me	t Ile	Gln 100		Phe	Phe	Phe	Cys 105		Cys	Val	Ile	Ser 110	Glu	Cys		
Tyr Me	t Leu 115		Ala	Met	Ala	Cys 120		Arg	Tyr	Val	Ala 125	Ile	Cys	Ser		
Pro Le 13		Tyr	Arg	Val	11e 135		Ser	Pro	Arg	Val 140	. Cys	Ser	Leu	Leu	-	
Val Al 145	a Ala	Val	Phe	Ser 150		Gly	Phe	Thr	Asp 155	Ala	ı Val	Ile	His	Gly 160		
Gly Cy	s Ile	e Leu	165		Ser	Phe	Cys	Gly 170	Ser	Asr	ılle	: Ile	Lys 175	His		
Tyr Ph	e Cys	180		· Val	Pro	Leu	11e 185		Leu	ı Sei	Cys	3 Ser 190	Ser	Thr		
Tyr Il	e Asp 195		ı Leu	Leu	Ile	200		. Ile	: Gl	/ Gl	7 Phe 205	e Asn	Met	: Val		

16U 200 PCT FINAL.ST25 Ala Thr Ser Leu Thr Ile Ile Ile Ser Tyr Ala Phe Ile Leu Thr Ser 210 215 220	
Ile Leu Arg Ile His Ser Lys Lys Gly Arg Cys Lys Ala Phe Ser Thr 225 230 235 240	
Cys Ser Ser His Leu Thr Ala Val Leu Met Phe Tyr Gly Ser Leu Met 245 250 255	
Ser Met Tyr Leu Lys Pro Ala Ser Ser Ser Ser Leu Thr Gln Glu Lys 260 265 270	
Val Ser Ser Val Phe Tyr Thr Thr Val Ile Leu Met Leu Asn Pro Leu 275 280 285	
Ile Tyr Ser Leu Arg Asn Asn Glu Val Arg Asn Ala Leu Met Lys Leu 290 295 300	
Leu Arg Arg Lys Ile Ser Leu Ser Pro Gly 305 310	
<210> 154 <211> 34 <212> DNA <213> Homo sapiens	
<400> 154 ctgtgatcca tggaggttgt atactcaggt tgtc	34
<210> 155 <211> 36 <212> DNA <213> Homo sapiens	
<400> 155 ' tcatcagage atttcttact tcattgttcc tcagac	36
<210> 156 <211> 50 <212> DNA <213> Homo sapiens	
<400> 156 caggagaatt aaatataaga gtggtcagtg tgtttgtaac actcaggaca	50
<210> 157 <211> 50 <212> DNA <213> Homo sapiens	
<400> 157 aaaacatgct ttaaaaaacc catgatatta aagacaaaaa actgagcata	50
<210> 158 <211> 50 <212> DNA <213> Homo sapiens	
<400> 158 atgaacagct tattaaatag ccaggtagct gggcagaatg agaaaatgca	∵ 50
<pre><210> 159 <211> 50 <212> DNA <213> Homo sapiens</pre>	
<400> 159	

16U 200 PCT FINAL.ST25 gcccaacact aaataaaggg tcagctttct cagagataag gccatgattg	50										
<210> 160 <211> 50 <212> DNA <213> Homo sapiens											
<400> 160 tgctataaaa tgtttttaaa aagtgtgaag ttggcctatc accaagtaag	50										
<210> 161 <211> 50 <212> DNA <213> Homo sapiens											
<400> 161 taaatattgt atttatatag teetteagga ggaetgagge ateeteeagt											
<210> 162 <211> 957 <212> DNA <213> Homo sapiens											
<220> <221> CDS <222> (1)(957) <223>											
<400> 162 atg aat cca gca aat cat tcc cag gtg gca gga ttt gtt cta ctg ggg Met Asn Pro Ala Asn His Ser Gln Val Ala Gly Phe Val Leu Leu Gly 1 5 10 15	48										
ctc tct cag gtt tgg gag ctt cgg ttt gtt ttc ttc act gtt ttc tct Leu Ser Gln Val Trp Glu Leu Arg Phe Val Phe Phe Thr Val Phe Ser 20 25 30	96										
gct gtg tat ttt atg act gta gtg gga aac ctt ctt att gtg gtc ata Ala Val Tyr Phe Met Thr Val Val Gly Asn Leu Leu Ile Val Val Ile 35 40 45	144										
gtg acc tcc gac cca cac ctg cac aca acc atg tat ttt ctc ttg ggc Val Thr Ser Asp Pro His Leu His Thr Thr Met Tyr Phe Leu Leu Gly 50 55 60	192										
aat ctt tct ttc ctg gac ttt tgc tac tct tcc atc aca gca cct agg Asn Leu Ser Phe Leu Asp Phe Cys Tyr Ser Ser Ile Thr Ala Pro Arg 65 70 75 80	240										
atg ctg gtt gac ttg ctc tca ggc aac cct acc att tcc ttt ggt gga Met Leu Val Asp Leu Leu Ser Gly Asn Pro Thr Ile Ser Phe Gly Gly 85 90 95	288										
tgc ctg act caa ctc ttc ttc ttc cac ttc att gga ggc atc aag atc Cys Leu Thr Gln Leu Phe Phe His Phe Ile Gly Gly Ile Lys Ile 100 105 110	336										
ttc ctg ctg act gtc atg gcg tat gac cgc tac att gcc att tcc cag Phe Leu Leu Thr Val Met Ala Tyr Asp Arg Tyr Ile Ala Ile Ser Gln 115 120 125	384										
ccc ctg cac tac acg ctc att atg aat cag act gtc tgt gca ctc ctt Pro Leu His Tyr Thr Leu Ile Met Asn Gln Thr Val Cys Ala Leu Leu 130 135 140	432										
atg gca gcc tcc tgg gtg ggg ggc ttc atc cac tcc ata gta cag att Met Ala Ala Ser Trp Val Gly Gly Phe Ile His Ser Ile Val Gln Ile 145 150 155 160	480										
gca ttg act atc cag ctg cca ttc tgt ggg cct gac aag ctg gac aac Ala Leu Thr Ile Gln Leu Pro Phe Cys Gly Pro Asp Lys Leu Asp Asn 165 170 175	528										
ttt tat tgt gat gtg cct cag ctg atc aaa ttg gcc tgc aca gat acc Phe Tyr Cys Asp Val Pro Gln Leu Ile Lys Leu Ala Cys Thr Asp Thr 180 185 190	576										

160	200	PCT	FINAL	ST25

ttt Phe	gtc Val	tta Leu 195	gag Glu	ctt Leu	tta Leu	atg Met	gtg Val 200	tct Ser	aac Asn	aat Asn	ggc Gly	ctg Leu 205	gtg Val	acc Thr	ctg Leu	624
atg Met	tgt Cys 210	Phe	ctg Leu	gtg Val	ctt Leu	ctg Leu 215	gga Gly	tcg Ser	tac Tyr	aca Thr	gca Ala 220	ctg Leu	cta Leu	gtc Val	atg Met	672
ctc Leu 225	Arg	agc Ser	cac His	tca Ser	cgg Arg 230	gag Glu	ggc Gly	cgc Arg	agc Ser	aag Lys 235	gcc Ala	ctg Leu	tct Ser	acc Thr	tgt Cys 240	720
gcc Ala	tct Ser	cac His	att Ile	gct Ala 245	gtg Val	gtg Val	acc Thr	tta Leu	atc Ile 250	ttt Phe	gtg Val	cct Pro	tgc Cys	atc Ile 255	tac Tyr	768
									ccc Pro							816
									ctg Leu							864
		Asn							atg Met							912
	Lys								cac His					tag		957
<210 <211 <211 <211	1 > 2 >	163 318 PRT Homo	sapi	.ens												
< 400	0>	163														
Met 1	Asn	Pro	Ala	Asn 5	His	Ser	Gln	Val	Ala 10	Gly	Phe	Val	Leu	Leu 15	Gly	
Leu	Ser	Gln	Val 20	Тгр	Glu	Leu	Arg	Phe 25	Val	Phe	Phe	Thr	Val 30	Phe	Ser	
Ala	Val	Tyr 35	Phe	Met	Thr	Val	Val 40	Gly	Asn	Leu	Leu	11e 45	Val	Val	Ile	
Val	Thr 50	Ser	Asp	Dro												
Aen				F10	His	Leu 55	His	Thr	Thr	Met	Tyr 60	Phe	Leu	Leu	Gly	
65	Leu	Ser				55			Thr		60					
65			Phe	Leu	Asp 70	55 Phe	Cys	Туг		Ser 75	60 Ile	Thr	Ala	Pro	Arg 80	
65 Met	Leu	Val	Phe Asp	Leu Leu 85	Asp 70 Leu	55 Phe Ser	Cys Gly	Tyr Asn	Ser Pro	Ser 75 Thr	60 Ile Ile	Thr Ser	Ala Phe	Pro Gly 95	Arg 80 Gly	
65 Met Cys	Leu Leu	Val Thr	Phe Asp Gln 100	Leu Leu 85 Leu	Asp 70 Leu Phe	55 Phe Ser Phe	Cys Gly Phe	Tyr Asn His 105	Ser Pro 90	Ser 75 Thr	<pre>11e Ile Gly</pre>	Thr Ser Gly	Ala Phe Ile 110	Pro Gly 95 Lys	Arg 80 Gly Ile	
65 Met Cys Phe	Leu Leu Leu	Val Thr Leu 115	Phe Asp Gln 100	Leu 85 Leu Val	Asp 70 Leu Phe	Phe Ser Phe	Cys Gly Phe Tyr 120	Tyr Asn His 105 Asp	Ser Pro 90 Phe	Ser 75 Thr Ile	<pre>11e Ile Gly Ile</pre>	Thr Ser Gly Ala 125	Ala Phe Ile 110	Gly 95 Lys Ser	Arg 80 Gly Ile	

Ala	Leu	Thr	Ile	Gln 165	Leu	Pro	Phe	Cys	Gly 170	Pro	Asp	Lys	Leu	Asp 175	Asn		
Phe	Туr	Cys	Asp 180	Val	Pro	Gln	Leu	Ile 185	Lys	Leu	Ala	Cys	Thr 190	Asp	Thr		
Phe	Val	Leu 195	Glu	Leu	Leu	Met	Val 200	Ser	Asn	Asn	Gly	Leu 205	Val	Thr	Leu		
Met	Cys 210		Leu	Val	Leu	Leu 215	G1 y	Ser	Tyr	Thr	Ala 220	Leu	Leu	Val	Met		
Leu 225	Arg	Ser	His	Ser	Arg 230	Glu	Gly	Arg	Ser	Lys 235	Ala	Leu	Ser	Thr	Cys 240		
Ala	Ser	His	Ile	Ala 245	Val	Val	Thr	Leu	11e 250	Phe	Val	Pro	Cys	Ile 255	Tyr		
Val	Туr	Thr	Arg 260	Pro	Phe	Arg	Thr	Phe 265	Pro	Met	Asp	Lys	Ala 270	Val	Ser		
Val	Leu	T yr 275	Thr	Ile	Val	Thr	Pro 280	Met	Leu	Asn	Pro	Ala 285	Ile	Tyr	Thr		
Leu	Arg 290	Asn	Lys	Glu	Val	Ile 295	Met	Ala	Met	Lys	Lys 300	Leu	Trp	Arg	Arg		
Lys 305	Lys	Asp	Pro	Ile	Gly 310	Pro	Leu	Glu	His	Arg 315	Pro	Leu	His				
<210 <211 <212 <213	l> : ?> !	164 26 DNA Homo	sap:	iens													
<400 gaat		164 gca a	aatca	attc	cc a	ggtgg	9										26
<210 <211 <212 <213	l> : !> !	165 29 DNA Homo	san	iane													
<400)> :	165 aag (ct co	agg	ggac										29
<210 <211 <212 <213	! > ! !> !	166 50 DNA Homo	sapi	iens													
<400)> :	166 ctt (a go	gggg	cagto	g ati	ctctt	ttc	tttt	ctt	tt				50
<210 <211 <212 <213	> ! >	167 972 ONA Homo	sapi	iens												.*	
<220 <221 <222 <223	> (!>	DS	. (972	2)													

Page 97

< 40	0 > -	167								160	200	PCT	FINA	AL.SI	r25	
atg	aac	cct										gtc Val				48
												ctg Leu				96
												att Ile 45				144
												ttc Phe				192
												gtg Val				240
												tca Ser				288
												acc Thr				336
ttc Phe	ctc Leu	ttg Leu 115	gcc Ala	gtc Val	atg Met	tct Ser	ctg Leu 120	gat Asp	cgt Arg	tac Tyr	ctg Leu	gca Ala 125	atc Ile	tgc Cys	cga Arg	384
												tgt Cys				432
												ctt Leu				480
												ggt Gly				528
												tgt Cys				576
												ttg Leu 205				624
												att Ile				672
												gcg Ala				720
												ggc Gly				768
												ctg Leu				816
												ttg Leu 285				864
												ctg Leu				912
												agg Arg				960

305 310 16U 200 PCT FINAL.ST25 315 32

caa agg aaa tga Gln Arg Lys 972

<210> 168 <211> 323 <212> PRT

<213> Homo sapiens

<400> 168

Met Asn Pro Glu Asn Trp Thr Gln Val Thr Ser Phe Val Leu Leu Gly $1 \hspace{1cm} 5 \hspace{1cm} 10 \hspace{1cm} 15$

Phe Pro Ser Ser His Leu Ile Gln Phe Leu Val Phe Leu Gly Leu Met 20 25 30

Val Thr Tyr lle Val Thr Ala Thr Gly Lys Leu Leu Ile Ile Val Leu 35 40

Ser Trp Ile Asp Gln Arg Leu His Ile Gln Met Tyr Phe Phe Leu Arg 50 60

Asn Phe Ser Phe Leu Glu Leu Leu Leu Val Thr Val Val Val Pro Lys 65 70 75 80

Met Leu Val Val Ile Leu Thr Gly Asp His Thr Ile Ser Phe Val Ser 85 90 95

Cys Ile Ile Gln Ser Tyr Leu Tyr Phe Phe Leu Gly Thr Thr Asp Phe 100 105 110

Phe Leu Leu Ala Val Met Ser Leu Asp Arg Tyr Leu Ala Ile Cys Arg 115 120 125

Pro Leu Arg Tyr Glu Thr Leu Met Asn Gly His Val Cys Ser Gln Leu 130 135 140

Val Leu Ala Ser Trp Leu Ala Gly Phe Leu Trp Val Leu Cys Pro Thr 145 150150155

Val Leu Met Ala Ser Leu Pro Phe Cys Gly Pro Asn Gly Ile Asp His 165 170 175

Phe Phe Arg Asp Ser Trp Pro Leu Leu Arg Leu Ser Cys Gly Asp Thr 180 185 190

His Leu Leu Lys Leu Val Ala Phe Met Leu Ser Thr Leu Val Leu Leu 195 200 205

Gly Ser Leu Ala Leu Thr Ser Val Ser Tyr Ala Cys Ile Leu Ala Thr 210 215 220

Val Leu Arg Ala Pro Thr Ala Ala Glu Arg Arg Lys Ala Phe Ser Thr 225 230 235 240

Cys Ala Ser His Leu Thr Val Val Val Ile Ile Tyr Gly Ser Ser Ile 245 250 255

Phe Leu Tyr Ile Arg Met Ser Glu Ala Gln Ser Lys Leu Leu Asn Lys Page 99

City Ala Ser Val Leu Ser Cys lie lie Thr Pro Leu Leu Asn Pro Phe 285 The Phe Thr Leu Arg Asn Asp Lys Val Gln Gln Ala Leu Arg Glu Ala 290 m of the Color of the C		260		265	160	200 PCT	FINAL.ST	r25
Leu Gly Trp Pro Arg Leu Thr Ala Val Met Lys Leu Arg Val Thr Ser 305 Gln Arg Lys C210> 169 C211> 25 C212> BNA C213> Homo sapiens C400> 169 tgtgctcagc tggatagacc aacgc			Ser Cys		Thr Pro		Asn Pro	Phe
Gin Arg Lys Carry 169 Carry 211 25 25		Leu Arg		Lys Val	Gln Gln		Arg Glu	Ala
C210 169		Pro Arg		Ala Val			Val Thr	
<pre> <211> DIN <213> Homo sapiens 4400> 169 tgtgctcagc tggatagacc aacgc</pre>	Gln Arg Lys							
<pre>c210> 170 c211> 30 c212> DNA c213> Homo sapiens <400> 170 ctgagaacag tggcaagaat gcaggcatag</pre>	<211> 25 <212> DNA	sapiens						
<pre> c211> 30 c122> DNA c213> Homo sapiens <400> 170 ctgaqaacaq tggcaagaat gcaggcatag 210> 171 c211> 450 c212> DNA c213> Homo sapiens c220> c221> DNA c213> Homo sapiens c220> c221> CDS c222> (11)(450) c223> c222> (11)(450) c223> c400> 171 atg gac ctt ccc cat gtc cca gct ctg gac gcc cca ctc ttt gga gtc gcc c22> (11)(450) c223> c400> 171 atg gac ctt ccc cat gtc cca gct ctg gac gcc cca ctc ttt gga gtc gcc gcc gcc gcc gcc gcc gcc gcc gcc</pre>		tggataga	cc aacgc					25
ctgagaacag tggcaagaat gcaggcatag 30 <pre> <210> 171 <211> 450 <212> DNA <213> Homo sapiens </pre> <pre> <220> <222> (1) (450) </pre> <pre> <222> (1) (450) </pre> <pre> <222> (1) (450) </pre> <pre> <223> (1) (450) </pre> <pre> <2240</pre>	<211> 30 <212> DNA	sapiens						
<pre> <211> 450 <212> DNA <213> Homo sapiens </pre> <pre> <220> <221> CDS <222> (1) (450) <222> (1) (450) </pre> <pre> <400> 171 atg gac ctt ccc cat gtc cca gct ctg gac gcc cca ctc ttt gga gtc Met Asp Leu Pro His Val Pro Ala Leu Asp Ala Pro Leu Phe Gly Val 1</pre>		tggcaagaa	at gcaggo	catag				30
<pre> <221> CDS <222> (1)(450) <222></pre>	<211> 450 <212> DNA	sapiens						
Asp dec ctt ccc cat gtc cca gct ctg gac gcc cca ctc ttt gga gtc degree ctg level his val pro Ala Leu Asp Ala Pro Leu Phe Gly val 15 ttc ctg gtg gtt tat gtg ctt act gtg ctg ggg aac ctc ctc atc ctg degree ctg gtg gtg act act ctg leu act leu Thr val Leu Gly Asn Leu Leu lle Leu all leu all leu val lle Leu val lle Arg val Tyr Ser His Leu His Thr Pro Lys Tyr Tyr Phe act act gtg degree ccc aact gtc atg gtg tac ttc act gtg tgg ttc ttc act gtc atg gtg leu act ctg leu his Thr Pro Lys Tyr Tyr Phe act ctc acc aact ctg tcc acc act gtc atg gtg trouble leu Thr Asn Leu Ser Phe lle Asp Leu Trp Phe Phe Thr Val Met Val Ser leu Cys Gly Lys Ala val Ser leu Cys Gly Lys Ala val Ser acc acc act cac act gtc atg gtg leu ttc cac acc gtc act gtg ggg acc leu ctg trouble leu Cys Gly Lys Ala val Ser acc acc acc cac act ttc ttc leu ttc leu Ser Leu Cys Gly Lys Ala val Ser leu Cys ggg acc acc leu ctg ttg trouble leu Gly Ser go acc gtg ttc ttc cac acc acc ctc ttc tac acc gtc tat acc gtc gtg tcc leu cac acc acc ctc ttc tac acc gtc tat acc gtc acc leu cys ggg acc leu acc gac gtg ttc ttc cac acc gac gtc acc leu cys ggg acc leu acc ggt acc leu cys ggg acc leu acc gly ser glu leu leu Tyr Thr val Met Ser Tyr Asp Arg Tyr Arg Gly lilo acc cac act cac cac acc ccc cac gaa gtg acc leu leu cys lilo acc cac gac gtg acc acc cac ccc cac gaa gtg acc acc leu cys glu acc cac acc ccc cac gaa gtg acc acc leu cys glu acc cac acc ccc cac gaa gtg acc acc leu cac acc gac acc leu cac acc acc ccc cac gaa gtg acc acc leu cac acc ggt acc acc acc ccc cac gaa gtg acc acc leu cac acc acc ccc acc gaa gtg acc acc leu cac acc acc acc acc acc acc acc acc ac	<221> CDS <222> (1).	. (450)						
Phe Leu Val Val Tyr Val Leu Thr Val Leu Gly Asn Leu Leu Ile Leu 25 ctg gtg atc agg gtg tac tct cac ctc cac acc ccc aag tac tac ttc Leu Val Ile Arg Val Tyr Ser His Leu His Thr Pro Lys Tyr Tyr Phe 35 ctc acc aat ctg tcc ttc att gac ttg tgg ttc ttc act gtc atg gtg Leu Thr Asn Leu Ser Phe Ile Asp Leu Trp Phe Phe Thr Val Met Val 50 ccc aaa atg ccg agg acc ttg ttg tcc ctg tgt ggc aag gct gtg tcc Pro Lys Met Pro Arg Thr Leu Leu Ser Leu Cys Gly Lys Ala Val Ser 65 ctc acc agt tgt atg acc caa ctc tat ttc ttc tac ttc ctg ggg agc Ber Phe His Ser Cys Met Thr Gln Leu Tyr Phe Phe Tyr Phe Leu Gly Ser 95 acc gag tgt ttg ctc tac acg gtc atg tcc tat gat ccc tat gat cgc tat aga gga Thr Glu Cys Leu Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Arg Gly 110 aat act cag cac ttc cca ggt agt gaa aac act ccc cac gaa gtg agc Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser 384	atg gac ctt Met Asp Leu	Pro His			Asp Ala		Phe Gly	
Leu Val Ile Arg Val Tyr Ser His Leu His Thr Pro Lys Tyr Tyr Phe 45 ctc acc aat ctg tcc ttc att gac ttg tgg ttc ttc act gtc atg gtg 192 Leu Thr Asn Leu Ser Phe Ile Asp Leu Trp Phe Phe Thr Val Met Val 50 ccc aaa atg ccg agg acc ttg ttg tcc ctg tgt ggc aag gct gtg tcc Pro Lys Met Pro Arg Thr Leu Leu Ser Leu Cys Gly Lys Ala Val Ser 70 ttc Cac agt tgt atg acc caa ctc tat ttc ttc tac ttc ctg ggg agc Phe His Ser Cys Met Thr Gln Leu Tyr Phe Phe Tyr Phe Leu Gly Ser 95 acc gag tgt ttg ctc tac acg gtc atg tcc tat gat cgc tat aga gga 336 Thr Glu Cys Leu Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Arg Gly 110 aat act cag cac ttc cca ggt agt gaa aac act ccc cac gaa gtg agc 384 Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser		Val Tyr		Thr Val			Leu Ile	
Leu Thr Asn Leu Ser Phe Ile Asp Leu Trp Phe Phe Thr Val Met Val Ser Ccc aaa atg ccg agg acc ttg ttg tcc ctg tgt ggc aag gct gtg tcc Pro Lys Met Pro Arg Thr Leu Leu Ser Leu Cys Gly Lys Ala Val Ser 70	Leu Val Ile			His Leu		Pro Lys		
Pro Lys Met Pro Arg Thr Leu Leu Ser Leu Cys Gly Lys Ala Val Ser 80 ttc cac agt tgt atg acc caa ctc tat ttc ttc tac ttc ctg ggg agc Phe His Ser Cys Met Thr Gln Leu Tyr Phe Phe Tyr Phe Leu Gly Ser 95 acc gag tgt ttg ctc tac acg gtc atg tcc tat gat cgc tat aga gga 336 Thr Glu Cys Leu Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Arg Gly 100 aat act cag cac ttc cca ggt agt gaa aac act ccc cac gaa gtg agc Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser	Leu Thr Asn		Phe Ile			Phe Thr		
Phe His Ser Cys Met Thr Gln Leu Tyr Phe Phe Tyr Phe Leu Gly Ser 95 . acc gag tgt ttg ctc tac acg gtc atg tcc tat gat cgc tat aga gga Thr Glu Cys Leu Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Arg Gly 100 105 110 384 aat act cag cac ttc cca ggt agt gaa aac act ccc cac gaa gtg agc Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser	Pro Lys Met		Thr Leu		Leu Cys			Ser
Thr Glu Cys Leu Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Arg Gly 100 105 110 aat act cag cac ttc cca ggt agt gaa aac act ccc cac gaa gtg agc Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser		Cys Met			Phe Phe		Leu Gly	Ser
Asn Thr Gln His Phe Pro Gly Ser Glu Asn Thr Pro His Glu Val Ser		Leu Leu		Val Met			Tyr Arg	
	Asn Thr Gln			Ser Glu		Pro His		

Gln Met Le 130		a Alg	Gly 135	Ala	His	Gly	Leu	Pro 140	Leu	Ile	Ile	ctg Leu	432
gca gat ct Ala Asp Le 145													450
<210> 172 <211> 149 <212> PRT <213> Hom		s											
<400> 172													
Met Asp Le	u Pro Hi 5	s Val	Pro	Ala	Leu	Asp 10	Ala	Pro	Leu	Phe	Gly 15	Val	
Phe Leu Va	l Val Ту 20	r Val	Leu	Thr	Val 25	Leu	Gly	Asn	Leu	Leu 30	lle	Leu	
Leu Val II 35	e Arg Va	l Tyr	Ser	His 40	Leu	His	Thr	Pro	Lys 45	туr	Tyr	Phe	
Leu Thr As 50	n Leu Se	r Phe	Ile 55	Asp	Leu	Trp	Phe	Phe 60	Thr	Val	Met	Val	
Pro Lys Me 65	t Pro Ar	g Thr 70	Leu	Leu	Ser	Leu	Cys 75	Gly	Lys	Ala	Val	Ser 80	
Phe His Se	r Cys Me 85	t Thr	Gln	Leu	Tyr	Phe 90	Phe	Tyr	Phe	Leu	Gly 95	Ser	
Thr Glu Cy	s Leu Le 100	u Tyr	Thr	Val	Met 105	Ser	туг	Asp	Arg	Туг 110	Arg	Gly	
Asn Thr Gl	n His Ph 5	e Pro	Gly	Ser 120	Glu	Asn	Thr	Pro	His 125	Glu	Val	Ser	
Gln Met Le 130	val Al	a Arg	Gly 135	Ala	His	Gly	Leu	Pro 140	Leu	Ile	Ile	Leu	
Ala Asp Le 145	Ser Gl	У											
<210> 173 <211> 23 <212> DNA <213> Hom	o sapien	5											
<400> 173 agctctggac	gcccac	tct tt	:ġ										23
<210> 174 <211> 27 <212> DNA <213> Hom	sapien:	5											
<400> 174 acccactcag	atctgcc	agg at	gatg	a									27
<210> 175 <211> 936 <212> DNA <213> Home	o sapien:	;											

<22 <22 <22 <22	1> 2>	CDS (1).	. (93	6)												
<40 atg Met 1	tcc	175 aac Asn	gcc Ala	agc Ser 5	ctc Leu	gtg Val	aca Thr	gca Ala	ttc Phe 10	atc Ile	ctc Leu	aca Thr	ggc Gly	ctt Leu 15	ccc Pro	48
								ctc Leu 25								96
tac Tyr	gtg Val	ctc Leu 35	act Thr	gtg Val	ctg Leu	G] y	aac Asn 40	ctc Leu	ctc Leu	atc Ile	ctg Leu	ctg Leu 45	gtg Val	atc Ile	agg Arg	144
gtg Val	gat Asp 50	tct Ser	cac His	ctc Leu	cac His	acc Thr 55	ccc Pro	atg Met	tac Tyr	tac Tyr	ttc Phe 60	ctc Leu	acc Thr	aac Asn	ctg Leu	192
tcc Ser 65	ttc Phe	att Ile	gac Asp	atg Met	tgg Trp 70	ttc Phe	tcc Ser	act Thr	gtc Val	acg Thr 75	gtg Val	ccc Pro	aaa Lys	atg Met	ctg Leu 80	240
								agg Arg								288
								ttc Phe 105								336
ctc Leu	tac Tyr	aca Thr 115	gtc Val	atg Met	tcc Ser	tat Tyr	gat Asp 120	cgc Arg	tac Tyr	ttg Leu	gcc Ala	atc Ile 125	agt Ser	tac Tyr	ccg Pro	384
								ggg Gly								432
								ctg Leu								480
								gga Gly								528
								aaa Lys 185								576
gcc Ala	aac Asn	gtg Val 195	atg Met	gtc Val	atc Ile	ttt Phe	gtg Val 200	gac Asp	att Ile	999 Gly	ata Ile	gtg Val 205	gcc Ala	tca Ser	ggc Gly	624
								tat Tyr								672
								agg Arg								720
gcc Ala	tcc Ser	cac His	tgt Cys	att Ile 245	gtg Val	gtc Val	ctt Leu	tgc Cys	ttc Phe 250	ttt Phe	gtt Val	ccc Pro	tgt Cys	gtt Val 255	gtc Val	768
								gat Asp 265								816
								ctt Leu								864

912

936

ctg aga aac aag gag gtg aag aaa gct gtg ttg aaa ctt aga gac aaa Leu Arg Asn Lys Glu Val Lys Lys Ala Val Leu Lys Leu Arg Asp Lys 290 295 300
gta gca cat cct cag agg aaa taa Val Ala His Pro Gln Arg Lys 305 310
<210> 176 <211> 311 <212> PRT <213> Homo sapiens
<400> 176
Met Ser Asn Ala Ser Leu Val Thr Ala Phe Ile Leu Thr Gly Leu Pro 1 5 10 15
His Ala Pro Gly Leu Asp Ala Leu Leu Phe Gly Ile Phe Leu Val Val 20 25 30
Tyr Val Leu Thr Val Leu Gly Asn Leu Leu Ile Leu Leu Val Ile Arg 35 40
Val Asp Ser His Leu His Thr Pro Met Tyr Tyr Phe Leu Thr Asn Leu 50 60
Ser Phe Ile Asp Met Trp Phe Ser Thr Val Thr Val Pro Lys Met Leu 65 70 75 80
Met Thr Leu Val Ser Pro Ser Gly Arg Ala Ile Ser Phe His Ser Cys 85 90 95
Val Ala Gln Leu Tyr Phe Phe His Phe Leu Gly Ser Thr Glu Cys Phe 100 105 110
Leu Tyr Thr Val Met Ser Tyr Asp Arg Tyr Leu Ala Ile Ser Tyr Pro 115 120 125
Leu Arg Tyr Thr Ser Met Met Ser Gly Ser Arg Cys Ala Leu Leu Ala 130 135 140
Thr Gly Thr Trp Leu Ser Gly Ser Leu His Ser Ala Val Gln Thr Ile 145 150 155 160
Leu Thr Phe His Leu Pro Tyr Cys Gly Pro Asn Gln Ile Gln His Tyr 165 170 175
Phe Cys Asp Ala Pro Pro Ile Leu Lys Leu Ala Cys Ala Asp Thr Ser 180 185 190
Ala Asn Val Met Val Ile Phe Val Asp Ile Gly Ile Val Ala Ser Gly 195 200 205
Cys Phe Val Leu Ile Val Leu Ser Tyr Val Ser Ile Val Cys Ser Ile 210 215 220
Leu Arg Ile Arg Thr Ser Asp Gly Arg Arg Arg Ala Phe Gln Thr Cys 225 230 235 240
Ala Ser His Cys Ile Val Val Leu Cys Phe Phe Val Pro Cys Val Val 245 255

16U 200 PCT FINAL.ST25 Ile Tyr Leu Arg Pro Gly Ser Met Asp Ala Met Asp Gly Val Val Ala 265 Ile Phe Tyr Thr Val Leu Thr Pro Leu Leu Asn Pro Val Val Tyr Thr Leu Arg Asn Lys Glu Val Lys Lys Ala Val Leu Lys Leu Arg Asp Lys Val Ala His Pro Gln Arg Lys <210> 177 <211> 29 <212> DNA <213> Homo sapiens <400> 177 caaccagate cagcactact tetgtgacg 29 <210> 178 <211> 33 <212> DNA <213> Homo sapiens <400> 178 ttatttcctc tgaggatgtg ctactttgtc tct 33 <210> 179 <211> 50 <212> DNA <213> Homo sapiens <400> 179 taggagaagc cctttaaaag caggcaatag taaggacatc agtaacaata 50 <210> 180 <211> 50 <212> DNA <213> Homo sapiens <400> 180 gctgggtgct ctttatatcc ccagagggag agagaccaag ggtgagaaga 50 <210> 181 <211> 921 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (1)..(921) <223> <400> 181 atg gtg act gag ttt ctt ctt ctc ggc ttc tcc cac ctg gcc gac ctc Met Val Thr Glu Phe Leu Leu Gly Phe Ser His Leu Ala Asp Leu 48 cag ggc ttg ctc ttc tct gtc ttt ctc act atc tac ctg ctg acc gtg Gln Gly Leu Phe Ser Val Phe Leu Thr Ile Tyr Leu Leu Thr Val 96 gca ggc aat ttc ctc att gtg gtg ctg gtc tcc act gat gct gcc ctc Ala Gly Asn Phe Leu Ile Val Val Leu Val Ser Thr Asp Ala Ala Leu

cag tcc cct atg tac ttc ttc ctg cgc acc ctc tcg gcc ttg gag att Gln Ser Pro Met Tyr Phe Phe Leu Arg Thr Leu Ser Ala Leu Glu Ile

Page 104

192

													ctc Leu				240
													atg Met				288
													gcc Ala 110				336
													cca Pro				384
													tgg Trp				432
ggg Gly 145	gtg Val	ctg Leu	gtg Val	ggg Gly	ctg Leu 150	ggc Gly	cac His	acc Thr	cct Pro	ttc Phe 155	atc Ile	ttc Phe	tct Ser	ttg Leu	ccc Pro 160		480
													atc Ile				528
													ctg Leu 190				576
													ctc Leu				624
													cca Pro				672
													ctg Leu				720
													cgc Arg				768
													ttc Phe 270				816
	_										-		cgg Arg				864
		Lys											gtg Val				912
	att Ile																921
<21 <21 <21 <21	1 > 2 >	182 306 PRT Homo	sap:	iens													
< 40	0>	182												,	•	,	
Met 1	Val	Thr	Glu	Phe 5	Leu	Leu	Leu	Gly	Phe 10	Ser	His	Leu	Ala	Asp 15	Leu		
Gln	Gly	Leu	Leu 20	Phe	Ser	Val	Phe	Leu 25	Thr	Ile	Туr	Leu	Leu 30	Thr	Val		

Ala Gly Asn Phe Leu Ile Val Val Leu Val Ser Thr Asp Ala Ala Leu $35 \hspace{1.5cm} 40 \hspace{1.5cm} 45$

Gln Ser Pro Met Tyr Phe Phe Leu Arg Thr Leu Ser Ala Leu Glu Ile 50 55

Gly Tyr Thr Ser Val Thr Val Pro Leu Leu Leu His His Leu Leu Thr 70 75 80

Gly Arg Arg His Ile Ser Arg Ser Gly Cys Ala Leu Gln Met Phe Phe 85 90 95

Phe Leu Phe Phe Gly Ala Thr Glu Cys Cys Leu Leu Ala Ala Met Ala 100 105 110

Tyr Asp Arg Tyr Ala Ala Ile Cys Glu Pro Leu Arg Tyr Pro Leu Leu 115 120 125

Leu Ser His Arg Val Cys Leu Gln Leu Ala Gly Ser Ala Trp Ala Cys 130 $$135\$

Gly Val Leu Val Gly Leu Gly His Thr Pro Phe Ile Phe Ser Leu Pro 145 150150155

Phe Cys Gly Pro Asn Thr Ile Pro Gln Phe Phe Cys Glu Ile Gln Pro 165 170 175

Val Leu Gln Leu Val Cys Gly Asp Thr Ser Leu Asn Glu Leu Gln Ile 180 185 190

Ile Leu Ala Thr Ala Leu Leu Ile Leu Cys Pro Phe Gly Leu Ile Leu 195 200 205

Gly Ser Tyr Gly Arg Ile Leu Val Thr Ile Phe Arg Ile Pro Ser Val 210 215 220

Ala Gly Arg Arg Lys Ala Phe Ser Thr Cys Ser Ser His Leu Ile Val 225 230 240

Val Ser Leu Phe Tyr Gly Thr Ala Leu Phe Ile Tyr Ile Arg Pro Lys 245 250 255

Ala Ser Tyr Asp Pro Ala Thr Asp Pro Leu Val Ser Leu Phe Tyr Ala 260 265 270

Val Val Thr Pro Ile Leu Asn Pro Ile Ile Tyr Ser Leu Arg Asn Thr 275 280 285

Glu Val Lys Ala Ala Leu Lys Arg Thr Ile Gln Lys Thr Val Pro Met 290 295 300

Glu Ile 305

<210> 183

<211> 20 <212> DNA

<213> Homo sapiens

16U 200 PCT FINAL.ST25 <400> 183 ctcggcttct cccacctggc 20 <210> 184 23 <211> <212> DNA <213> Homo sapiens <400> 184 ggcgccaaag aagaggaaga aga 23 <210> 185 <211> 897 <212> DNA <213> Homo sapiens <220> CDS <221> <222> (1)..(897) <223> <400> 185 atg ggt cga gga aac agc act gaa gtg act gaa ttc cat ctt ctg gga Met Gly Arg Gly Asn Ser Thr Glu Val Thr Glu Phe His Leu Leu Gly 1048 10 ttt ggt gtc caa cac gaa ttt cag cat gtc ctt ttc att gta ctt ctt Phe Gly Val Gln His Glu Phe Gln His Val Leu Phe Ile Val Leu Leu 96 ctt atc tat gtg acc tcc ctg ata gga aat att gga atg atc tta ctc Leu Ile Tyr Val Thr Ser Leu Ile Gly Asn Ile Gly Met Ile Leu Leu 35 40 45144 atc aag acc gat tcc aga ctt caa aca ccc atg tac ttt ttt cca caa Ile Lys Thr Asp Ser Arg Leu Gln Thr Pro Met Tyr Phe Phe Pro Gln 50192 cat ttg gct ttt gtt gat atc tgt tat act tct gct atc act ccc aag His Leu Ala Phe Val Asp Ile Cys Tyr Thr Ser Ala Ile Thr Pro Lys 65 70 75 80240 atg ctc caa agc ttc aca gaa gaa aat aat ttg ata aca ttt cgg ggc Met Leu Gln Ser Phe Thr Glu Glu Asn Asn Leu Ile Thr Phe Arg Gly 85 90 95 288 tgt gtg ata caa ttc tta gtt tat gca aca ttt gca acc agt gac tgt Cys Val Ile Gln Phe Leu Val Tyr Ala Thr Phe Ala Thr Ser Asp Cys 336 105 tac ctc cta gct att atg gca atg gat tgt tat gtt gcc atc tgt aag Tyr Leu Leu Ala Ile Met Ala Met Asp Cys Tyr Val Ala Ile Cys Lys 115 120 125384 ccc ctt cgc tat ccc atg atc atg tcc caa aca gtc tac atc caa ctc Pro Leu Arg Tyr Pro Met Ile Met Ser Gln Thr Vall Tyr Ile Gln Leu 432 135 gta gct ggc tca tat att ata ggc tca ata aat gcc tct gta cat aca Val Ala Gly Ser Tyr Ile Ile Gly Ser Ile Asn Ala Ser Val His Thr 145 150 155 160480 ggt ttt aca ttt tca ctg tcc ttc tgc aag tct aat aaa atc aat cac 528 Gly Phe Thr Phe Ser Leu Ser Phe Cys Lys Ser Asn Lys Ile Asn His ttt ttc tgt gat ggt ctc cca att ctt gcc ctt tca tgc tcc aac att 576 Phe Phe Cys Asp Gly Leu Pro Ile Leu Ala Leu Ser Cys Ser Asn Ile 185 gac atc aac atc att cta gat gtt gtc ttt gtg gga ttt gac ttg;atg Asp Ile Asn Ile Ile Leu Asp Val Val Phe Val Gly Phe Asp Leu Met 195 200 205

ttc act gag ttg gtc atc atc ttt tcc tac atc tac att atg gtc acc Phe Thr Glu Leu Val Ile Ile Phe Ser Tyr Ile Tyr Ile Met Val Thr

			gct Ala						720
			gta Val						768
			tct Ser						816
			act Thr 280						864
			gaa Glu		taa				897

<210> 186

<211> 298

<212> PRT

<213> Homo sapiens

<400> 186

Met Gly Arg Gly Asn Ser Thr Glu Val Thr Glu Phe His Leu Leu Gly 1 $$ 5 $$ 10 $$ 15

Phe Gly Val Gln His Glu Phe Gln His Val Leu Phe Ile Val Leu Leu 20 25 30

Leu Ile Tyr Val Thr Ser Leu Ile Gly Asn Ile Gly Met Ile Leu Leu 35 40 45

Ile Lys Thr Asp Ser Arg Leu Gln Thr Pro Met Tyr Phe Phe Pro Gln 50 60

His Leu Ala Phe Val Asp Ile Cys Tyr Thr Ser Ala Ile Thr Pro Lys 65 70 75 80

Met Leu Gln Ser Phe Thr Glu Glu Asn Asn Leu Ile Thr Phe Arg Gly 85 90 95

Cys Val Ile Gln Phe Leu Val Tyr Ala Thr Phe Ala Thr Ser Asp Cys 100 105 110

Tyr Leu Leu Ala Ile Met Ala Met Asp Cys Tyr Val Ala Ile Cys Lys 115 120 125

Pro Leu Arg Tyr Pro Met Ile Met Ser Gln Thr Val Tyr Ile Gln Leu 130 135 140

Val Ala Gly Ser Tyr Ile Ile Gly Ser Ile Asn Ala Ser Val His Thr 145 150 155 160

Gly Phe Thr Phe Ser Leu Ser Phe Cys Lys Ser Asn Lys Ile Asn His 165 170 175

Phe Phe Cys Asp Gly Leu Pro Ile Leu Ala Leu Ser Cys Ser Asn Ile 180 \$190\$

Asp Ile Asn Ile Ile Leu Asp Val Val Phe Val Gly Phe Asp Leu Met 195 200 205

Phe	Thr 210	Glu	Leu	Val	lle	Ile 215	Phe	Ser	Tyr	Ile	Tyr 220	Ile	Met	Val	Thr	
11e 225	Leu	Lys	Met	Ser	Ser 230	Thr	Ala	Gly	Arg	Lys 235	Lys	Ser	Phe	Ser	Thr 240	
Cys	Ala	Ser	His	Leu 245	Thr	Ala	Val	Thr	11e 250	Phe	Tyr	Gly	Thr	Leu 255	Ser	
Туr	Met	Tyr	Le u 260	Gln	Pro	Gln	Ser	Asn 265	Asn	Ser	Gln	Glu	Asn 270	Met	Lys	
Val	Ala	Ser 275	lle	Phe	Tyr	Gly	Thr 280	Val	Ile	Pro	Met	Leu 285	Asn	Pro	Leu	
Ile	Tyr 290		Leu	Arg	Asn	Lys 295	Glu	G1 y	Lys							
<21 <21 <21 <21	1> 2>	187 930 DNA Homo	sap:	iens												
<22 <22 <22 <22	1> (2>	CDS (1).	. (936	0)												
	aca					act Thr										48
						ttt Phe										96
						ata Ile										144
atc Ile	aac Asn 50	aca Thr	gat Asp	tcc Ser	aga Arg	ttt Phe 55	caa Gln	aca Thr	ctc Leu	acg Thr	tac Tyr 60	ttt Phe	ttt Phe	cta Leu	caa Gln	192
						atc Ile										240
						gaa Glu										288
tgt Cys	gtg Val	ata Ile	caa Gln 100	ttc Phe	tta Leu	gtt Val	tat Tyr	gca Ala 105	aca Thr	ttt Phe	gca Ala	acc Thr	agt Ser 110	gac Asp	tgt Cys	336
						gca Ala										384
						atc Ile 135								Arg	Leu	432
gta Val 145	gct Ala	ggt Gly	tca Ser	tac Tyr	atc Ile 150	atg Met	ggc Gly	tca Ser	ata Ile	aat Asn 155	gcc Ala	tct Ser	gta Val	caa Gln	aca	480
						tcc Ser					Asn		Ile			528

				165					170	1 6U	200	PCT	FINA	L.ST	25	
									gct Ala							576
									ttt Phe							624
									tac Tyr							672
									agg Arg							720
									att Ile 250							768
									aat Asn							816
									att Ile							8 64
									aaa Lys							912
	aaa Lys				taa											930
<210 <213 <213	1> 3	188 309 PRT														
<21	1 >		sapi	iens												
<213 <213	1> 3 2> 1 3> 1	309 PRT	sapi	iens												
<213 <213 <213 <400	1> 3 2> 1 3> 1	309 PRT Homo			Ser	Thr	Glu	Val	Thr 10	Glu	Phe	туг	Leu	Leu 15	Gly	
<211 <212 <211 <400 Met 1	1> (2> 1) 3> 1 0> (1) Thr	309 PRT Homo 188 Leu	Gly	Asn 5										15		
<211 <212 <211 <400 Met 1	1> 1 2> 1 3> 1 0> 1 Thr	309 PRT Homo 188 Leu	Gly Gln 20	Asn 5 His	Glu	Phe	Trp	Cys 25	10	Leu	Phe	Ile	Val 30	15 Phe	Leu	
<211 <212 <211 <400 Met 1 Phe	1> 3 2> 1 3> 1 7 Thr Gly	309 PRT Homo 188 Leu Ala Tyr 35	Gly Gln 20 Val	Asn 5 His	Glu Ser	Phe Ile	Trp Met 40	Cys 25 Gly	10	Leu Ser	Phe Gly	Ile Ile 45	Val 30	15 Phe Leu	Leu Leu	
<211 <212 <211 <400 Met 1 Phe Leu	1> 5 2> 1 3> 1 0> 1 Thr Gly Ile	309 PRT Homo 188 Leu Ala Tyr 35	Gly Gln 20 Val	Asn 5 His Thr	Glu Ser Arg	Phe Ile Phe 55	Trp Met 40 Gln	Cys 25 Gly Thr	10 Ile Asn	Leu Ser Thr	Phe Gly Tyr 60	Ile Ile 45 Phe	Val 30 Ile Phe	15 Phe Leu Leu	Leu Leu Gln	
<21: <21: <21: <400 Met 1 Phe Leu Ile	11> (12> 13) (13) (14) (15) (15) (15) (15) (15) (15) (15) (15	309 PRT HOMO 108 Leu Ala Tyr 35 Thr	Gly Gln 20 Val	Asn 5 His Thr Ser	Glu Ser Arg Asp 70	Phe Ile Phe 55	Trp Met 40 Gln Cys	Cys 25 Gly Thr	10 Ile Asn Leu	Leu Ser Thr Ser 75	Phe Gly Tyr 60	Ile Ile 45 Phe	Val 30 Ile Phe	Phe Leu Leu Pro	Leu Gln Lys 80	
<21: <21: <21: <400 Met 1 Phe Leu Ile His 65	11> (2) 12 12 13	309 PRT Homo 188 Leu Ala Tyr 35 Thr	Gly Gln 20 Val Asp	Asn 5 His Thr Ser Val	Glu Ser Arg Asp 70	Phe Ile Phe 55 Ile Glu	Trp Met 40 Gln Cys	Cys 25 Gly Thr Tyr	10 Ile Asn Leu Thr	Leu Ser Thr Ser 75 Leu	Phe Gly Tyr 60 Ala	Ile Ile 45 Phe Ile	Val 30 Ile Phe Thr	Phe Leu Pro Gln 95	Leu Gln Lys 80 Gly	
<21: <21: <400 Met 1 Phe Leu Ile His 65 Met	11> (12> 13) (13) (14) (15) (15) (15) (15) (15) (15) (15) (15	309 PRT Homo 188 Leu Ala Tyr 35 Thr Ala Gln	Gly Gln 20 Val Asp Phe Ser Gln 100	Asn 5 His Thr Ser Val Phe 85	Glu Ser Arg Asp 70 Thr	Phe Ile Phe 55 Ile Glu Val	Trp Met 40 Gln Cys Glu	Cys 25 Gly Thr Tyr Lys	10 Ile Asn Leu Thr	Leu Ser Thr Ser 75 Leu	Tyr 60 Ala	Ile Ile 45 Phe Ile Leu	Val 30 Ile Phe Thr Phe	Phe Leu Pro Gln 95 Asp	Leu Gln Lys 80 Gly Cys	

130	. 1	135	16U 200 140	PCT FINAL.S	Т25
Val Ala Gly S 145	Ser Tyr Ile N 150	Met Gly Ser	Ile Asn Ala 155	Ser Val Glr	Thr 160
Gly Phe Thr C	Cys Ser Leu S 165	Ser Phe Cys	Lys Ser Asn 170	Ser Ile Asn 175	
Phe Phe Cys A	Asp Val Pro B 180	Pro Ile Leu 185	Ala Leu Ser	Cys Ser Asr 190	Val
Asp Ile Asn I 195	le Met Leu I	Leu Val Val 200	Phe Val Gly	Ser Asn Leu 205	Ile
Phe Thr Gly I 210		Tle Phe Ser 215	Tyr Ile Tyr 220	Ile Met Ala	Thr
Ile Leu Lys M 225	det Ser Ser S 230	Ser Ala Gly	Arg Lys Lys 235	Ser Phe Ser	Thr 240
Cys Ala Ser H	lis Leu Thr A 245	Ala Val Thr	Ile Phe Tyr 250	Gly Thr Leu 255	
Tyr Met Tyr L	Leu Gln Ser F 260	dis Ser Asn 265	Asn Ser Gln	Glu Asn Met 270	Lys
Val Ala Phe I 275	le Phe Tyr C	Gly Thr Val 280	Ile Pro Met	Leu Asn Pro 285	Leu
Ile Tyr Ser L 290		ys Glu Val 295	Lys Glu Ala 300	Leu Lys Val	Ile
Gly Lys Lys L 305	eu Phe				
<210> 189 <211> 522 <212> DNA <213> Homo s	apiens				
<220> <221> CDS <222> (1)(<223>	522)				
<400> 189 atg ctc caa a Met Leu Gln S 1		ilu Glu Lys			
tgc atg ata c Cys Met Ile G 2					
tat ctc ctg g Tyr Leù Leu A 35					
ccc ctt cac t Pro Leu His T 50	yr Thr Val I			Cys Ile His	
gta gct ggt t Val Ala Gly S 65					
ggt ttt gca t	tt tca ctg t	ct ttc tgc		aac atc aac age 111	cac 288

WO 03/089583 PCT/US03/11497

Gly	Phe	Ala	Phe	Ser 85	Leu	Ser	Phe	Cys	Lys 90				FINA Ile			
		tgt Cys														336
		aac Asn 115														384
		ggg Gly														432
		ccc Pro														480
		gaa Glu											taa			522
<210 <211 <212 <213	! > ! >	190 173 PRT Homo	sapi	ens												
< 400)>	190														
Met 1	Leu	Gln	Ser	Phe 5	Thr	Glu	Glu	Lys	Asn 10	Leu	Ile	Ser	Phe	Trp 15	Gly	
Cys	Met	Ile	Gln 20	Leu	Leu	Val	Tyr	Ala 25	Thr	Phe	Ala	Thr	Ser 30	Asp	Cys	
Tyr	Leu	Leu 35	Ala	Met	Ile	Ala	Val 40	Asp	His	Tyr	Val	Ala 45	Ile	Cys	Lys	
Pro	Leu 50	His	Tyr	Thr	Val	Ile 55	Thr	Ser	Gln	Thr	Val 60	Cys	Ile	His	Leu	
Val 65	Ala	Gly	Ser	Туr	Ile 70	Met	Gly	Ser	Ile	Asn 75	Ala	Ser	Val	His	Thr 80	
Gly	Phe	Ala	Phe	Ser 85	Leu	Ser	Phe	Cys	Lys 90	Ser	Asn	Asn	Ile	Asn 95	His	
Phe	Phe	Cys	Asp 100	Gly	Pro	Pro	Ile	Leu 105	Ala	Leu	Ser	Cys	Ser 110	Asn	Ile	
Asp	Ile	Asn 115	Ile	Met	Leu	Leu	Val 120	Val	Phe	Val	Gly	Phe 125	Asn	Leu	Met	
Phe	Thr 130	Gly	Leu	Glu	Asn	Met 135	Lys	Val	Ala	Ser	Ile 140	Phe	Tyr	Gly	Thr	
Val 145	Ile	Pro	Met	Leu	Asn 150	Pro	Leu	Ile	Tyr	Ser 155	Leu	Arg	Asn	Lys	Glu 160	
Val	Lys	Glu	Ala	Leu 165	Lys	Leu	Ile	Gly	Lys 170	Lys	Phe	Phe				
<210 <211 <212 <213	>	191 499 DNA Homo	sapi	.ens												

<220> <221> CDS				
<221> CDS <222> (43)(3 <223>	372)			
<400> 191 cgttatgtgg cctt	ctgtaa cccac	tccat tatccagggg	g tt atg tcc cag aga Met Ser Gln Arg l	54
			atg ggt ttc cta aat Met Gly Phe Leu Asn 20	102
			g aac ttc tgc aaa tcc u Asn Phe Cys Lys Ser 35	150
			. cca att att gcc cta) Pro Ile Ile Ala Leu 50	198
			a tta aca gta ttt gtg 1 Leu Thr Val Phe Val 65	246
			atc att tcc tgc ata . Ile Ile Ser Cys Ile 80	294
			gct gca ggg aag aaa Ala Ala Gly Lys Lys 100	342
aaa gtc tct cta Lys Val Ser Leu			pcagtcac cattttctat	392
ggggttctct ctta	catgta tctat	gccat cgtattaatg	agtotoaaaa acaagaaaaa	452
		gccat cgtattaatg		452 499
	ttatgg catta			
gtggcctctg tgtt <210> 192 <211> 109 <212> PRT	ttatgg catta			
<pre>gtggcctctg tgtt <210> 192 <211> 109 <212> PRT <213> Homo sap <400> 192</pre>	ttatgg catta iens	ttatt cccatgttaa		
<pre>gtggcctctg tgtt <210> 192 <211> 109 <212> PRT <213> Homo sap <400> 192 Met Ser Gln Arg 1</pre>	ttatgg catta niens Leu Cys Ile 5	ttatt cccatgttaa Lys Leu Leu Val 10	acccett Ser Ser Tyr Val Met	
gtggcctctg tgtt <210> 192 <211> 109 <212> PRT <213> Homo sap <400> 192 Met Ser Gln Arg 1 Gly Phe Leu Asn 20	iens Leu Cys Ile 5 Ala Ser Ile	Lys Leu Leu Val 10 Asn Ile Ser Phe 25	Ser Ser Tyr Val Met 15	
Style="font-size: 150;"> Style="font-size: 1	iens Leu Cys Ile 5 Ala Ser Ile Lys Thr Ile	Lys Leu Leu Val 10 Asn Ile Ser Phe 25 Asn His Phe Phe	Ser Ser Tyr Val Met 15 Thr Phe Ser Leu Asn 30 Cys Asp Glu Pro Pro	
Style="font-size: 150;"> Style="font-size: 1	ttatgg catta riens Leu Cys Ile 5 Ala Ser Ile Lys Thr Ile Pro Cys Ser 55	Lys Leu Leu Val 10 Asn Ile Ser Phe 25 Asn His Phe Phe 40 Asn Ile Asp Leu	Ser Ser Tyr Val Met 15 Thr Phe Ser Leu Asn 30 Cys Asp Glu Pro Pro 45	
Style="font-size: 150;"> Style="font-size: 1	ttatgg catta fiens Leu Cys Ile 5 Ala Ser Ile Lys Thr Ile Pro Cys Ser 55 Gly Leu Asn 70	Lys Leu Leu Val 10 Asn Ile Ser Phe 25 Asn His Phe Phe 40 Asn Ile Asp Leu Leu Met Cys Thr	Ser Ser Tyr Val Met 15 Thr Phe Ser Leu Asn 30 Cys Asp Glu Pro Pro 45 Asn Ile Met Leu Leu 60 Val Met Val Val Ile	

<210> 193 <211> 681 <212> DNA <213> Homo	sapiens		160	200 PCT	FINAL.ST	25
<220> <221> CDS <222> (1) <223>	. (681)					
	gac cgc tac Asp Arg Tyr 5					
	ttc cca gag Phe Pro Glu 20					
	ttt gta aac Phe Val Asn					
	tgt ggc agc Cys Gly Ser					
	gta aag ttg Val Lys Leu 70				Tyr Gln	
	ttt atg ctt Phe Met Leu . 85					
	cat ctc ttc His Leu Phe 100		Ala Ile			
	cgc ctc cag Arg Leu Gln					
	ttg tac tat Leu Tyr Tyr					
	tat gcc tta Tyr Ala Leu 150				Val Phe	
	att cca atg Ile Pro Met 165					
	aaa gat gcc Lys Asp Ala 180		Met Leu			
	aaa tat tgt Lys Tyr Cys					
ctc ctg gct Leu Leu Ala 210	gcc atg ggt Ala Met Gly	tat gac tgc Tyr Asp Cys 215	tat gag Tyr Glu	gca atc Ala Ile 220	tcc aag Ser Lys	ccc 672 Pro
ctg ctt taa Leu Leu 225					<i>y</i> *	681
<210> 194 <211> 226 <212> PRT						
	sapiens					
13002 134						

Page 114

16U 200 PCT FINAL.ST25 Met Ala Tyr Asp Arg Tyr Met Ala Ile Ser Lys Pro Leu Leu Tyr Ser 1 5 10 Arg Ala Thr Phe Pro Glu Leu Cys Ala Ser Leu Val Glu Ala Ser His 20 25 30 Leu Gly Gly Phe Val Asn Ser Thr Ile Ile Thr Ser Glu Thr Pro Thr Leu Ser Phe Cys Gly Ser Asn Ile Ile Asp Asp Phe Phe Cys Asp Leu 50 55 60 Pro Pro Leu Val Lys Leu Val Cys Asp Val Lys Glu Arg Tyr Gln Ala Val Leu His Phe Met Leu Ala Ser Asn His His Ser His Cys Thr Tyr Ser Cys Val His Leu Phe Ile Ile Ala Ala Ile Ser Lys Ile Arg Ser 100 105 110Ile Lys Gly Arg Leu Gln Val Phe Ser Thr Cys Gly Ser Pro Leu Thr 115 120 125 Ala Leu Thr Leu Tyr Tyr Gly Ala Ile Phe Phe Ile Tyr Ser Gln Pro 130 135 140 Arg Thr Ser Tyr Ala Leu Lys Met Asp Lys Leu Gly Ser Val Phe Tyr 145 150 160 Thr Val Val Ile Pro Met Leu Asn Pro Leu Ile Tyr Ser Leu Arg Asn 165 170 175 Lys Asp Val Lys Asp Ala Leu Lys Lys Met Leu Asp Arg Leu Gln Phe 180 190Leu Lys Glu Lys Tyr Cys Arg Tyr Gly Leu Ala Cys Ser Glu Arg Tyr 195 200 205 Leu Leu Ala Ala Met Gly Tyr Asp Cys Tyr Glu Ala Ile Ser Lys Pro 210 215 220 <211> 1095 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (1)..(1095) <223> <400> 195 atg gcc tct gag acc ttc aac act gaa gac cca gcc ggg ttg atg cac Met Ala Ser Glu Thr Phe Asn Thr Glu Asp Pro Ala Gly Leu Met His 1 5 10 1548 tcg gat gcc ggc acc agc tgc ccc gtc ctt tgc aca tgc cgt aac cag Ser Asp Ala Gly Thr Ser Cys Pro Val Leu Cys Thr Cys Arg Asn Gln 20 25 3096

										1 6U	200	PCT	FINA	AL.ST	25	
			tgt Cys													144
			acc Thr													192
			ggc Gly													240
			aac Asn													288
			ttg Leu 100													336
			gac Asp													384
			aac Asn													432
			cag Gln													480
			ctg Leu													528
			ggc Gly 180													576
			ctg Leu													624
			ccg Pro													672
			gag Glu													720
			aac Asn													768
aac Asn	ttt Phe	999 Gly	cac His 260	agg Arg	caa Gln	gcg Ala	gtt Val	gtg Val 265	ggt Gly	gga Gly	ctt Leu	tcc Ser	aat Asn 270	ccc Pro	ctc Leu	816
			Gly ggg													864
			gag Glu													912
			act Thr													960
			tac Tyr													1008
			gcc Ala 340													1056

Page 116

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

	cgc Arg											tga				1095
<210 <211 <212 <213	l> 3 2> I	196 364 PRT Homo	sap:	iens												
< 400)> 1	196														
Met 1	Ala	Ser	Glu	Thr 5	Phe	Asn	Thr	Glu	Asp 10	Pro	Ala	Gly	Leu	Met 15	His	
Ser	Asp	Ala	Gly 20	Thr	Ser	Суѕ	Pro	Val 25	Leu	Суѕ	Thr	Cys	Arg 30	Asn	Gln	
Val	Val	Asp 35	Cys	Ser	Ser	Gln	Arg 40	Lev	Phe	Ser	Val	Pro 45	Pro	Asp	Leu	
Pro	Met 50	Asp	Thr	Arg	Asn	Leu 55	Ser	Leu	Ala	His	Asn 60	Arg	Ile	Thr	Ala	
Val 65	Pro	Pro	Gly	Tyr	Leu 70	Thr	Cys	Tyr	Met	Glu 75	Leu	Gln	Val	Leu	Asp 80	
Leu	His	Asn	Asn	Ser 85	Leu	Met	Glu	Leu	Pro 90	Arg	Gly	Leu	Phe	Leu 95	His	
Ala	Lys	Arg	Leu 100	Ala	His	Leu	Asp	Leu 105	Ser	Туr	Asn	Asn	Phe 110	Ser	His	
Val	Pro	Ala 115	Asp	Met	Phe	Gln	Glu 120	Ala	His	Gly	Leu	Val 125	His	Ile	Asp	
Leu	Ser 130	His	Asn	Pro	Trp	Leu 135	Arg	Arg	Val	His	Pro 140	Gln	Ala	Phe	Gln	
Gly 145	Leu	Met	Gln	Leu	Arg 150	Asp	Leu	Asp	Leu	Ser 155	Tyr	G1 y	Gly	Leu	Ala 160	
Phe	Leu	Ser	Leu	Glu 165	Ala	Leu	Glu	Gly	Leu 170	Pro	Gly	Leu	Val	Thr 175	Leu	
Gln	Ile	Gly	Gly 180	Asn	Pro	Trp	Val	Cys 185	Gly	Cys	Thr	Met	Glu 190	Pro	Leu	
Leu	Lys	Trp 195	Leu	Arg	Asn	Arg	11e 200	Gln	Arg	Cys	Thr	Ala 205	Glu	Ser	Gly	
Ser	Gly 210	Leu	Pro	Glu	Glu	Ser 215	Glu	Pro	Glu	Ser	Trp 220	Thr	Gly	Gln	Arg	
Ala 225	Ala	Val	Glu	Phe	Gln 230	Asp	Leu	Met	Gln	Leu 235	Gln	Asp	Leu	,	,240	
Ser	Tyr	Glu	Asn	Leu 245	Ala	Phe	Leu	Lys	Leu 250	Lys	Ala	Leu	Ser		Val	
Asn	Phe	Gly	His 260	Arg	Gln	Ala	Val	Val 265	Gly	Gly	Leu	Ser	Asn 270	Pro	Leu	

Ser Ph	e Pro 275		Туr	Leu	Thr	Leu 280	Pro	Gly	Phe	Cys	Val 285	Thr	Asp	Ser	
Gln Le 29		Glu	Cys	Arg	Gly 295	Pro	Pro	Glu	Val	Glu 300	Gly	Ala	Pro	Leu	
Phe Se 305	r Leu	Thr	Glu	Glu 310	Ser	Phe	Lys	Ala	Cys 315	His	Leu	Thr	Leu	Thr 320	
Leu As	p Asp	Тyr	Leu 325	Phe	Ile	Ala	Phe	Val 330	Gly	Phe	Val	Val	Ser 335	Ile	
Ala Se	r Val	Ala 340	Thr	Asn	Phe	Leu	Leu 345	Gly	Ile	Thr	Ala	Asn 350	Cys	Cys	
His Ar	g Trp 355		Lys	Ala	Ser	Glu 360	Glu	Glu	Glu	Ile					
<210> <211> <212> <213>	197 27 DNA Homo	sap	iens												
<400> gggatt	197 tggt	gtcc	aaca	eg aa	attto	ca .									2
<210> <211> <212> <213>	198 34 DNA Homo	sap	iens												
<400> gagccta	198 ataa	tata	tgago	cc aç	gctad	cgagt	t t g	ga							3
<210> <211> <212> <213>	199 32 DNA Homo	sap:	iens												
<400> gtcacto	199 gaat	tctai	tette	ct go	gat t	tggt	gc								3:
<210><211><211><212><213>	200 32 DNA Homo	sap	iens												
<400> aaacct	200 gttt	gtaca	agago	gc at	ttat	tgag	g cc								3:
<210> <211> <212> <213>	201 35 DNA Homo	sapi	iens												
<400> gatatea	201 attt	·	gctgo	a to	atac	aatt	att	gg							3:
<210> <211> <212> <213>	202 33 DNA Homo	sapi	iens												
<400> ctccaac	202 cca (gtgaa	cato	ca aç	ıttaa	atco	cac	:							3:

<210>	203	
<211>	36	
<212>	DNA	
<213>	Homo sapiens	
<400>	203	
		2.0
ctaage	tatt agttagttca tatgtcatgg gtttcc	36
<210>	204	
<211>	36	
<212>	DNA	
<213>	Homo sapiens	
< 400>	204	
		3.6
Ctcact	aata cgatggcata gatacatgta agagag	36
<210>	205	
<211>	50	
<212>		
<213>	Homo sapiens	
< 400>	205	
	cato taaatgaago otgagaaaco cagcactaco cacttgttag	50
argere	care cadargaage ergagaaace eageaceace eacetgerag	30
<210>	206	
<211>	50	
<212>	DNA	
<213>	Homo sapiens	
<400>	206	
acatco	atta tataacaggg ttaatatact tgtaaagaat agcacctaga	50
40000	tata tataadayyy taatatata tytaaayaat ageacetaya	50
<210>	207	
<211>	50	
<212>	DNA	
<213>		
(213)	Homo sapiens	
<400>	207	
		50
	207 ataa attotgoatg aaattggggg tggggottgt actacttttg	50
		50
aaatgt	ataa attotgoatg aaattggggg tggggottgt actacttttg	50
aaatgt <210>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208	50
aaatgt	ataa attotgoatg aaattggggg tggggottgt actacttttg	50
aaatgt <210>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50	50
<210><211><212>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50 DNA	50
<210><211>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50	50
<210><211><212><213>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50 DNA Homo sapiens	50
<210><211><212>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50 DNA	
<210><211><212><213><400>	ataa attotgoatg aaattggggg tggggcttgt actacttttg 208 50 DNA Homo sapiens 208	50
<210><211><212><213><400>	ataa attotgoatg aaattggggg tggggottgt actacttttg 208 50 DNA Homo sapiens	
<210><211><212><213><400>	ataa attotgoatg aaattggggg tggggcttgt actacttttg 208 50 DNA Homo sapiens 208	
<pre><210> <211> <212> <213> <400> atgttc</pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag	
<pre><210> <211> <212> <213> <400> atgttc</pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag	
<pre><210> <211> <212> <213> <400> atgttc</pre> <pre><210><210></pre>	208 50 DNA Homo sapiens 208 catc taaatgaage etgagaaace cagcactace cacttgttag	
<pre><210> <211> <212> <213> <400> atgttc</pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag	
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag	
<pre><210> <211> <212> <213> <400> atgttc</pre> <pre><210><210></pre>	208 50 DNA Homo sapiens 208 catc taaatgaage etgagaaace cagcactace cacttgttag	
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens	
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 50 DNA Homo sapiens	50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens	
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 50 DNA Homo sapiens	50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 50 DNA Homo sapiens	50
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213> <400 atgttc <210> <211> <212> <213> <400> acatcc</pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 50 DNA Homo sapiens 209 301 209 302 303 304 305 305 306 307 307 308 308 308 308 308 308 308 308 308 308	50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 209 209 209 209 209 209 209 209 209	50
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213> <400> atgttc <211> <212> <213> <400> acatcc <210> <211></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 400 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga	50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga	50
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213> <400> atgttc <211> <212> <213> <400> acatcc</pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 400 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga	50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga	50
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213> <400> acatcc <210> <212> <213> <400> acatcc</pre>	208 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens	50
<pre></pre>	208 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 50 DNA Homo sapiens	50 50
<pre></pre>	208 50 DNA Homo sapiens 209 Atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 Atat tttaaattgg ccagggggg tggggcttgt actatatece	50 50
<pre></pre>	208 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 50 DNA Homo sapiens	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaage etgagaaace cagcactace cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ecaggegegg tggetcacge etataatece	50 50
<pre><210> <211> <212> <213> <400> atgttc <210> <211> <212> <213> <400> acatcc <210> <212> <213> <400> acatcc <210> <211> <212> <213></pre>	208 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc	50 50
<pre></pre>	208 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggegegg tggetcacge ctataatcce 211	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc 211 50	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc 211 50 DNA	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc 211 50	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc 211 50 DNA	50 50
<pre></pre>	208 50 DNA Homo sapiens 208 catc taaatgaagc ctgagaaacc cagcactacc cacttgttag 209 50 DNA Homo sapiens 209 atta tataacaggg ttaatatact tgtaaagaat agcacctaga 210 50 DNA Homo sapiens 210 atat tttaaattgg ccaggcgcgg tggctcacgc ctataatccc 211 50 DNA	50 50

16U 200 PCT FINAL.ST25 ggctcacgcc tataatccca gçactttggg aggccgaggc aggtggatca	50
<210> 212 <211> 50 <212> DNA <213> Homo sapiens	
<400> 212 tcccaaatat atatatatac acacacacac acacacac	50
<210> 213 <211> 50 <212> DNA <213> Homo sapiens	
<400> 213 cacacacaca tatatataca cacacatata tttataatca tttaacaaca	50
<210> 214 <211> 933 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (1)(930) <223>	
<400> 214 atg gca gag atg aac ctc acc ttg gtg acc gag ttc ctc ctt att gca Met Ala Glu Met Asn Leu Thr Leu Val Thr Glu Phe Leu Leu Ile Ala 1 5 10 15	48
ttc act gaa tat cct gaa tgg gca ctc cct ctc ttc ctc ttg ttt tta Phe Thr Glu Tyr Pro Glu Trp Ala Leu Pro Leu Phe Leu Leu Phe Leu 20 25 30	96
ttt atg tat ctc atc acc gta ttg ggg aac tta gag atg att att ctg Phe Met Tyr Leu Ile Thr Val Leu Gly Asn Leu Glu Met Ile Ile Leu 35 40 45	144
atc ctc atg gat cac cag ctc cac gct cca atg tat ttc ctt ctg agt Ile Leu Met Asp His Gln Leu His Ala Pro Met Tyr Phe Leu Leu Ser 50 55 60	192
cac ctc gct ttc atg gac gtc tgc tac tca tct atc act gtc ccc cag His Leu Ala Phe Met Asp Val Cys Tyr Ser Ser Ile Thr Val Pro Gln 65 70 75 80	240
atg ctg gca gtg ctg ctg gag cat ggg gca gct tta tct tac aca cgc Met Leu Ala Val Leu Leu Glu His Gly Ala Ala Leu Ser Tyr Thr Arg 85 90 95	288
tgt gct gct cag ttc ttt ctg ttc acc ttc ttt ggt tcc atc gac tgc Cys Ala Ala Gln Phe Phe Leu Phe Thr Phe Phe Gly Ser Ile Asp Cys 100 105 110	336
tac ctc ttg gcc ctc atg gcc tat gac cgc tac ttg gct gtg tgc cag Tyr Leu Leu Ala Leu Met Ala Tyr Asp Arg Tyr Leu Ala Val Cys Gln 115 120 125	384
ccc ctg ctt tat gtc acc atc ctg aca cag cag gcc cgc ttg agt ctt Pro Leu Leu Tyr Val Thr Ile Leu Thr Gln Gln Ala Arg Leu Ser Leu 130 135 140	432
gtg gct ggg gct tac gtt gct ggt ctc atc agt gcc ttg gtg cgg aca Val Ala Gly Ala Tyr Val Ala Gly Leu Ile Ser Ala Leu Val Arg Thr 145 150 155 . 160	480
gtc tca gcc ttc act ctc tcc ttc tgt gga acc agt gag att gac ttt Val Ser Ala Phe Thr Leu Ser Phe Cys Gly Thr Ser Glu Ile Asp Phe 165 170 175	528
att ttc tgt gac ctc cct cct ctg tta aag ttg acc tgt ggg gag agc Ile Phe Cys Asp Leu Pro Pro Leu Leu Lys Leu Thr Cys Gly Glu Ser 180 185 190	576

16U 200 PCT	LINAL	. ST25
-------------	-------	--------

		caa Gln 195														624
		atg Met														672
		ggg Gly														720
		cac His														768
		ttg Leu														816
		gtg Val 275														864
		ctg Leu														912
		gcc Ala				taa										933
<210 <211 <211 <211	1 > 1 2 > 1	215 310 PRT Homo	sap	iens												
<400)> :	215														
Met 1	Ala	Glu	Met	Asn	Leu	Thr	1 412	v. i					_	- 1.		
				5			Deu	vaı	Thr 10	Glu	Phe	Leu	Leu	11e 15	Ala	
Phe	Thr	Glu	Tyr 20	5					10					15		
		Glu Tyr 35	20	5 Pro	Glu	Trp	Ala	Leu 25	10 Pro	Leu	Phe	Leu	Leu 30	15 Phe	Leu	
Phe	Met	Туг	20 Leu	5 Pro Ile	Glu Thr	Trp Val	Ala Leu 40	Leu 25 Gly	10 Pro Asn	Leu Leu	Phe Glu	Leu Met 45	Leu 30	15 Phe Ile	Leu Leu	
Phe Ile	Met Leu 50	Tyr 35	20 Leu Asp	5 Pro Ile His	Glu Thr Gln	Trp Val Leu 55	Ala Leu 40 His	Leu 25 Gly Ala	10 Pro Asn Pro	Leu Leu Met	Phe Glu Tyr 60	Leu Met 45 Phe	Leu 30 Ile Leu	15 Phe Ile Leu	Leu Leu Ser	
Phe Ile His	Met Leu 50 Leu	Tyr 35 Met	20 Leu Asp Phe	Pro Ile His	Glu Thr Gln Asp 70	Trp Val Leu 55 Val	Ala Leu 40 His	Leu 25 Gly Ala Tyr	10 Pro Asn Pro Ser	Leu Leu Met Ser 75	Phe Glu Tyr 60	Leu Met 45 Phe	Leu 30 Ile Leu Val	Phe Ile Leu Pro	Leu Ser Gin 80	
Phe Ile His 65	Met Leu 50 Leu	Tyr 35 Met	20 Leu Asp Phe	Pro Ile His Met Leu 85	Glu Thr Gln Asp 70 Leu	Trp Val Leu 55 Val	Ala Leu 40 His Cys	Leu 25 Gly Ala Tyr	Pro Asn Pro Ser Ala	Leu Leu Met Ser 75	Phe Glu Tyr 60	Leu Met 45 Phe Thr	Leu 30 Ile Leu Val	Phe Ile Leu Pro Thr	Leu Ser Gln 80 Arg	
Phe Ile His 65 Met	Met Leu 50 Leu Leu	Tyr 35 Met Ala	20 Leu Asp Phe Val Gln 100	Pro Ile His Met Leu 85	Glu Thr Gln Asp 70 Leu	Trp Val Leu S5 Val	Ala Leu 40 His Cys	Leu 25 Gly Ala Tyr Gly	Pro Asn Pro Ser Ala 90 Phe	Leu Met Ser 75 Ala	Phe Glu Tyr 60 Ile Leu Gly	Leu Met 45 Phe Thr Ser	Leu 30 Ile Leu Val Tyr	Phe Ile Leu Pro Thr 95	Leu Ser Gin 80 Arg	
Phe Ile His 65 Met Cys	Met Leu 50 Leu Ala	Tyr 35 Met Ala Ala	20 Leu Asp Phe Val Gln 100	Pro Ile His Met Leu B5 Phe	Glu Thr Gln Asp 70 Leu Phe	Trp Val Leu S5 Val Glu Leu	Ala Leu 40 His Cys Phe	Leu 25 Gly Ala Tyr Gly Thr 105	Pro Asn Pro Ser Ala 90 Phe	Leu Met Ser 75 Ala Phe	Phe Glu Tyr 60 Ile Leu Gly	Leu Met 45 Phe Thr Ser Ser	Leu 30 Ile Leu Val Tyr Ile 110	Phe Ile Leu Pro Thr 95 Asp	Leu Ser Gln 80 Arg Cys	

741 501 1114	Phe Thr 165		r Phe	Cys	Gly 170	Thr	Ser	Glu	lle	Asp 175	Phe	
Ile Phe Cys	Asp Leu 180	Pro Pr	o Leu	Leu 185	Lys	Leu	Thr	Cys	G1 y 190	Glu	Ser	
Tyr Thr Gln 195	Glu Val	Leu Il	e Ile 200	Met	Phe	Ala	lie	Phe 205	Val	lle	Pro	
Ala Ser Met 210	Val Val	Ile Le 21		Ser	Туr	Leu	Phe 220	Ile	lle	Val	Ala	
lle Met Gly 225	Ile Pro	Ala Gl 230	y Ser	Gln	Ala	Lys 235	Thr	Phe	Ser	Thr	Cys 240	
Thr Ser His	Leu Thr 245	Ala Va) Ser	Leu	Phe 250	Phe	Gly	Thr	Leu	11e 255	Phe	
Met Tyr Leu	Arg Gly 260	Asn Se	r Asp	Gln 265	Ser	Ser	Glu	Lys	Asn 270	Arg	Val	
Val Ser Val 275		Thr Gl	ı Val 280	Ile	Pro	Met	Leu	Asn 285	Pro	Leu	Ile	
Tyr Ser Leu 290	Arg Asn	Lys Gl 29		Lys	Glu	Ala	Leu 300	Arg	Lys	lle	Leu	
Asn Arg Ala 305	Lys Leu	Ser 310				ė.						
<210> 216 <211> 933 <212> DNA <213> Homo	sapiens											
<220>												
<221> CDS	. (930)											
<221> CDS <222> (1).	aac aag											48
<221> CDS <222> (1). <223> <400> 216 atg gaa ggc Met Glu Gly	aac aag Asn Lys 5 cca gca	Thr Tr	lle gatt	Thr	Asp 10 ctc	lle tgt	Thr gga	Leu	Pro	Arg 15 tct	Phe gcc	48 96
<221> CDS <222> (1). <223> <400> 216 atg gaa ggc Met Glu Gly 1 cag gtt ggt	aac aag Asn Lys 5 cca gca Pro Ala 20	Ctg gad Leu Glo	att lle	Thr ctc Leu 25	Asp 10 ctc Leu	lle tgt Cys gtc	Thr gga Gly	ctt Leu	Pro ttc Phe 30	Arg 15 tct Ser	Phe gcc Ala atc	
<221> CDS <222> (1). <223> <400> 216 atg gaa ggc Met Glu Gly 1 cag gtt ggt Gln Val Gly ttc tat aca Phe Tyr Thr	aac aag Asn Lys 5 cca gca Pro Ala 20 ctc acc Leu Thr	Thr Tr	g att lle ggg Gly 40 caca	Thr ctc Leu 25 aat Asn	Asp 10 ctc Leu ggg Gly	tgt Cys gtc val	Thr gga Gly atc Ile	ctt Leu ttt Phe 45	ttc Phe 30 ggg Gly	Arg 15 tct Ser att Ile	Phe gcc Ala atc Ile	96
<221> CDS <222> (1). <223> <400> 216 atg gaa ggc Met Glu Gly 1 cag gtt ggt Gln Val Gly ttc tat aca Phe Tyr Thr 35 tgc ctg gac Cys Leu Asp	aac aag Asn Lys 5 cca gca Pro Ala 20 ctc acc Leu Thr tgt aag Cys Lys	ctg gac Leu Gli ctg ctc Leu Leu ctt cac Leu His 55	g att g att g ggg g ggy 40 c aca a Thr	Thr ctc Leu 25 aat Asn ccc Pro	Asp 10 ctc Leu ggg Gly atg Met	tgt Cys gtc val tac Tyr	gga Gly atc Ile ttc Phe 60 tat	ctt Leu ttt Phe 45 ttc Phe	ttc Phe 30 ggg Gly ctc Leu	Arg 15 tct Ser att Ile tca Ser	Phe gcc Ala atc Ile cac His	96 144
<221> CDS <222> (1). <223> <400> 216 atg gaa ggc Met Glu Gly 1 cag gtt ggt Gln Val Gly ttc tat aca Phe Tyr Thr 35 tgc ctg gac Cys Leu Asp 50 ctg gcc att Leu Ala Ile	aac aag Asn Lys 5 cca gca Pro Ala 20 ctc acc Leu Thr tgt aag Cys Lys gtt gac Val Asp	ctg gar Leu Gli ctg ctc Leu Leu ctt car Leu Hi: 55 ata tcc 70	g att le ggg g 40 acas Thr	Thr ctc Leu 25 aat Asn ccc Pro gct Ala agc	Asp 10 ctc Leu ggg Gly atg Met tcc Ser	tgt Cys gtc val tac Tyr aac Asn 75	Thr gga Gly atc Ile ttc Phe 60 tat Tyr	ctt Leu ttt Phe 45 ttc Phe gtc Val	pro ttc Phe 30 ggg Gly ctc Leu ccc Pro	arg 15 tct Ser att Ile tca Ser aag Lys	gcc Ala atc Ile cac His atg Met 80 tgc	96 144 192

10		105	16U 200 PCT	FINAL.ST25
att ttg gtg gt Ile Leu Val Va 115				
tta cgt tac aa Leu Arg Tyr As 130				
gtg gct tcc tg Val Ala Ser Tr 145				
ctc atc ctg ag Leu Ile Leu Se				
ttc tgt gaa at Phe Cys Glu Il 18	le Leu Ser Val			
ctc aac cag gt Leu Asn Gln Va 195				
cca ctc tgc ct Pro Leu Cys Le 210				
ttg agg atc ca Leu Arg Ile Gl 225				
tcc tcc cac ct Ser Ser His Le	tt tgc gtg gtg eu Cys Val Val 245	gga ctc ttc Gly Leu Phe 250	ttt ggc agc Phe Gly Ser	gcc att gtc 768 Ala Ile Val 255
acg tac atg gc Thr Tyr Met Al 26	la Pro Lys Ser			
ctt tcc ctg tt Leu Ser Leu Ph 275				
tat agc cta ag Tyr Ser Leu Ar 290				
agg aag gag ag Arg Lys Glu Ar 305				933
<210> 217 <211> 310 <212> PRT <213> Homo sa	apiens			
<400> 217				
Met Glu Gly As 1	sn Lys Thr Trp 5	Ile Thr Asp 10	Ile Thr Leu	Pro Arg Phe 15
Gln Val Gly Pr 20		Ile Leu Leu 25	Cys Gly Leu	Phe Ser Ala 30
Phe Tyr Thr Le	eu Thr Leu Leu	Gly Asn Gly 40	Val Ile Phe 45	Gly Ile Ile
Cys Leu Asp Cy 50	ys Lys Leu His 55	Thr Pro Met	Tyr Phe Phe 60	Leu Ser His
Leu Ala Ile Va	al Asp Ile Ser	Tyr Ala Ser	Asn Tyr Val	Pro Lys Met

16U 200 PCT FINAL.ST25 65 70 Leu Thr Asn Leu Met Asn Gln Glu Ser Thr Ile Ser Phe Phe Pro Cys Ile Met Gln Thr Phe Leu Tyr Leu Ala Phe Ala His Val Glu Cys Leu $100 \hspace{1cm} 105 \hspace{1cm} 110$ Leu Arg Tyr Asn Ser Leu Met Ser Trp Arg Val Cys Thr Val Leu Ala 130 $$135\$ Val Ala Ser Trp Val Phe Ser Phe Leu Leu Ala Leu Val Pro Leu Val Leu Ile Leu Ser Leu Pro Phe Cys Gly Pro His Glu Ile Asn His Phe 165 170 175Phe Cys Glu Ile Leu Ser Val Leu Lys Leu Ala Cys Ala Asp Thr Trp 180 185 190 Leu Asn Gln Val Val Ile Phe Ala Ala Cys Val Phe Ile Leu Val Gly 195 200 205 Pro Leu Cys Leu Val Leu Val Ser Tyr Leu Arg Ile Leu Ala Ala Ile 210 215 220 Leu Arg Ile Gln Ser Gly Glu Gly Arg Arg Lys Ala Phe Ser Thr Cys 225 230 235 240Ser Ser His Leu Cys Val Val Gly Leu Phe Phe Gly Ser Ala Ile Val 245 250 255 Thr Tyr Met Ala Pro Lys Ser Arg His Pro Glu Glu Gln Gln Lys Val 260 265 270Leu Ser Leu Phe Tyr Ser Leu Phe Asn Pro Met Leu Asn Pro Leu Ile Tyr Ser Leu Arg Asn Ala Glu Val Lys Gly Ala Leu Arg Arg Ala Leu 290 295 300Arg Lys Glu Arg Leu Thr <210> 218 <211> 927 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (1)..(924) <223> <400> 218 48 atg gcc atg gac aat gtc aca gca gtg ttt cag ttt ctc ctt att ggc Met Ala Met Asp Asn Val Thr Ala Val Phe Gln Phe Leu Leu Ile Gly att tot aac tat oot caa tgg aga gad acg ttt tto aca tta gtg otg Page 124

Ile	Ser	Asn	туг 20	Pro	Gln	Trp	Arg	Asp 25	Thr		PCT Thr			
								ggg Gly						144
								act Thr						192
								tat Tyr						240
								cat His						288
								ttg Leu 105						336
								gac Asp						384
								aat Asn						432
								ctt Leu						480
								ggg Gly						528
								aag Lys 185						576
								acc Thr						624
								tac Tyr						672
								agg Arg						720
								atc Ile						768
								tcc Ser 265						816
								aca Thr						864
								aaa Lys				Val	Met	912
		agg Arg		tga								f	:	927
<210 <211 <212	l> 3	219 308 PRT												

<213> Homo sapiens

<400> 219

Met Ala Met Asp Asn Val Thr Ala Val Phe Gln Phe Leu Leu Ile Gly 1 5 15

lle Ser Asn Tyr Pro Gln Trp Arg Asp Thr Phe Phe Thr Leu Val Leu $20 \hspace{1cm} 25 \hspace{1cm} 30$

lle lle Tyr Leu Ser Thr Leu Leu Gly Asn Gly Phe Met Ile Phe Leu $35 \hspace{1cm} 40 \hspace{1cm} 45$

lle His Phe Asp Pro Asn Leu His Thr Pro Ile Tyr Phe Phe Leu Ser 50 60

Asn Leu Ser Phe Leu Asp Leu Cys Tyr Gly Thr Ala Ser Met Pro Gln 65 70 75 80

Ala Leu Val His Cys Phe Ser Thr His Pro Tyr Leu Ser Tyr Pro Arg 85 90 95

Cys Leu Ala Gln Thr Ser Val Ser Leu Ala Leu Ala Thr Ala Glu Cys $100 \hspace{1.5cm} 105 \hspace{1.5cm} 110 \hspace{1.5cm}$

Leu Leu Leu Ala Ala Met Ala Tyr Asp Arg Val Val Ala Ile Ser Asn 115 120 125

Pro Leu Arg Tyr Ser Val Val Met Asn Gly Pro Val Cys Val Cys Leu 130 135 140

Val Ala Thr Ser Trp Gly Thr Ser Leu Val Leu Thr Ala Met Leu Ile 145 150 160

Leu Ser Leu Arg Leu His Phe Cys Gly Ala Asn Val Ile Asn His Phe 165 170 175

Ala Cys Glu Ile Leu Ser Leu Ile Lys Leu Thr Cys Ser Asp Thr Ser 180 185 190

Leu Asn Glu Phe Met Ile Leu Ile Thr Ser Ile Phe Thr Leu Leu Leu 195 200 205

Pro Phe Gly Phe Val Leu Leu Ser Tyr Ile Arg Ile Ala Met Ala Ile 210 215 220

Ile Arg Ile Arg Ser Leu Gln Gly Arg Leu Lys Ala Phe Thr Thr Cys 225 230 240

Gly Ser His Leu Thr Val Val Thr Ile Phe Tyr Gly Ser Ala Ile Ser 245 250 255

Met Tyr Met Lys Thr Gln Ser Lys Ser Tyr Pro Asp Gln Asp Lys Phe 260 265 270

Ile Ser Val Phe Tyr Gly Ala Leu Thr Pro Met Leu Asn Pro Leu Ile 275 280 285

Tyr Ser Leu Arg Lys Lys Asp Val Lys Arg Ala Ile Arg Lys Val Met 290 295 300

Leu Lys Arg Thr 305
<210> 220 <211> 1008 <212> DNA <213> Homo sapiens
<220> <221> CDS <222> (1)(1005) <223>
<pre><400> 220 atg gaa tca tct ttc tca ttt gga gtg atc ctt gct gtc ctg gcc tcc Met Glu Ser Ser Phe Ser Phe Gly Val Ile Leu Ala Val Leu Ala Ser 1</pre>
ctc atc att gct act aac aca cta gtg gct gtg gct gtg ctg ctg ttg Leu Ile Ile Ala Thr Asn Thr Leu Val Ala Val Ala Val Leu Leu Leu 20 25 30
atc cac aag aat gat ggt gtc agt ctc tgc ttc acc ttg aat ctg gct Ile His Lys Asn Asp Gly Val Ser Leu Cys Phe Thr Leu Asn Leu Ala 35 40 45
gtg gct gac acc ttg att ggt gtg gcc atc tct ggc cta ctc aca gac Val Ala Asp Thr Leu Ile Gly Val Ala Ile Ser Gly Leu Leu Thr Asp 50 55 60
cag ctc tcc agc cct tct cgg ccc aca cag aag acc ctg tgc agc ctg Gln Leu Ser Ser Pro Ser Arg Pro Thr Gln Lys Thr Leu Cys Ser Leu 65 70 75 80
cgg atg gca ttt gtc act tcc tcc gca gct gcc tct gtc ctc acg gtc Arg Met Ala Phe Val Thr Ser Ser Ala Ala Ala Ser Val Leu Thr Val 85 90 95
atg ctg atc acc ttt gac agg tac ctt gcc atc aag cag ccc ttc cgc Met Leu Ile Thr Phe Asp Arg Tyr Leu Ala Ile Lys Gln Pro Phe Arg 100 105 110
tac ttg aag atc atg agt ggg ttc gtg gcc ggg gcc tgc att gcc ggg Tyr Leu Lys Ile Met Ser Gly Phe Val Ala Gly Ala Cys Ile Ala Gly 115 120 125
ctg tgg tta gtg tct tac ctc att ggc ttc ctc cca ctc gga atc ccc Leu Trp Leu Val Ser Tyr Leu Ile Gly Phe Leu Pro Leu Gly Ile Pro 130 135 140
atg ttc cag cag act gcc tac aaa ggg cag tgc agc ttc ttt gct gta Met Phe Gln Gln Thr Ala Tyr Lys Gly Gln Cys Ser Phe Phe Ala Val 145 150 155 160
ttt cac cct cac ttc gtg ctg acc ctc tcc tgc gtt ggc ttc ttc cca . 528 Phe His Pro His Phe Val Leu Thr Leu Ser Cys Val Gly Phe Pro 165 170 175
gcc atg ctc ctc ttt gtc ttc ttc tac tgc gac atg ctc aag att gcc 576 Ala Met Leu Leu Phe Val Phe Phe Tyr Cys Asp Met Leu Lys Ile Ala 180 185 190
tcc atg cac agc cag cag att cga aag atg gaa cat gca gga gcc atg Ser Met His Ser Gln Gln Ile Arg Lys Met Glu His Ala Gly Ala Met 195 200 205
gct gga ggt tat cga tcc cca cgg act ccc agc gac ttc aaa gct ctc Ala Gly Gly Tyr Arg Ser Pro Arg Thr Pro Ser Asp Phe Lys Ala Leu 210 215 220
cgt act gtg tct gtt ctc att ggg agc ttt gct cta tcc tgg acc ccc Arg Thr Val Ser Val Leu Ile Gly Ser Phe Ala Leu Ser Trp Thr Pro 225 230 235 240
ttc ctt atc act ggc att gtg cag gtg gcc tgc cag gag tgt cac ctc Phe Leu Ile Thr Gly Ile Val Gln Val Ala Cys Gln Glu Cys His Leu 245 250 255

										160	200	PCT	FINA	L.ST	25	
tac Tyr	cta Leu	gtg Val	ctg Leu 260	gaa Glu	cgg Arg	tac Tyr	ctg Leu	tgg Trp 265	ctg Leu	ctc Leu	ggc Gly	gtg Val	ggc Gly 270	aac Asn	tcc Ser	816
		aac Asn 275														864
		tac Tyr														912
		ttt Phe														960
		tgt Cys													taa	1008
<21 <21 <21 <21	1 > . 2 >	221 335 PRT Homo	sap:	i ens												
< 40	3> :	221														
Met 1	Glu	Ser	Ser	Phe 5	Ser	Phe	Gly	Val	Ile 10	Leu	Ala	Val	Leu	Ala 15	Ser	
Leu	Ile	lle	Ala 20	Thr	Asn	Thr	Leu	Val 25	Ala	Val	Ala	Val	Leu 30	Leu	Leu	
Ile	His	Lys 35	Asn	Asp	Gl y	Val	Ser 40	Leu	Cys	Phe	Thr	Leu 45	Asn	Leu	Ala	
Val	Ala 50	Asp	Thr	Leu	Ile	Gly 55	Val	Ala	Ile	Ser	Gly 60	Leu	Leu	Thr	Asp	
Gln 65	Leu	Ser	Ser	Pro	Ser 70	Arg	Pro	Thr	Gln	Lys 75	Thr	Leu	Cys	Ser	Leu 80	
Arg	Met	Ala	Phe	Val 85	Thr	Ser	Ser	Ala	Ala 90	Ala	Ser	Val	Leu	Thr 95	Val	
Met	Leu	lle	Thr 100		Asp	Arg		Leu 105		Ile	Lys		Pro 110		Arg	
Туг	Leu	Lys 115	Ile	Met	Ser	Gly	Phe 120	Val	Ala	Gly	Ala	Cys 125	Ile	Ala	Gly	
Leu	Trp 130	Leu	Val	Ser	туr	Leu 135	Ile	Gly	Phe	Leu	Pro 140	Leu	Gly	Ile	Pro	
Met 145	Phe	Gln	Gln	Thr	Ala 150	Tyr	Lys	Gly	Gln	Cys 155	Ser	Phe	Phe	Ala	Val 160	
Phe	His	Pro	His	Phe 165	Val	Leu	Thr	Leu	Ser 170	Cys	Val	Gly	Phe	Phe 175		÷
Ala	Met	Leu	Leu 180	Phe	Val	Phe	Phe	Туг 185	Cys	Asp	Met	Leu	Lys 190	Ile	Ala	

Ser Met His Ser Gln Gln Ile Arg Lys Met Glu His Ala Gly Ala Met 195 200 205

Ala	Gly 210		Тyr	Arg	Ser	Pro 215	Arg	Thr	Pro	Ser	Asp 220		Lys	Ala	Leu	
Arg 225	Thr	· Val	Ser	Val	Leu 230		Gly	Ser	Phe	Ala 235		Ser	Trp	Thr	Pro 240	
Phe	Leu	Ile	Thr	Gly 245	Ile	Val	Gln	Val	Ala 250		Gln	Glu	Cys	His 255	Leu	
Tyr	Leu	Val	Leu 260		Arg	Туr	Leu	Trp 265	Leu	Leu	Gly	Val	Gly 270		Ser	
Leu	Leu	Asn 275		Leu	Ile	Туr	Ala 280	туг	Trp	Gln	Lys	Glu 285	Val	Arg	Leu	
Gln	Leu 290		His	Met	Ala	Leu 295	Gly	Val	Lys	Lys	Val 300	Leu	Thr	Ser	Phe	
Leu 305		Phe	Leu	Ser	Ala 310	Arg	Asn	Cys	Gly	Pro 315	Glu	Arg	Pro	Arg	Glu 320	
Ser	Ser	Cys	His	11e 325	Val	Thr	Ile	Ser	Ser 330	Ser	Glu	Phe	Asp	Gly 335		
<21 <21 <21 <21	1> 2>	222 975 DNA Homo	sap	iens												
<22 <22 <22 <22	1> 2>	CDS (1).	. (97	2)												
	cgg										ctc Leu					48
											act Thr					96
gcc Ala	gcc Ala	gag Glu 35	gca Ala	gct Ala	ggg Gly	gcg Ala	cca Pro 40	gac Asp	cgc Arg	gtg Val	ggc Gly	tct Ser 45	tta Leu	ttt Phe	gtt Val	144
											agt Ser 60					192
65											ata Ile					240
65 999	Phe	Gln	Met ggc	Leu cca	Val 70 aaa	Val ggt	Ile caa	Pro cga	Cys	Leu 75 ttc	ata	Tyr att	Arg	Lys aga	Thr 80 gga	240
65 ggg Gly gtc	Phe ttt Phe ctt	Gln ata Ile ggt	Met ggc Gly tct	cca Pro 85	Val 70 aaa Lys gcc	Val ggt Gly atg	lle caa Gln atg	Pro cga Arg	Cys att ile 90 ata	Leu 75 ttc Phe	ata Ile ctc	Tyr att Ile gct	arg ctc Leu	Lys aga Arg 95 cag	Thr 80 gga Gly aca	
ggg Gly gtc Val	Phe ttt Phe ctt Leu tcc	Gln ata Ile ggt Gly	Met ggc Gly tct Ser 100 gct	Leu cca Pro 85 acc Thr	Val 70 aaa Lys gcc Ala	yal ggt Gly atg Met	lle caa Gln atg Met	Pro cga Arg ctt Leu 105	Cys att Ile 90 ata Ile acg	Leu 75 ttc Phe tac Tyr	ata Ile ctc Leu	Tyr att Ile gct Ala	ctc Leu tac Tyr 110	Lys aga Arg 95 cag Gln	Thr 80 gga Gly aca Thr	288

	130					135				1 6U	200 140	PCT	FINA	L.SI	25		
	gct					ttc		atc Ile			gtg					4 8	80
								gac Asp								52	28
								ttc Phe 185								51	76
								cta Leu								62	24
								tat Tyr								67	72
								gga Gly								72	20
								ttc Phe								7 (68
								ctt Leu 265								81	16
								gtg Val								86	64
								acg Thr								91	12
								gga Gly								96	60
	-	tcc Ser	aaa Lys	tga												97	75
<210 <211 <211 <211	1> . ?> !	223 324 PRT Homo	sapi	ens													
< 400)> 2	223															
Met 1	Arg	Pro	Gln	Asp 5	Ser	Thr	Gly	Val	Ala 10	Glu	Leu	Gln	Glu	Pro 15	Gly		
Leu	Pro	Leu	Thr 20	Asp	Asp	Ala	Pro	Pro 25	Gly	Ala	Thr	Glu	Glu 30	Pro	Ala		
Ala	Ala	Glu 35	Ala	Ala	Gly	Ala	Pro 40	Asp	Arg	Val	Gly	Ser 45	Leu	Phe	Val		
Lys	Lys 50	Val	Gln	Asp	Val	His 55	Ala	Val	Glu	lle	Ser 60	Ala	Phe		,Cys	·	
Val 65	Phe	Gln	Met	Leu	Val 70	Val	Ile	Pro	Cys	Leu 75	Ile	Туг	Arg	Lys	Thr 80		
Gly	Phe	Ile	Gly	Pro	Lys	G] A	Gln	Arg	Ile	Phe		lle age		Arg	Gly		

48

96

				85					90	160	200	PCT	FIN	AL.S' 95	T25	
Val	Leı	g Gly	Ser 100	Thr	Ala	Met	Met	Leu 105		Tyr	Tyr	Ala	Туг 110		Thr	
Met	Ser	Leu 115		Asp	Ala	Thr	Val 120	Ile	Thr	Phe	Ser	Ser 125	Pro	Val	Phe	
Thr	Ser 130	lle	Phe	Ala	Trp	Ile 135	Cys	Leu	Lys	Glu	Lys 140		Ser	Pro	Trp	
Asp 145	Ala	Leu	Phe	Thr	Val 150	Phe	Thr	Ile	Thr	Gly 155	Val	Ile	Leu	Ile	Val 160	
Arg	Pro	Pro	Phe	Leu 165	Phe	Gly	Ser	Asp	Thr 170	Ser	Gly	Met	Glu	Glu 175	Ser	
туr	Ser	Gly	His 180	Leu	Lys	Gly	Thr	Phe 185	Ala	Ala	Ile	Gly	Ser 190	Ala	Val	
Phe	Ala	Ala 195	Ser	Thr	Leu	Val	Ile 200	Leu	Arg	Lys	Met	Gly 205	Lys	Ser	Val	
Asp	Tyr 210	Phe	Leu	Ser	Ile	Trp 215	Tyr	Туr	Val	Val	Leu 220	Gly	Leu	Val	Glu	
Ser 225	Val	Ile	Ile	Leu	Ser 230	Val	Leu	Gly	Glu	Trp 235	Ser	Leu	Pro	Tyr	Cys 240	
Gly	Leu	Asp	Arg	Leu 245	Phe	Leu	Ile	Phe	Ile 250	Gly	Leu	Phe	Gly	Leu 255	Gly	
Gly	Gl n	Ile	Phe 260	Ile	Thr	Lys	Ala	Leu 265	Gln	Ile	Glu	Lys	Ala 270	Gly	Pro	
Val	Ala	11e 275	Met	Lys	Thr	Met	Asp 280	Val	Val	Phe	Ala	Phe 285	Ile	Phe	Gln	
Ile	11e 290	Phe	Phe	Asn	Asn	Val 295	Pro	Thr	Trp	Trp	Thr 300	Val	Gly	Gly	Ala	
Leu 305	Cys	Val	Val	Ala	Ser 310	Asn	Val	Gly	Ala	Ala 315	Ile	Arg	Lys	Trp	Туг 320	
Gln	Ser	Ser	Lys							,						
<210 <211 <212 <213	> !>	224 876 DNA Homo	sapi	ens												
<220 <221 <222 <223	>	CDS (1)	(873	1)										,	:	·
	tac	224 aac Asn														
atc	cct	gga	ctt	gag	cag	tac	cac	gtc	tgg	atc	agc	atc	cca	ttc	tgc	

Ile	Pro	Gly	Leu 20	Glu	Gln	Туг	His	Val 25	Trp		FINA Pro 30		
		tat Tyr 35											144
		gta Val											192
		gcc Ala											240
		agc Ser											288
		ttt Phe											336
		ctg Leu 115											384
		agt Ser											432
		ttc Phe											480
		gtt Val											528
		ttt Phe											576
		gtc Val 195											624
		gat Asp											672
		atc Ile											720
		ttt Phe											768
		tat Tyr											816
		acc Thr 275											864
		cag Gln	tga										876
<210 <211 <212 <213	l> ?>	225 291 PRT Homo	sapi	iens								,	
<400)>	225											

Met Tyr Asn Met Ser Asp His Gly Thr Gly Leu Phe Ile Leu Leu Gly 1 5 10 15

Ile Pro Gly Leu Glu Gln Tyr His Val Trp Ile Ser Ile Pro Phe Cys $20 \hspace{1cm} 25 \hspace{1cm} 30$

Leu Ile Tyr Leu Met Ala Val Val Ala As
n Ser Ile Leu Leu Tyr Leu $35 \ 40 \ 45$

Ile Val Val Glu His Ser Leu His Ala Pro Met Phe Phe Leu Ser 50 55 60

Met Leu Ala Ile Thr Asp Leu Ile Leu Ser Thr Thr Cys Val Pro Lys 65 70 75 80

Thr Leu Ser Ile Phe Cys Phe Val Leu Asp Ser Ala Ile Leu Leu Ala 85 90 95

Met Ala Phe Asp Arg Tyr Met Ala Ile Cys Ser Pro Leu Arg Tyr Thr $100 \hspace{1cm} 105 \hspace{1cm} 110$

Thr Ile Leu Thr Pro Lys Thr Ile Val Lys Ile Ala Val Gly Ile Cys 115 120 125

Phe Arg Ser Phe Cys Val Phe Val Pro Cys Val Phe Leu Val Asn Arg 130 135 140

Leu Pro Phe Cys Arg Thr His Ile Ile Ser His Thr Tyr Cys Glu His 145 150 155 160

Ile Gly Val Ala Gln Leu Ala Cys Ala Asp Ile Ser Ile Asn Ile Trp 165 170 175

Cys Gly Phe Cys Val Pro Ile Met Thr Val Met Thr Asp Val Ile Leu 180 185 190

Ile Ala Val Ser Tyr Thr Leu Ile Leu Cys Ala Val Phe Cys Leu Pro 195 200 205

Ser Gln Asp Ala Arg Gln Lys Ala Leu Cys Ser Cys Gly Ser His Val 210 215 220

Cys Val Ile Leu Ile Phe Tyr Ile Pro Ala Phe Phe Ser Ile Leu Ala 225 230 235 240

His Cys Phe Gly His Asn Val Pro His Thr Phe His Ile Met Phe Ala 245 250 255

Asn Leu Tyr Val Ile Ile Pro Pro Ala Leu Asn Ser Ile Val Tyr Arg 260 265 270

Ile Lys Thr Lys Gln Ile Gln Asn Arg Ile Leu Leu Leu Phe Pro Lys 275 280 285

Gly Ser Gln 290

<210> 226 <211> 1949 <212> DNA

<21	.3>	Homo	sap	iens	;					160	200	PCT	FIN	AL.S	T25	
<22 <22 <22 <22	?1> ?2>	CDS (430))(1164)											
< 40 aga		226 jcgg	gact	tctc	ca a	gtca	agge	c ao	atct	cttc	cct	acto	aat	acta	tgttcc	60
															gacatt	
															ccagac	
ctt	ttct	ccg	tgag	ttcc	tc a	gcca	ggac	t gc	tgcc	atgc	cgg	tgac	tgt	tacc	caccca	240
acc	gtca	cga	ccac	catg	cg g	tece	ccac	c gt	cgta	gggt	cct	ctag	ggc	cctg	atccag	300
ccc	ctgg	gcc	tcct	ccgc	ct g	ctgc	agct	g gt	gtcc	acct	gcg	tggc	ctt	gtca	ctggtg	360
															tgtttc	
tgc	tttg	cc a M 1	et T	cc c hr L	tg g eu V	tc a al I 5	tc c le L	tc c eu L	tc g eu V	tg g al G	lu L	tg g eu G 0	gc g ly G	gc t ly S	cc cag er Gln	471
gcc Ala 15	cgc Arg	ttc Phe	ccc Pro	ttg Leu	ttt Phe 20	tgg Trp	cgc Arg	aac Asn	ttc Phe	ccc Pro 25	atc Ile	acc Thr	ttt Phe	gcc Ala	tgc Cys 30	519
tat Tyr	gcg Ala	gcc Ala	ctc Leu	ttg Leu 35	tgc Cys	ctc Leu	tcg Ser	gcc Ala	tcc Ser 40	atc Ile	atc Ile	tac Tyr	ccc Pro	acc Thr 45	acc Thr	567
tac Tyr	ttg Leu	cag Gln	ttc Phe 50	ctg Leu	tcc Ser	cac His	ggc Gly	cgt Arg 55	tcc Ser	cgc Arg	gac Asp	cac His	gcc Ala 60	atc Ile	gcc Ala	615
gcc Ala	atc Ile	gtc Val 65	ttc Phe	tct Ser	ggc Gly	atc Ile	gcc Ala 70	tgt Cys	gtg Val	gct Ala	tac Tyr	gcc Ala 75	acc Thr	gaa Glu	gta Val	663
acc Thr	tgg Trp 80	acc Thr	cgg	gcc Ala	cgg Arg	ccc Pro 85	ggc Gly	gag Glu	atc Ile	act Thr	gac Asp 90	tac Tyr	atg Met	gcc Ala	tcc Ser	711
gag Glu 95	ctg Leu	д9д Glу	ctg Leu	ctg Leu	aag Lys 100	gtg Val	ctg Leu	gag Glu	acc Thr	ttc Phe 105	gtg Val	gcc Ala	tgc Cys	ctc Leu	atc Ile 110	759
ttc Phe	gtg Val	ttc Phe	atc Ile	aat Asn 115	agc Ser	ccc Pro	tac Tyr	gtg Val	tac Tyr 120	cac His	aac Asn	cgg Arg	ccg Pro	gcc Ala 125	ctg Leu	807
gag Glu	tgg Trp	tgg Trp	gtg Val 130	gcy Ala	gtg Val	tac Tyr	gcc Ala	ctc Leu 135	tgc Cys	ttc Phe	gtc Val	ctg Leu	gcg Ala 140	gcc Ala	ctc Leu	855
act Thr	atc Ile	ctg Leu 145	ctg Leu	agc Ser	ctg Leu	ggg Gly	cac His 150	t <i>gc</i> Cys	acc Thr	aac Asn	atg Met	ctg Leu 155	ccc Pro	atc Ile	cgc Arg	903
ttc Phe	ccc Pro 160	agt Ser	ttc Phe	ctg Leu	ttg Leu	ggg Gly 165	ctg Leu	gcc Ala	ttg Leu	ctg Leu	tcc Ser 170	gtc Val	ctc Leu	ctc Leu	tat Tyr	951
gcc Ala 175	acț Thr	gcc Ala	ctt Leu	gtc Val	ct <i>c</i> Leu 180	tgg Trp	ccc Pro	ct <i>c</i> Leu	tac Tyr	cag Gln 185	ttc Phe	aac Asn	gag Glu	aag Lys	tat Tyr 190	999
ggt Gly	gtc Val	cag Gln	ccc Pro	tgg Trp 195	cag Gln	acg Thr	aga Arg	gat Asp	gtg Val 200	agc Ser	tgc Cys	agc Ser	gac Asp	aga Arg 205	aac Asn	1047
ccc Pro	tac Tyr	ctt Leu	gtg Val 210	tgt Cys	atc Ile	tgg Trp	gac Asp	cgc Arg 215	cga Arg	ctg Leu	gct Ala	gtg Val	acc Thr 220	aac Asn	ctg Leu	1095
acg	gcc	gtc	aac	ttg	ctg	gcc	tat	gtg	ggc	gac	ctg	gtg	tac	tct	gcc	1143
											P	age	134			

Thr Ala Val Asn Leu Leu Ala Tyr Val Gly Asp Leu Val Tyr Ser Ala 225 230 235	
cac ctg gtt ttt gtc aag gtc taagactccc aaagggcccc gtttgcctct His Leu Val Phe Val Lys Val 240 245	1194
ccaacctctt catcctgccc ccgctgagtt ttctttattg agtattcatt tcctgggttt	1254
tcctcttccc tatctcccct cctccccttt tctttccttc ccaattcatc gcactttccc	1314
agttctctga tgtatgttct tccctttcct ctgctgtttc cttcttgttt tgttctgttg	1374
cccacaacct gttttcaccc gtttctcttt ttccactctc tcttttgttt ctttcctctc	1434
aattettte taggttteet gttggttte ttatetgeet attteeegae catettetee	1494
tatttcctgg ggagccctga ggcttttctt ctcctgcccc caagcacctc cagcggtgat	1554
gagetecaca eccecacace cattgeaget gtggegeeae gteeteceaa ggggeettet	1614
gcccgccccc gccctagctg tgccttagtc agtgtgtact tgtgtgtgtt tgggggagtg	1674
ggaattgggc cccctttctc ccagtggagg aaggtgtgct gtgcacctcc cctttaaatt	1734
aaaaaaaatg tatgtatctc tggaagtcaa taatttccag tgagcgggag gcttcaagcg	1794
cagaccetgg gteectagae etegeetage actetgeett gecagagatt ggetecagaa	1854
tttgtgccag acttacagaa aacccactgc ctagaggcca tcttaaagga agcaatggat	1914
ggatcccttt catcccaact gttcttcgcg gtatc	1949
<210> 227 <211> 245 <212> PRT <213> Homo sapiens	
<400> 227	
Met Thr Leu Val Ile Leu Leu Val Glu Leu Gly Gly Ser Gln Ala Arg 1 5 10 15	
Phe Pro Leu Phe Trp Arg Asn Phe Pro Ile Thr Phe Ala Cys Tyr Ala 20 25 30	
Ala Leu Leu Cys Leu Ser Ala Ser Ile Ile Tyr Pro Thr Thr Tyr Leu 35 40 45	
Gln Phe Leu Ser His Gly Arg Ser Arg Asp His Ala Ile Ala Ala Ile 50 55 60	
Val Phe Ser Gly Ile Ala Cys Val Ala Tyr Ala Thr Glu Val Thr Trp 65 70 75 80	
Thr Arg Ala Arg Pro Gly Glu Ile Thr Asp Tyr Met Ala Ser Glu Leu 85 90 95	
Gly Leu Leu Lys Val Leu Glu Thr Phe Val Ala Cys Leu Ile Phe Val	
Phe Ile Asn Ser Pro Tyr Val Tyr His Asn Arg Pro Ala Leu Glu Trp 115 120 125	
Trp Val Ala Val Tyr Ala Leu Cys Phe Val Leu Ala Ala Leu Thr Ile 130 135 140	
Leu Leu Ser Leu Gly His Cys Thr Asn Met Leu Pro Ile Arg Phe Pro 145 150 155 160	
Page 135	

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Ser Phe Leu Leu Gly Leu Ala Leu Leu Ser Val Leu Leu Tyr Ala Thr 165 170 175	
Ala Leu Val Leu Trp Pro Leu Tyr Gln Phe Asn Glu Lys Tyr Gly Val 180 185 190	
Gln Pro Trp Gln Thr Arg Asp Val Ser Cys Ser Asp Arg Asn Pro Tyr 195 200 205	
Leu Val Cys Ile Trp Asp Arg Arg Leu Ala Val Thr Asn Leu Thr Ala 210 215 220	
Val Asn Leu Leu Ala Tyr Val Gly Asp Leu Val Tyr Ser Ala His Leu 225 230 235 240	
Val Phe Val Lys Val 245	
<210> 228 <211> 2980 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (213)(2336) <223>	
<400> 228 cccagagacc caggccgcgg aactggcagg cgtttcagag cgtcagaggc tgcggatgag	60
cagacttgga ggactccagg ccagagacta ggctgggcga agagtcgagc gtgaaggggg	120
	120 1 80
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg	
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1 5	180
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1 5 gcc ctc ctc gtc ctc ctc ttc tgc ttc aga ggg aga gca ggc ccg tcg Ala Leu Leu Val Leu Phe Cys Phe Arg Gly Arg Ala Gly Pro Ser 10 15 20	180 233
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1 5 gcc ctc ctc gtc ctc ctc ttc tgc ttc aga ggg aga gca ggc ccg tcg Ala Leu Leu Val Leu Leu Phe Cys Phe Arg Gly Arg Ala Gly Pro Ser 10 20 ccc cat ttc ctg caa cag cca gag gac ctg gtg gtg ctg ctg ggg gag Pro His Phe Leu Gln Gln Pro Glu Asp Leu Val Val Leu Leu Gly Glu 25 30 35	180 233 281
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1 5 gcc ctc ctc gtc ctc ctc ttc tgc ttc aga ggg aga gca ggc ccg tcg Ala Leu Leu Val Leu Leu Phe Cys Phe Arg Gly Arg Ala Gly Pro Ser 10 15 20 ccc cat ttc ctg caa cag cca gag gac ctg gtg gtg ctg ctg ggg gag Pro His Phe Leu Gln Gln Pro Glu Asp Leu Val Val Leu Leu Gly Glu 25 30 35 gaa gcc cgg ctg ccg tgt gct ctg ggc gcc tac tgg ggg cta gtt cag Glu Ala Arg Leu Pro Cys Ala Leu Gly Ala Tyr Trp Gly Leu Val Gln 45 50 55	180 233 281
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1	180 233 281 329
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1	180 2233 2281 3329 3377
ctccgggcca gggtgacagg aggcgtgctt gagaggaaga agttgacggg aaggccagtg cgacggcaaa tctcgtgaac cttgggggac ga atg ctc agg atg cgg gtc ccc Met Leu Arg Met Arg Val Pro 1	180 233 281 329 377 425

										tgt	cgg	agc	cgt	AL.ST 999 Gly 150	gat	665
														ctg Leu		713
														cct Pro		761
														gat Asp		809
														gga Gly		857
														act Thr 230		905
	-	_										-		t t c Phe	-	953
														gca Ala		1001
														gtc Val		1049
														agc Ser		1097
														ttt Phe 310		1145
														ggg Gly		1193
														cgg Arg		1241
acc Thr	tgg Trp 345	Thr	cgc Arg	Arg	ggt Gly	Gly	Ala	cag Gln	Val	Leu	Gly	tct Ser	gga Gly	gcc Ala	aca Thr	1289
														gtg Val		1337
														gag Glu 390		1385
														tct Ser		1433
														ttc Phe		1481
														ttc [.] Phe		-1529
														gcc Ala		1577

														cac His 470		1625
														agt Ser		1673
														cgt Arg		1721
gac Asp	ttg Leu 505	ctg Leu	ccc Pro	act Thr	gtg Val	cgg Arg 510	ata Ile	gtg Val	gcc Ala	gga Gly	gtg Val 515	gcc Ala	gct Ala	gcc Ala	acc Thr	1769
														tgg Trp		1817
														atg Met 550		1865
														gaa Glu		1913
														gac Asp		1961
														gac Asp		2009
														agc Ser		2057
														ccc Pro 630		2105
														ctg Leu		2153
														ctc Leu		2201
														tcc Ser		2249
														gct Ala		2297
			cct Pro										tgad	catci	tt	2346
cca	atgga	aag a	agtc	ctgg	ga to	tcca	actt	gco	cataa	atgg	atte	gttct	tga t	ttct	gagga	2406
gcc	aggad	caa q	gttg	gcgad	c tt	acto	ctc	aaa	acto	gaac	acaa	gggg	gag q	ggaaa	agatca	2466
tta	catt	tgt (cagga	agcat	t tç	gtata	cagt	caç	gctca	gcc	aaaq	gaga	atg o	ccca	aagtgg	2526
gage	caaca	atg q	gccad	ccaa	at at	gcc	cacct	att	ccc	ggt	gtaa	aaga	aga 1	tca	agatgg	2586
cag	gtago	gec d	ettt	gagga	ag ag	gatgo	gggad	ago	ggcaq	gtgg	gtgi	tgg	gag t	ttg	ggccg	2646
gga	ggaa	agt 1	tgtti	tcta	go da	ectga	aaga	a aga	atatt	tca	agat	gaco	cat o	ctgca	attgag	2706
agga	aaag	gta q	gcata	aggat	ta ga	itgaa	gato	g aag	gagca	atac	cag	geced	cac o	ctg	gctctc	2766

Page 138

16U 200 PCT FINAL.ST25 cctgagggga actttgctcg gccaatggaa atgcagccaa gatggccata tactccctag	2026
gaacccaaga tggccaccat cttgatttta ctttccttaa agactcagaa agacttggac	2826
ccaaggagtg gggatacagt gagaattacc actgttgggg caaaatattg ggataaaaat	2886 2946
atttatgttt aataataaaa aaaagtcaaa gagg	
<210> 229 <211> 708 <212> PRT	2980
<213> Homo sapiens	
<400> 229	
Met Leu Arg Met Arg Val Pro Ala Leu Leu Val Leu Leu Phe Cys Phe 1 10 15	
Arg Gly Arg Ala Gly Pro Ser Pro His Phe Leu Gln Gln Pro Glu Asp 20 25 30	
Leu Val Val Leu Leu Gly Glu Glu Ala Arg Leu Pro Cys Ala Leu Gly 35 40 45	
Ala Tyr Trp Gly Leu Val Gln Trp Thr Lys Ser Gly Leu Ala Leu Gly 50 60	
Gly Gln Arg Asp Leu Pro Gly Trp Ser Arg Tyr Trp Ile Ser Gly Asn 65 70 75 80	
Ala Ala Asn Gly Gln His Asp Leu His Ile Arg Pro Val Glu Leu Glu 85 90 95	
Asp Glu Ala Ser Tyr Glu Cys Gln Ala Thr Gln Ala Gly Leu Arg Ser 100 105 110	
Arg Pro Ala Gln Leu His Val Leu Val Pro Pro Glu Ala Pro Gln Val 115 120 125	
Leu Gly Gly Pro Ser Val Ser Leu Val Ala Gly Val Pro Ala Asn Leu 130 135 140	
Thr Cys Arg Ser Arg Gly Asp Ala Arg Pro Thr Pro Glu Leu Leu Trp 145 150 155 160	
Phe Arg Asp Gly Val Leu Leu Asp Gly Ala Thr Phe His Gln Thr Leu 165 170 175	
Leu Lys Glu Gly Thr Pro Gly Ser Val Glu Ser Thr Leu Thr 180 185 190	
Pro Phe Ser His Asp Asp Gly Ala Thr Phe Val Cys Arg Ala Arg Ser 195 200 205	
Gln Ala Leu Pro Thr Gly Arg Asp Thr Ala Ile Thr Leu Ser Leu Gln 210 215 220	
Tyr Pro Pro Glu Val Thr Leu Ser Ala Ser Pro His Thr Val Glu 225 230 235 240	
Gly Glu Lys Val Ile Phe Leu Cys Gln Ala Thr Ala Gln Pro Pro Val 245 250 255	

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Thr Gly Tyr Arg Trp Ala Lys Gly Gly Ser Pro Val Leu Gly Ala Arg 260 265 270

Gly Pro Arg Leu Glu Val Val Ala Asp Ala Ser Phe Leu Thr Glu Pro 275 280 285

Val Ser Cys Glu Val Ser Asn Ala Val Gly Ser Ala Asn Arg Ser Thr 290 295 300

Ala Leu Asp Val Leu Phe Gly Pro Ile Leu Gln Ala Lys Pro Glu Pro 305 310 315 320

Val Ser Val Asp Val Gly Glu Asp Ala Ser Phe Ser Cys Ala Trp Arg 325 330 335

Gly Asn Pro Leu Pro Arg Val Thr Trp Thr Arg Arg Gly Gly Ala Gln 340 345 350

Val Leu Gly Ser Gly Ala Thr Leu Arg Leu Pro Ser Val Gly Pro Glu 355 360 365

Asp Ala Gly Asp Tyr Val Cys Arg Ala Glu Ala Gly Leu Ser Gly Leu 370 380

Arg Gly Gly Ala Ala Glu Ala Arg Leu Thr Val Asn Ala Pro Pro Val 385 390395500

Val Thr Ala Leu His Ser Ala Pro Ala Phe Leu Arg Gly Pro Ala Arg 405 410 415

Leu Gln Cys Leu Val Phe Ala Ser Pro Ala Pro Asp Ala Val Val Trp 420 425 430

Ser Trp Asp Glu Gly Phe Leu Glu Ala Gly Ser Gln Gly Arg Phe Leu 435 440 445

Val Glu Thr Phe Pro Ala Pro Glu Ser Arg Gly Gly Leu Gly Pro Gly 450 455 460

Leu Ile Ser Val Leu His 1le Ser Gly Thr Gln Glu Ser Asp Phe Ser 465 470 475 480

Arg Ser Phe Asn Cys Ser Ala Arg Asn Arg Leu Gly Glu Gly Gla Ala 405 490 495

Gln Ala Ser Leu Gly Arg Arg Asp Leu Leu Pro Thr Val Arg Ile Val 500 505 510

Ala Gly Val Ala Ala Ala Thr Thr Thr Leu Leu Met Val Ile Thr Gly 515 520 525

Val Ala Leu Cys Cys Trp Arg His Ser Lys Ala Ser Ala Ser Phe Ser 530 535 540

Glu Gln Lys Asn Leu Met Arg Ile Pro Gly Ser Ser Asp Gly Ser Ser 545 550 555 560

Ser Arg Gly Pro Glu Glu Glu Glu Thr Gly Ser Arg Glu Asp Arg Gly
565 570 575

Pro Ile Val His Thr Asp His Ser Asp Leu Val Leu Glu Glu Lys Gly 585

Thr Leu Glu Thr Lys Asp Pro Thr Asn Gly Tyr Tyr Lys Val Arg Gly

Val Ser Val Ser Leu Ser Leu Gly Glu Ala Pro Gly Gly Gly Leu Phe

Leu Pro Pro Pro Ser Pro Leu Gly Pro Pro Gly Thr Pro Thr Phe Tyr

Asp Phe Asn Pro His Leu Gly Met Val Pro Pro Cys Arg Leu Tyr Arg

Ala Arg Ala Gly Tyr Leu Thr Thr Pro His Pro Arg Ala Phe Thr Ser

Tyr Ile Lys Pro Thr Ser Phe Gly Pro Pro Asp Leu Ala Pro Gly Thr

Pro Pro Phe Pro Tyr Ala Ala Phe Pro Thr Pro Ser His Pro Arg Leu

Gln Thr His Val 705

<210> 230

5188 <211> <212> DNA

<213> Homo sapiens

<220>

<221> <222> CDS

(887)..(2974)

<223>

<400> 230

egegetetet tecteetea gacaactege eeeeegeest eegeeeest eeaegtaatt 60 ccgaaagagc agaagaaaga gaaggagaac aggaaaagaa gagctagtaa gcgagagcga 120 gagCacagaa aagaaaaaa aaagccttaa gaggaccgaa ggggaggaaa ggaaaaggat 180 ggacaaccac aaaacgcagc gattgcggaa attttccagc gccattggct gggcagcgtg 240 agtccttcgg tcgggcgtga tttcagcacc gggggaactg gacagcacct cggggggact 300 totgggcaac cogcaaccac agcaagaact ccaccagcag cotcaacaac agaagcogcg 360 420 gaaaaccctg ctttgtatca gagaggcaag gtcagtccga cgcacagcca tgcacaggca gtgcgcctgt actacgctgc aaaccctctg cttgtttctc taacatgcac ttgcttctaa 480 ttactagcat tgtttcattt ctgatcagtg aagatcagta gatgagattc tgtaagggtg 540 tacttttaat ttatatgtat atatttaact tctttttctg ttattttaa agtgttgtgg 600 gggagtgggg ttttttcct acttttttt ttttttttt ttctttgctt gccttgcact 660 720 acgtgcctgg atagtttgtg gatataatta ttgactggcg tctgggctat tgcagtgcgg gggggttagg gaggaaggaa tccaccccca ccccccaaa ccctttctt ctcctttcct 780 ggcttcggac attggagcac taaatgaact tgaattgtgt ctgtggcgag caggatggtc 840 895 gctgttactt tgtgatgaga tcggggatga attgctcgct ttaaaa atg ctg ctt Met Leu Leu

												•	•			
					gag Glu											943
					aaa Lys 25											991
					gac Asp											1039
					act Thr											1087
					ctt Leu											1135
					atg Met											1183
					ctg Leu 105											1231
					ttt Phe											1279
					gct Ala											1327
					ttg Leu											1375
					cta Leu											1423
					cgg Arg 185											1471
					atc Ile											1519
					acc Thr											1567
gaa Glu	aac Asn	att Ile 230	ccc Pro	aag Lys	aat Asn	gcc Ala	ctg Leu 235	atc Ile	ggc Gly	cga Arg	gtg Val	gtc Val 240	tgc Cys	gaa Glu	gcc Ala	1615
					ggt Gly											1663
					aac Asn 265											1711
					ttt Phe										Lys	1759
					gat Asp											1807
ggt	aca	aag	atc	cca	ggc	aac	tgg	cag	atc	aaa	atc	aga	ccc	aca	gca	1855

Gly	Thr	Lys 310		Pro	Gly	Asn	Trp 315	Gln	Ile					AL.S. Thr		
												Ala		agt Ser		1903
ccc Pro 340	Cys	cct Pro	ggg Gly	ggc G1y	tgc Cys 345	agc Ser	tgc Cys	gac Asp	cac His	atc Ile 350	Pro	G1 y	tcg Ser	ggt Gly	tta Leu 355	1951
aag Lys	atg Met	aac Asn	tgc Cys	aac Asn 360	aac Asn	agg Arg	aac Asn	gtg Val	agc Ser 365	agc Ser	ttg Leu	gct Ala	gat Asp	ttg Leu 370	aag Lys	1999
														aag Lys		2047
														att Ile		2095
														act Thr		2143
aag Lys 420	aac Asn	ctt Leu	ttg Leu	gac Asp	ctc Leu 425	agg Arg	tgg Trp	cta Leu	tac Tyr	atg Met 430	gat Asp	agc Ser	aat Asn	tac Tyr	ctg Leu 435	2191
														gag Glu 450		2239
														act Thr		2287
														ctg Leu		2335
agg Arg	tcc Ser 485	ctg Leu	cct Pro	gtg Val	gac Asp	gtg Val 490	ttc Phe	gct Ala	ggg Gly	gtc Val	tcg Ser 495	ctc Leu	tct Ser	aaa Lys	ctc Leu	2383
														gtg Val		2431
gac Asp	cag Gln	tta Leu	acc Thr	tcc Ser 520	atc Ile	atc Ile	cag Gln	ata Ile	gac Asp 525	ctc Leu	cac His	gga Gly	aac Asn	ccc Pro 530	tgg Trp	2479
														cgc Arg		2527
ggt Gly	tcc Ser	gaa Glu 550	gtg Val	ctg Leu	atg Met	agc Ser	gac Asp 555	ctc Leu	aag Lys	tgt Cys	gag Glu	acg Thr 560	ccg Pro	gtg Val	aac Asn	2575
														tgc Cys		2623
cag Gln 580	ctg Leu	tac Tyr	gct Ala	agg Arg	atc Ile 585	tcg Ser	ccc Pro	acg Thr	tta Leu	act Thr 590	tcg Ser	cac His	agt Ser	aaa Lys	aac Asn 595	2671
agc Ser	act Thr	GJ À āāā	ttg Leu	gcg Ala 600	gag Glu	acc Thr	ggg Gly	acg Thr	cac His 605	tcc Ser	aac Asn	tcc Ser	tac Tyr	cta' Leu 610	gac Asp	2719
														ctg Leu		2767

Page 143

16U 200 PCT FINAL.ST25 ttt gtc acc tcc gcc ttc acc gtg gtg ggc atg ctc gtg ttt atc ctg Phe Val Thr Ser Ala Phe Thr Val Val Gly Met Leu Val Phe Ile Leu 630 635 640	2815
agg aac cga aag cgg tcc aag aga cga gat gcc aac tcc tcc gcg tcc Arg Asn Arg Lys Arg Ser Lys Arg Arg Asp Ala Asn Ser Ser Ala Ser 645 650 655	2863
gag att aat tcc cta cag aca gtc tgt gac tct tcc tac tgg cac aat Glu lle Asn Ser Leu Gln Thr Val Cys Asp Ser Ser Tyr Trp His Asn 660 665 670 675	2911
ggg cct tac aac gca gat ggg gcc cac aga gtg tat gac tgt ggc tct Gly Pro Tyr Asn Ala Asp Gly Ala His Arg Val Tyr Asp Cys Gly Ser 680 685 690	2959
cac tog oto toa gao taagacooca accocaatag gggagggcag agggaaggog His Ser Leu Ser Asp 695	3014
atacatcett ecceacegea ggeaceeegg gggetggagg ggegtgtace caaateeeeg	3074
cgccatcagc ctggatgggc ataagtagat aaataactgt gagctcgcac aaccgaaagg	3134
gcctgacccc ttacttagct ccctccttga aacaaagagc agactgtgga gagctgggag	3194
agegeageea getegetett tgetgagage eeettttgae agaaageeea geacgaeeet	3254
gctggaagaa ctgacagtgc cctcgccctc ggccccgggg cctgtggggt tggatgccgc	3314
ggttctatac atatacat atatccacat ctatatagag agatagatat ctatttttcc	3374
cctgtggatt agccccgtga tggctccctg ttggctacgc agggatgggc agttgcacga	3434
aggcatgaat gtattgtaaa taagtaactt tgacttctga caaaaaacaa aaagtgctgc	3494
atggetegea tggaateeae gegeteeagg gaetetgeee geeeeegga etggagaegg	3554
catctcgttc acagcaccca ccctcttacc tgataagttc catcgtatca aactttctat	3614
aaacaaaata cagtataatc agaaagtgcc atttcgccat tatttgtgat cggtaggcag	3674
ttcagagcat aagttaactg tgaaaaaaat gtaaaggttt tatttaggac atttgcatgg	3734
ctagtcatca gtccatttta tgagttaaca atgtattttg ttgagggaag tttttagggg	3794
ttgttttggg ttcttttatt ttgatggtga tgttttattt tattttattt ttttcagggg	3854
gtcttttttt taatacatat ccaataatgc cttccatctg aatgtaaaat aagtacccat	3914
gatttctatt atagtatcag tgtaattatt taaaaaatga ttttgaggca gttaagcatg	3974
accaattaat gtcactctag tgcttaggct gcgatcctat ggtagcaatt ctgtgctggt	4034
ataaatetta ettataaagt aggaaaagag aaccgaggaa gcacgtgaaa ettactaatt	4094
ctattcgagg attttataat ggcatatttt ttcagtatta aagcgaaaat gttttcaact	4154
ctgggtcctt accttttcc agcttcatat ttgcaagtgg taaattggat ttgcggtgga	4214
agagacaggg gagggaaacg gttggggtta gatcccttcc tgagctacat taaggctctt	4274
tctctaatcg ccttacttag ctttttaccc tttaagtagc tcctcttccc tcgcccccac	4334
cctctacccc acccccacct tcgctcagac tttaccggct ttccccagtc cataaaggtc	4394
ttgccccaac actcacccct tcttttttc ccctctccaa atgcagcagt gaatcccttt	4454
attaatactg gaaatccctc tctgctgctt ttgttggtgc tgcccacact gcagatatat	4514
taaggatgtt aggagagatt tgatttaatt gactctgcct agataggtct cattaaacag	4574
agtggagatt tcattggtca gcactcctca atgaaagaca gacctaatga ctggcatttg	4634
agatgctgct ggcattttga attcaacatc tgctgaaaac ggtaaaacta attagtgccc	4694
acceaecete ecegececag caactgeata ttgaaatttg ttaaageaet catettatg	4754
gaaattaatc attatcctaa agaagtgttt ctctcccatc atccggattt ctggttgtgg	4814

cccagcaatt	aacaaaaaca	gcttcaactg	ttcgaatttt	atgaaccaat	gtaactctgg	4874
cctcaatcat	attcctctgg	gatttctaaa	cagcagttaa	gctacaaaaa	gcaaacaaaa	4934
ccacacatat	tgatggagtc	tgcattccac	cacatatcca	cccttgagaa	gtatgtcaaa	4994
agactgcaga	ctatagattt	ttttttaata	taggattata	aatcagctag	tgaaagacct	5054
cagagcagtt	gtaagtagat	ctgccatcta	gaactcatat	tctaaaggga	agtgatttct	5114
cagaacagtg	atgttctgga	atatgtatta	tttattttaa	cacttttta	ataaaatctt	5174
tattataaac	catg					5188

<210> 231 <211> 696 PRT <213> Homo sapiens

<400> 231

Met Leu Leu Trp Ile Leu Leu Glu Thr Ser Leu Cys Phe Ala Ala 1 5 10 15

Gly Asn Val Thr Gly Asp Val Cys Lys Glu Lys Ile Cys Ser Cys Asn 20 25 30

Glu Ile Glu Gly Asp Leu His Val Asp Cys Glu Lys Lys Gly Phe Thr $35 \hspace{1cm} 40 \hspace{1cm} 45$

Ser Leu Gln Arg Phe Thr Ala Pro Thr Ser Gln Phe Tyr His Leu Phe 50 60

Leu His Gly Asn Ser Leu Thr Arg Leu Phe Pro Asn Glu Phe Ala Asn 65 70 75 80

Phe Tyr Asn Ala Val Ser Leu His Met Glu Asn Asn Gly Leu His Glu 85 90 95

Ile Val Pro Gly Ala Phe Leu Gly Leu Gln Leu Val Lys Arg Leu His 100 105 110

Ile Asn Asn Asn Lys Ile Lys Ser Phe Arg Lys Gln Thr Phe Leu Gly 115 120 125

Leu Asp Asp Leu Glu Tyr Leu Gln Ala Asp Phe Asn Leu Leu Arg Asp 130 135 140

Ile Asp Pro Gly Ala Phe Gln Asp Leu Asn Lys Leu Glu Val Leu Ile 145 150 160

Leu Asn Asp Asn Leu Ile Ser Thr Leu Pro Ala Asn Val Phe Gln Tyr 165 170 175

Val Pro Ile Thr His Leu Asp Leu Arg Gly Asn Arg Leu Lys Thr Leu 180 185 190

Pro Tyr Glu Glu Val Leu Glu Gln Ile Pro Gly Ile Ala Glu Ile Leu 195 200 205

Leu Glu Asp Asn Pro Trp Asp Cys Thr Cys Asp Leu Leu Ser Leu Lys 210 215 220

Glu 225	Trp	Leu	Glu	Asn	11e 230	Pro	Lys	Asn	Ala	16U Leu 235	200 Ile	PCT Gly	FINA Arg	AL.ST Val	25 Val 240
Cys	Glu	Ala	Pro	Thr 245	Arg	Leu	Gln	Gly	Lys 250	Asp	Leu	Asn	Glu	Thr 255	Thr
Glu	Gln	Asp	Leu 260	Cys	Pro	Leu	Lys	Asn 265	Arg	Val	Asp	Ser	Ser 270	Leu	Pro
Ala	Pro	Pro 275	Ala	Gln	Glu	Glu	Thr 280	Phe	Ala	Pro	Gly	Pro 285	Leu	Pro	Thr
Pro	Phe 290	Lys	Thr	Asn	G1 y	Gln 295	Glu	Asp	His	Ala	Thr 300	Pro	Gly	Ser	Ala
Pro 305	Asn	Gly	Gly	Thr	Lys 310	lle	Pro	G1 y	Asn	Trp 315	Gln	Ile	Lys	Ile	Arg 320
Pro	Thr	Ala	Ala	11e 325	Ala	Thr	Gl y	Ser	Ser 330	Arg	Asn	Lys	Pro	Leu 335	Ala
Asn	Ser	Leu	Pro 340	Cys	Pro	Gly	Gly	Cys 345	Ser	Cys	Asp	His	11e 350	Pro	Gly
Ser	Gly	Leu 355	Lys	Met	Asn	Cys	Asn 360	Asn	Arg	Asn	Val	Ser 365	Ser	Leu	Ala
Asp	Leu 370	Lys	Pro	Lys	Leu	Ser 375	Asn	Val	Gln	Glu	Leu 380	Phe	Leu	Arg	Asp
Asn 385	Lys	Ile	His	Ser	Ile 390	Arg	Lys	Ser	His	Phe 395	Val	Asp	Tyr	Lys	Asn 400
Leu	Ile	Leu	Leu	Asp 405	Leu	Gly	Asn	Asn	Asn 410	Ile	Ala	Thr	Val	Glu 415	Asn
Asn	Thr	Phe	Lys 420	Asn	Leu	Leu	Asp	Leu 425	Arg	Trp	Leu	Туr	Met 430	Asp	Ser
Asn	туг	Leu 435	Asp	Thr	Leu	Ser	Arg 440	Glu	Lys	Phe	Ala	Gly 445	Leu	Gln	Asn
Leu	Glu 450	Tyr	Leu	Asn	Val	Glu 455	Tyr	Asn	Ala	Ile	Gln 460	Leu	lle	Leu	Pro
Gly 465	Thr	Phe	Asn	Ala	Met 470	Pro	Lys	Leu	Arg	11e 475	Leu	Ile	Leu	Asn	Asn 480
Asn	Leu	Leu	Arg	Ser 485	Leu	Pro	Val	Asp	Val 490	Phe	Ala	Gly	Val	Ser 495	Leu
			500					Tyr 505			-		510		•
Gly	Val	Leu 515	Asp	Gln	Leu	Thr	Ser 520	Ile	Ile	Gln	Ile	Asp 525	Leu	His	Gly
Asn	Pro 530	Trp	Glu	Cys	Ser	Cys 535	Thr	Ile	Val	Pro	Phe 540	Lys	Gln	Trp	Ala

Page 146

										160	200	PCT	FINA	L.SI	25	
Glu 545	Arg	Leu	Gly	Ser	Glu 550	Val	Leu	Met	Ser	Asp 555	Leu	Lys	Cys	Glu	Thr 560	
Pro	Val	Asn	Phe	Phe 565	Arg	Lys	Asp	Phe	Met 570	Leu	Leu	Ser	Asn	Asp 575	Glu	
Ile	Cys	Pro	Gln 580	Leu	Tyr	Ala	Arg	Ile 585	Ser	Pro	Thr	Leu	Thr 590	Ser	His	
Ser	Lys	Asn 595	Ser	Thr	Gly	Leu	Ala 600	Glu	Thr	Gly	Thr	His 605	Ser	Asn	Ser	
Tyr	Leu 610	Asp	Thr	Ser	Arg	Val 615	Ser	Ile	Ser	Val	Leu 620	Val	Pro	Gly	Leu	
Leu 625	Leu	Val	Phe	Val	Thr 630	Ser	Ala	Phe	Thr	Val 635	Val	Gly	Met	Leu	Val 640	
Phe	lle	Leu	Arg	Asn 645	Arg	Lys	Arg	Ser	Lys 650	Arg	Arg	Asp	Ala	Asn 655	Ser	
Ser	Ala	Ser	Glu 660	Ile	Asn	Ser	Leu	Gln 665	Thr	Val	Cys	Asp	Ser 670	Ser	Tyr	
Trp	His	Asn 675	Gly	Pro	Туr	Asn	Ala 680	Asp	Gly	Ala	His	Arg 685	Val	туг	Asp	
Cys	Gly 690	Ser	His	Ser	Leu	Ser 695	Asp									
<210 <211 <212 <213	.> 5 !> [232 506 DNA Homo	sap:	iens												
<220 <221 <222 <223	.> (!>	CDS (32).	. (3	46)												
<400 ttca		232 agc t	ctg	gatg	gt to	gaact	tgtag	ı			aag a Lys N	1et l				52
					atc Ile											100
					gtt Val											148
					aag Lys 45											196
					tgt Cys										Thr	244
					cca Pro									aga	atg	292
					tcc Ser											340

ggc ctc tgacttggga aagctgggca caaaaatctt catgagcaat atttctttct Gly Leu 105	396											
taatagaatg ttttattatt caagtcaagt tctagagtgt ttacatacta ttatataatg												
tacagtgtta ttttctgtac ttctgaataa atgtgcaata ttggaaataa	506											
<210> 233 <211> 105 <212> PRT <213> Homo sapiens												
<400> 233												
Met Ala Lys Met Phe Asp Leu Arg Thr Lys Ile Met Ile Gly Ile Gly 1 5 10 15												
Ser Ser Leu Val Ala Ala Met Val Leu Leu Ser Val Val Phe Cys 20 25 30												
Leu Tyr Phe Lys Val Ala Lys Ala Leu Lys Ala Ala Lys Asp Pro Asp 35 40 45												
Ala Val Ala Val Lys Asn His Asn Pro Asp Lys Val Cys Trp Ala Thr 50 55 60												
Asn Ser Gln Ala Lys Ala Thr Thr Met Glu Ser Cys Pro Ser Leu Gln 65 70 75 80												
Cys Cys Glu Gly Cys Arg Met His Ala Ser Ser Asp Ser Leu Pro Pro 90 95												
Cys Cys Cys Asp Ile Asn Glu Gly Leu 100 105												
<210> 234 <211> 1037 <212> DNA <213> Homo sapiens												
<220> <221> CDS <222> (180)(560) <223>												
<400> 234 gagcgaaggg aacatttaac cttgactttc cacagtcctg aggttcccaa aataaagggg	60											
aaccggaaat accaaaggat tatctccaat attccagggc cttctttctc atctctgtct	120											
ttaccatact tactggcctt ggctggctct tcagctcttg gatccttaat cgaggaagc	179											
atg acc acc aac ttg gat ctg aag gta tcc atg ctc agc ttc atc tca Met Thr Thr Asn Leu Asp Leu Lys Val Ser Met Leu Ser Phe Ile Ser 1 5 10 15	227											
gct acc tgc ttg ctc ctc tgc ctc aac ctg ttt gtg gca cag gtt cac Ala Thr Cys Leu Leu Cys Leu Asn Leu Phe Val Ala Gln Val His 20 25 30	275											
tgg cat act agg gat gcc atg gag tca gat ctc cta tgg acc tat tat Trp His Thr Arg Asp Ala Met Glu Ser Asp Leu Leu Trp Thr Tyr Tyr 35 40 45	`323											
ctt aac tgg tgc agt gac atc ttt tac atg ttt gct ggg atc atc tct Leu Asn Trp Cys Ser Asp Ile Phe Tyr Met Phe Ala Gly Ile Ile Ser 50 55 60	371											
ctt ctc aac tac tta act tcc aga tcg cct gcc tgt gat gaa aac gtc Page 148	419											

BNSDOCID: <WO ___ _ 03089583A2 1 >

16U 200 PCT FINAL.ST25 Leu Leu Asn Tyr Leu Thr Ser Arg Ser Pro Ala Cys Asp Glu Asn Val 65 70 75 80	
act gtg att cca aca gag aga tca agg ctg ggg gtt ggt ccg gtg act Thr Val Ile Pro Thr Glu Arg Ser Arg Leu Gly Val Gly Pro Val Thr 85 90 95	467
aca gta tca cct gct aaa gat gaa ggg cca agg tct gag atg gaa tct Thr Val Ser Pro Ala Lys Asp Glu Gly Pro Arg Ser Glu Met Glu Ser 100 105 110	515
Cta agt gtg aga gag aaa aat tta cca aag tca gga ctg tgg tgg Leu Ser Val Arg Glu Lys Asn Leu Pro Lys Ser Gly Leu Trp Trp 115 120 125	560
tgataggaaa acctaactat agcttgtctt aaaagcaggg gagaagctga gttgggaatg	620
gtcacataaa ttctgggaaa ctctcctaat atcatgtcca tattacttga ggagacagca	680
ttaaagctga tgaaatgtct tttgcgtgca ttggatccaa aatatatatg atagtcataa	740
agtaaataac tcacttaaga aaaacattto taaaagaaaa caacaatgtt tagagtcatg	800
aatgaaagaa actagtgaaa gatgcagtgt gtagaccaga gacctctttg ggtatcaggg	860
atctcatgga ccagaatggc ccgtggagaa gaatgttaat tacttctgtt tggaattttc	920
tttattatgt gtggctttgg gtatactcag gatggaaagc acttggacaa atactgttga	980
atctgaactt aatagcatta ccagaaatgg aataaatatc aatggatata agaccta	037
<210> 235 <211> 127 <212> PRT <213> Homo sapiens <400> 235	
Met Thr Thr Asn Leu Asp Leu Lys Val Ser Met Leu Ser Phe Ile Ser 1 5 10 15	
Ala Thr Cys Leu Leu Cys Leu Asn Leu Phe Val Ala Gln Val His 20 25 30	
Trp His Thr Arg Asp Ala Met Glu Ser Asp Leu Leu Trp Thr Tyr Tyr 35 40 45	
Leu Asn Trp Cys Ser Asp Ile Phe Tyr Met Phe Ala Gly Ile Ile Ser 50 55 60	
Leu Leu Asn Tyr Leu Thr Ser Arg Ser Pro Ala Cys Asp Glu Asn Val 65 70 75 80	
Thr Val 11e Pro Thr Glu Arg Ser Arg Leu Gly Val Gly Pro Val Thr 85 90 95	
Thr Val Ser Pro Ala Lys Asp Glu Gly Pro Arg Ser Glu Met Glu Ser 100 105 110	
Leu Ser Val Arg Glu Lys Asn Leu Pro Lys Ser Gly Leu Trp Trp 115 120 125	
<210> 236 <211> 1054 <212> DNA <213> Homo sapiens	
<220> <221> CDS <222> (152)(895)	

16U 200 PCT FINAL.ST25	
<400> 236 gttggcattc ggtggtcctg gcagttagct gagcacgccc tctgagccgc tcggtggaca	60
ccaggcactc tagtaggcct ggcctaccca gaaacagcag gagagagaag aaacaggcca	120
gctgtgagaa gccaaggaca ccgagtcagt c atg gca cct aag gcg gca aag Met Ala Pro Lys Ala Ala Lys 1 5	172
ggg gcc aag cca gag cca gca cca gct cca cct cca ccc ggg gcc aaa Gly Ala Lys Pro Glu Pro Ala Pro Ala Pro Pro Pro Gly Ala Lys 10 15 20	220
ccc gag gaa gac aag gac ggt aag gag cca tcg gac aaa cct caa Pro Glu Glu Asp Lys Lys Asp Gly Lys Glu Pro Ser Asp Lys Pro Gln 25 30 35	268
and gcg gtg cag gac cat and gag cca tcg gac and cct can and gcg Lys Ala Val Gln Asp His Lys Glu Pro Ser Asp Lys Pro Gln Lys Ala 40 45 50 55	316
gtg cag ccc aag cac gaa gtg ggc acg agg agg ggg tgt cgc cgc tac Val Gln Pro Lys His Glu Val Gly Thr Arg Arg Gly Cys Arg Arg Tyr 60 65 . 70	364
cgg tgg gaa tta aaa gac agc aat aaa gag ttc tgg ctc ttg ggg cac Arg Trp Glu Leu Lys Asp Ser Asn Lys Glu Phe Trp Leu Leu Gly His 75 80 85	412
gct gag atc aag att cgg agt ttg ggc tgc cta ata gct gca atg ata Ala Glu Ile Lys Ile Arg Ser Leu Gly Cys Leu Ile Ala Ala Met Ile 90 95 100	460
ctg ttg tcc tca ctc acc gtg cac ccc atc ttg agg ctt atc acc Leu Leu Ser Ser Leu Thr Val His Pro Ile Leu Arg Leu Ile Ile Thr 105 110 115	508
atg gag ata tcc ttc ttc agc ttc ttc atc tta ctg tac agc ttt gcc Met Glu Ile Ser Phe Phe Ser Phe Phe Ile Leu Leu Tyr Ser Phe Ala 120 125 130 135	556
att cat aga tac ata ccc ttc atc ctg tgg ccc att tct gac ctc ttc Ile His Arg Tyr Ile Pro Phe Ile Leu Trp Pro Ile Ser Asp Leu Phe 140 145 150	604
aac gac ctg att gct tgt gcg ttc ctt gtg gga gcc gtg gtc ttt gct Asn Asp Leu Ile Ala Cys Ala Phe Leu Val Gly Ala Val Val Phe Ala 155 160 165	652
gtg aga agt cgg cga tcc atg aat ctc cac tac tta ctt gct gtg atc Val Arg Ser Arg Arg Ser Met Asn Leu His Tyr Leu Leu Ala Val Ile 170 175 180	700
Ctt att ggt gcg gct gga gtt ttt gct ttt atc gat gtg tgt ctt caa Leu Ile Gly Ala Ala Gly Val Phe Ala Phe Ile Asp Val Cys Leu Gln 185 190 195	748
aga aac cac ttc aga ggc aag aag gcc aaa aag cat atg ctg gtt cct Arg Asn His Phe Arg Gly Lys Lys Ala Lys Lys His Met Leu Val Pro 200 205 210 215	796
cct cca gga aag gaa aaa gga ccc cag cag ggc aag gga cca gaa ccc Pro Pro Gly Lys Glu Lys Gly Pro Gln Gln Gly Lys Gly Pro Glu Pro 220 225 230	844
gcc aag cca cca gaa cct ggc aag cca cca ggg cca gca aag gga aag Ala Lys Pro Pro Glu Pro Gly Lys Pro Pro Gly Pro Ala Lys Gly Lys 235 240 245	892
	945
atatgtttcc actctcttcc ttgtcttctt tctggaatgg ttttcttttc	005
accacctttg cttggaaaag aatggattaa tggattctaa aagcctaaa 10	054

```
<210> 237

<211> 248

<212> PRT

<213> Homo sapiens

<400> 237
```

Met Ala Pro Lys Ala Ala Lys Gly Ala Lys Pro Glu Pro Ala Pro Ala l $1 \\ 5 \\ 10 \\ 15$

Pro Pro Pro Gly Ala Lys Pro Glu Glu Asp Lys Lys Asp Gly Lys 20 25 30

Glu Pro Ser Asp Lys Pro Gln Lys Ala Val Gln Asp His Lys Glu Pro 35 40 45

Ser Asp Lys Pro Gln Lys Ala Val Gln Pro Lys His Glu Val Gly Thr 50 60

Arg Arg Gly Cys Arg Arg Tyr Arg Trp Glu Leu Lys Asp Ser Asn Lys 65 70 75 80

Glu Phe Trp Leu Leu Gly His Ala Glu Ile Lys Ile Arg Ser Leu Gly 85 90 95

Cys Leu Ile Ala Ala Met Ile Leu Leu Ser Ser Leu Thr Val His Pro $100 \hspace{1.5cm} 100 \hspace{1.5cm} 105 \hspace{1.5cm} 110 \hspace{1.5cm}$

Ile Leu Arg Leu Ile Ile Thr Met Glu Ile Ser Phe Phe Ser Phe Phe 115 120 125

Ile Leu Leu Tyr Ser Phe Ala Ile His Arg Tyr Ile Pro Phe Ile Leu 130 135 140

Trp Pro Ile Ser Asp Leu Phe Asn Asp Leu Ile Ala Cys Ala Phe Leu 145 150 155 160

Val Gly Ala Val Val Phe Ala Val Arg Ser Arg Arg Ser Met Asn Leu 165 170 175

His Tyr Leu Leu Ala Val Ile Leu Ile Gly Ala Ala Gly Val Phe Ala 180 185 190

Phe Ile Asp Val Cys Leu Gln Arg Asn His Phe Arg Gly Lys Lys Ala 195 200 205

Lys Lys His Met Leu Val Pro Pro Pro Gly Lys Glu Lys Gly Pro Gln 210 $$ 215 $$ 220

Gln Gly Lys Gly Pro Glu Pro Ala Lys Pro Pro Glu Pro Gly Lys Pro 225 230 235 240

Pro Gly Pro Ala Lys Gly Lys Lys 245

<210> 238 <211> 487

<212> DNA <213> Homo sapiens

<220> <221> CDS

Page 152

16U 200 PCT FINAL.ST25

<pre></pre>	
<pre><400> 238 agtggcagct tggctg atg agc tat aag cca gcc ttg ttt ggg ttc cta ttc</pre>	
Ctt ctg ctg ttg ctt agc aac tgg ttg gtc aag tat gaa cac aag ctc Leu Leu Leu Leu Ser Asn Trp Leu Val Lys Tyr Glu His Lys Leu 15 20 25	100
acc ctc cca gag ccc cag cag gag gaa gag aaa cca aag act tct gaa Thr Leu Pro Glu Pro Gln Gln Glu Glu Glu Lys Pro Lys Thr Ser Glu 30 35 40	148
aac gac too aag aac ago aag goo gtg aac aca aaa gaa gto aat aga Asn Asp Ser Lys Asn Ser Lys Ala Val Asn Thr Lys Glu Val Asn Arg 45 50 55 60	196
acg cat gcc tgc ttt gcc ctc cag gac gag atc ctc caa cgg ctg ttg Thr His Ala Cys Phe Ala Leu Gln Asp Glu Ile Leu Gln Arg Leu Leu 65 70 75	244
ttc agt gaa atg aag atg aag gtc cta gaa aat cag atg ttc atc ata Phe Ser Glu Met Lys Met Lys Val Leu Glu Asn Gln Met Phe Ile Ile 80 85 90	292
tgg aat aaa atg aat cac cac ggg cgg tca agc aga cat cgg aat ttt Trp Asn Lys Met Asn His His Gly Arg Ser Ser Arg His Arg Asn Phe 95 100 105	340
ccc atg aaa aaa cac aga atg agg agg cat gag tca att tgc ccc acc Pro Met Lys Lys His Arg Met Arg Arg His Glu Ser Ile Cys Pro Thr 110 115 120	388
ctg tct gac tgt act tcg agt tcc ccc agc taatgaggcc gaggcgggct Leu Ser Asp Cys Thr Ser Ser Ser Pro Ser 125 130	438
ggcctctgcc gatgttacct tttacctcag taaaacccag tcacagcct	487
<210> 239 <211> 134 <212> PRT <213> Homo sapiens	
<211> 134 <212> PRT	
<211> 134 <212> PRT <213> Homo sapiens	
<211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu	
<211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu 1	
<pre><211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu 1</pre>	
<pre> <211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu Leu 1</pre>	
<pre><211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu Leu 1</pre>	
<pre> <211> 134 <212> PRT <213> Homo sapiens <400> 239 Met Ser Tyr Lys Pro Ala Leu Phe Gly Phe Leu Phe Leu Leu Leu Leu 1</pre>	

Thr Ser Ser Ser Pro Ser

16U 200 PCT FINAL.ST25 115 120 125

Inr	130	Ser	ser	Pro	Ser											
<21 <21 <21 <21	1> 2>	240 846 DNA Homo	sap	iens												
<22 <22 <22 <22	1> 2>	CDS (108) (725)												
<40 att		240 aga	ctct	gatt	gg c	tggg	caga	t gg	gctg	actg	gct	gggc	aga	tggg	tgggtg	60
agt	tccc	tct	cccc	agag	cc a	tcgg	ccag	g ta	ccaa	agct	cag	ctgt		gat Asp		116
caa Gln	cag Gln 5	gag Glu	gac Asp	ctg Leu	cgc Arg	ttc Phe 10	cct Pro	G] A aaa	atg Met	tgg Trp	gtc Val 15	tca Ser	ttg Leu	tac Tyr	ttt Phe	164
														atc Ile		212
														cag Gln 50		260
														ttg Leu		308
														aac Asn		356
														aat Asn		404
gat Asp 100	gtt Val	ctg Leu	tgg Trp	gat Asp	ttg Leu 105	gac Asp	atc Ile	ccc Pro	gaa Glu	ggc Gly 110	agg Arg	agc Ser	cat His	gct Ala	gac Asp 115	452
														cct Pro 130		500
														cca Pro		548
														cta Leu		596
ccc Pro	ctg Leu 165	ctg Leu	aac Asn	cac His	tct Ser	gtc Val 170	tcc Ser	tat Tyr	cct Pro	ttg Leu	gcc Ala 175	acc Thr	tgt Cys	cct Pro	gaa Glu	644
														gac Asp		692
agc Ser	ttc Phe	aat Asn	gcc Ala	aag Lys 200	cct Pro	ttt Phe	cct Pro	tca Ser	gaa Glu 205	ctg Leu	tago	cctcc	etc t	cac,t	gaagg	745
tggg	agct	gc a	ggaa	itcaç	ıg tç	caga	igtag	gaa	atgg	aac	taad	ctca	igg a	aggt	ggtat	805
tgac	agaç	igt c	agga	ccca	c ct	ggat	gtca	tgo	tatg	aaa	С					846

<211> 206 <212> PRT <213> Homo sapiens <400> 241 Met Asp Ser Gln Glu Asp Leu Arg Phe Pro Gly Met Trp Val Ser Leu Tyr Phe Gly Ile Leu Gly Leu Cys Ser Val Ile Thr Gly Gly Cys 20 25 30Ile Ile Phe Leu His Trp Arg Lys Asn Leu Arg Arg Glu Glu His Ala 35 40 45 Gln Gln Trp Val Glu Val Met Arg Ala Ala Thr Phe Thr Tyr Ser Pro 50 55 60 Leu Leu Tyr Trp Ile Asn Lys Arg Arg Arg Tyr Gly Met Asn Ala Ala 65 70 75 80 lle Asn Thr Gly Pro Ala Pro Ala Val Thr Lys Thr Glu Thr Glu Val $85 \\ 90 \\ 95$ Gln Asn Pro Asp Val Leu Trp Asp Leu Asp Ile Pro Glu Gly Arg Ser 100 105 110 His Ala Asp Gln Asp Ser Asn Pro Lys Ala Glu Ala Pro Ala Pro Leu 115 120 125 Gln Pro Ala Leu Gln Leu Ala Pro Gln Gln Pro Gln Ala Arg Ser Pro Phe Pro Leu Pro Ile Phe Gln Glu Val Pro Phe Ala Pro Pro Leu Cys Asn Leu Pro Pro Leu Leu Asn His Ser Val Ser Tyr Pro Leu Ala Thr Cys Pro Glu Arg Asn Val Leu Phe His Ser Leu Leu Asn Leu Ala Gln Glu Asp His Ser Phe Asn Ala Lys Pro Phe Pro Ser Glu Leu <210> 242 <211> 663 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (40)..(585) <400> 242 tagtteetag agetgetget tattaaaatg teaacatet tea tet tet age tgg Ser Ser Ser Trp gac aac ctc tta gag tct ctc tct ctc agc aca gta tgg aat tgg ata Asp Asn Leu Leu Glu Ser Leu Ser Leu Ser Thr Val Trp Asn Trp Ile 10 15 20 102

Page 154

<210>

Gln Ala Ser Phe Leu Gly Glu Thr Ser Ala Pro Gln Gln Thr Ser Lo 25 30 35	tg 150 eu
gga cta tta gat aat ctt gct cca gct gtg caa atc atc ttg agg at Gly Leu Leu Asp Asn Leu Ala Pro Ala Val Gln Ile Ile Leu Arg II 40 45 50	tt 198 le
tct ttc ttg att tta ttg gga ata gga ata tat gcc tta tgg aaa cg Ser Phe Leu Ile Leu Leu Gly Ile Gly Ile Tyr Ala Leu Trp Lys Ar 55 60 65	ga 246 cg
agt att cag tca att cag aaa aca ttg ttg ttt gta atc aca ctc ta Ser Ile Gln Ser Ile Gln Lys Thr Leu Leu Phe Val Ile Thr Leu Ty 70 75 80 85	/r
aaa ctt tac aag aag ggc tca cat att ttt gag gct ttg cta gcc aa Lys Leu Tyr Lys Lys Gly Ser His Ile Phe Glu Ala Leu Leu Ala As 90 95 100	ac 342 sn
cca gaa gga agt ggt ctc cga att caa gac aat aat aat ctt ttc ct Pro Glu Gly Ser Gly Leu Arg Ile Gln Asp Asn Asn Asn Leu Phe Le 105 110 115	.g 390 eu
tcc ttg ggt ctg caa gag aaa att ttg aaa aaa ctt aag aca gtg ga Ser Leu Gly Leu Gln Glu Lys Ile Leu Lys Lys Leu Lys Thr Val Gl 120 125 130	ia 438 u
aac aaa atg aag aac cta gaa ggg ata atc gtt gct caa aaa cct gc Asn Lys Met Lys Asn Leu Glu Gly Ile Ile Val Ala Gln Lys Pro Al 135 140 145	c 486 a
acg aag agg gat tgc tcc tct gag ccc tac tgc agc tgc tct gac tg Thr Lys Arg Asp Cys Ser Ser Glu Pro Tyr Cys Ser Cys Ser Asp Cy 150 160 16	s
Cag agt ccc ttg tcc aca tca ggg ttt act tcc ccc att tga aat gt Gln Ser Pro Leu Ser Thr Ser Gly Phe Thr Ser Pro Ile Asn Va 170 175 18	1
atg gactccaatc ttttccagga aagcactgtt tccctcatgt gtgcagtggt Met	635
gtatcaataa agatagagaa cgctattg	663
<210> 243 <211> 178 <212> PRT <213> Homo sapiens	
<210> 243 <211> 178 <212> PRT <213> Homo sapiens <400> 243	663
<210> 243 <211> 178 <212> PRT <213> Homo sapiens	663
<210> 243 <211> 178 <212> PRT <213> Homo sapiens <400> 243 Ser Ser Ser Ser Trp Asp Asn Leu Leu Glu Ser Leu Ser Th	663 r
<pre> <210> 243 <211> 178 <212> PRT <213> Homo sapiens <400> 243 Ser Ser Ser Ser Trp Asp Asn Leu Leu Glu Ser Leu Ser Leu Ser Th</pre>	663 r
<pre> <210> 243 <211> 178 <212> PRT <213> Homo sapiens <400> 243 Ser Ser Ser Ser Trp Asp Asn Leu Leu Glu Ser Leu Ser Leu Ser Th 1</pre>	663 r
<pre> <210> 243 <211> 178 <212> PRT <213> Homo sapiens <400> 243 Ser Ser Ser Ser Trp Asp Asn Leu Leu Glu Ser Leu Ser Leu Ser Th 1</pre>	663

16U 200 PCT FINAL.ST25 Ala Leu Leu Ala Asn Pro Glu Gly Ser Gly Leu Arg Ile Gln Asp Asn Asn Asn Leu Phe Leu Ser Leu Gly Leu Gln Glu Lys Ile Leu Lys Lys Leu Lys Thr Val Glu Asn Lys Met Lys Asn Leu Glu Gly Ile Ile Val Ala Gln Lys Pro Ala Thr Lys Arg Asp Cys Ser Ser Glu Pro Tyr Cys Ser Cys Ser Asp Cys Gln Ser Pro Leu Ser Thr Ser Gly Phe Thr Ser Pro Ile <210> 244 541 <211> <212> DNA Homo sapiens <220> <221> CDS <222> (62)..(349) <223> <400> 244 catatotgtt gocactotga gtttggcatt atgtgttcga attttaacca catatotatt 60 g atg tgc ttt gct ggt ttt agt ttt aag gag aaa ata ttt att gct tta Met Cys Phe Ala Gly Phe Ser Phe Lys Glu Lys Ile Phe Ile Ala Leu 1 15 109 gca tgg atg ccc aaa gct aca gta cag gct gtg tta ggt cct ctg gct Ala Trp Met Pro Lys Ala Thr Val Gln Ala Val Leu Gly Pro Leu Ala 157 cta gaa aca gca aga gtc tct gca ccc cac ttg gaa cca tat gcg aag Leu Glu Thr Ala Arg Val Ser Ala Pro His Leu Glu Pro Tyr Ala Lys 205 40 gat gtg atg tca gta gca ttt tta gcc atc tcg atc aca gct cca aat Asp Val Met Ser Val Ala Phe Leu Ala Ile Ser Ile Thr Ala Pro Asn 253 gga gct cta ctt atg ggc att ctg ggg cct aaa atg ctt aca cgc cat Gly Ala Leu Leu Met Gly Ile Leu Gly Pro Lys Met Leu Thr Arg His 65 70 70 75 80301 tat gat cca agc aaa ata aaa ctg caa ttg tca aca tta gaa cat cat Tyr Asp Pro Ser Lys Ile Lys Leu Gln Leu Ser Thr Leu Glu His His 85 90 95 349 taaaaagttt acctgtcatc atctgcctgc ttcttttaat gaattatttc acatgacaga 409 agaattttaa agtagaaata tgtagggact gtacagaaaa tccaggattt agtaaacatg 469 tgatttcagt acagggcttt tcttggactt tttactccaa agttaattta ataaaaataa 529 tattaaatgg aa 541 <210> <211> 96 <212> PRT <213> Homo sapiens <400> Met Cys Phe Ala Gly Phe Ser Phe Lys Glu Lys Ile Phe Ile Ala Leu

Ala Trp Met Pro Lys Ala Thr Val Gln Ala Val Leu Gly Pro Leu Ala Leu Glu Thr Ala Arg Val Ser Ala Pro His Leu Glu Pro Tyr Ala Lys Asp Val Met Ser Val Ala Phe Leu Ala Ile Ser Ile Thr Ala Pro Asn Gly Ala Leu Leu Met Gly Ile Leu Gly Pro Lys Met Leu Thr Arg His 65 70 75 80 Tyr Asp Pro Ser Lys Ile Lys Leu Gln Leu Ser Thr Leu Glu His His 85 90 95 <210> 246 <211> 2499 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (128) . . (2284) <223> <400> 246 gcaagaggcc ccttgtggcc accgagtcct ccgacgccct cgccaggctg gcctttgggt 60 tggcccaggc aggacgggct gccgagagca ctcgggccgc gtcgccagga gccgcccagg gtgagcc atg ttc gta ggc gtc gcc cgg cac tct ggg agc cag gat gaa Met Phe Val Gly Val Ala Arg His Ser Gly Ser Gln Asp Glu 169 gtc tca agg gga gta gag ccg ctg gag gcc gcg cgg gcc cag cct gct Val Ser Arg Gly Val Glu Pro Leu Glu Ala Ala Arg Ala Gln Pro Ala aag gac agg agg gcc aag gga acc ccg aag tcc tcg aag ccc ggg aaa Lys Asp Arg Arg Ala Lys Gly Thr Pro Lys Ser Ser Lys Pro Gly Lys 35 40 45265 aaa cac cgg tat ctg aga cta ctt cca gag gcc ttg ata agg ttc ggc Lys His Arg Tyr Leu Arg Leu Leu Pro Glu Ala Leu Ile Arg Phe Gly 50 55 60ggt ttc cga aaa agg aaa aaa gcc aag tcc tca gtt tcc aag aag ccg Gly Phe Arg Lys Arg Lys Lys Ala Lys Ser Ser Val Ser Lys Lys Pro 65 70 75 361 gga gaa gtg gat gac agt ttg gag cag ccc tgt ggt ttg ggc tgc tta Gly Glu Val Asp Asp Ser Leu Glu Gln Pro Cys Gly Leu Gly Cys Leu 80 85 409 gtc agc acc tgc tgt gag tgt tgc aat aac att cgc tgc ttc atg att Val Ser Thr Cys Cys Glu Cys Cys Asn Asn Ile Arg Cys Phe Met Ile 457 ttc tac tgc atc ctg ctc ata tgt caa ggt gtg gtg ttt ggt ctt ata Phe Tyr Cys Ile Leu Leu Ile Cys Gln Gly Val Val Phe Gly Leu Ile 115 120 125 505 gat gtc agc att ggc gat ttt cag aag gaa tat caa ctg aaa acc att Asp Val Ser Ile Gly Asp Phe Gln Lys Glu Tyr Gln Leu Lys Thr Ile 130 135 140 553 gag aag ttg gca ttg gaa aag agt tac gat att tca tct ggc ctg gta Glu Lys Leu Ala Leu Glu Lys Ser Tyr Asp Ile Ser Ser Gly Leu Val 145 $$ 150 $$ 155 $$ 601 gca ata ttt ata gca ttc tat gga gac aga aaa aaa gta ata tgg ttt 649 Ala Ile Phe Ile Ala Phe Tyr Gly Asp Arg Lys Lys Val Ile Trp Phe

	160					165				160	200 170	PCT	FINA	L.ST	25	
gta Val 175	gct Ala	tcc Ser	tcc Ser	ttt Phe	tta Leu 180	ata Ile	gga Gly	ctt Leu	gga Gly	tca Ser 185	ctt Leu	tta Leu	tgt Cys	gct Ala	ttt Phe 190	697
cca Pro	tcc Ser	att Ile	aat Asn	gaa Glu 195	gaa Glu	aat Asn	aaa Lys	caa Gln	agt Ser 200	aag Lys	gta Val	gga Gly	att Ile	gaa Glu 205	gat Asp	745
att Ile	tgc Cys	gaa Glu	gaa Glu 210	ata Ile	aag Lys	gtt Val	gtc Val	agt Ser 215	ggt Gly	tgc Cys	cag Gln	agc Ser	agt Ser 220	ggt Gly	ata Ile	793
tça Ser	ttc Phe	caa Gln 225	tca Ser	aaa Lys	tac Tyr	ctg Leu	tct Ser 230	ttc Phe	ttc Phe	atc Ile	ctt Leu	ggg G1 y 235	cag Gln	act Thr	gtg Val	841
cag Gln	gga Gly 240	ata Ile	gca Ala	gga Gly	atg Met	cct Pro 245	ctt Leu	tat Tyr	atc Ile	ctt Leu	gga Gly 250	ata Ile	acc Thr	ttt Phe	att Ile	889
gat Asp 255	gag Glu	aat Asn	gtt Val	gct Ala	aca Thr 260	cac His	tca Ser	gct Ala	ggt Gly	atc Ile 265	tat Tyr	tta Leu	ggt Gly	att Ile	gca Ala 270	937
gaa Glu	tgt Cys	aca Thr	tca Ser	atg Met 275	att Ile	gga Gly	tat Tyr	gct Ala	ctg Leu 280	ggt Gly	tat Tyr	gtg Val	cta Leu	gga Gly 285	gca Ala	985
cca Pro	cta Leu	gtt Val	aaa Lys 290	gtc Val	cct Pro	gag Glu	aat Asn	act Thr 295	act Thr	tct Ser	gca Ala	aca Thr	aac Asn 300	act Thr	aca Thr	1033
gtc Val	aat Asn	aat Asn 305	ggt Gly	agt Ser	cca Pro	gaa Glu	tgg Trp 310	cta Leu	tgg Trp	act Thr	tgg Trp	tgg Trp 315	att Ile	aat Asn	ttt Phe	1081
ctt Leu	ttt Phe 320	gcc Ala	gct Ala	gtc Val	gtt Val	gca Ala 325	tgg Trp	tgt Cys	aca Thr	tta Leu	ata Ile 330	Pro	ttg Leu	tca Ser	tgc Cys	1129
ttt Phe 335	cca Pro	aac Asn	aat Asn	atg Met	cca Pro 340	ggt Gly	tca Ser	aca Thr	cgg Arg	ata Ile 345	aaa Lys	gct Ala	agg Arg	aaa Lys	cgt Arg 350	1177
aaa Lys	cag Gln	ctt Leu	cat His	ttt Phe 355	ttt Phe	gac Asp	agc Ser	aga Arg	ctt Leu 360	aaa Lys	gat Asp	ctg Leu	aaa Lys	ctt Leu 365	gga Gly	1225
act Thr	aat Asn	atc Ile	aag Lys 370	Asp	tta Leu	tgt Cys	gct Ala	gct Ala 375	ctt Leu	tgg Trp	att	ctg Leu	atg Met 380	Arg	aat Asn	1273
cca Pro	gtg Val	ctc Leu 385	lle	t gc Cys	cta Leu	gct Ala	ctg Leu 390	Ser	aaa Lys	gct Ala	aca Thr	gaa Glu 395	Tyr	tta Leu	gtt Val	1321
att Ile	att Ile 400	Gly	gct Ala	tct Ser	gaa Glu	ttt Phe 405	Leu	cct Pro	ata Ile	tat Tyr	Leu 410	Glu	aat Asn	cag Gln	ttt Phe	1369
ata Ile 415	Leu	aca Thr	ccc Pro	act	gtg Val 420	Ala	act Thr	aca Thr	ctt Leu	gca Ala 425	G1 7	ctt Leu	gtt Val	tta Leu	att Ile 430	1417
cca Pro	gga Gly	ggt Gly	gca Ala	Leu 435	Gly	cag Gln	ctt Leu	ctg Leu	gga Gly 440	Gly	gto Val	att Ile	gtt Val	Ser 445	Thr	1465
tta Leu	gaa Glu	atg Met	Ser 450	Cys	aaa Lys	gcc Ala	ctt Leu	atg Met 455	Arg	ttt Phe	ata Ile	ato Met	gtt Val 460	Thr	tct Ser	1513
gtg Val	ata Ile	tca Ser 465	Leu	ata Ile	ctg Leu	ctt Leu	gtg Val 470	Phe	att Ile	att Ile	ttt Phe	gta Val 475	Arg	tgt Cys	aat Asn	1561
cca	gtg	caa	ttt	gct	ggg	ato	aat	gaa	gat	tat	: gat		a aca : 158		aag	1609

Pro	Val 480	Gln	Phe	Ala	Gly	Ile 485	Asn	Glu	Asp	16U Tyr	200 Asp 490	PCT Gly	FIN) Thr	AL.ST Arg	r25 Lys	
ttg Leu 495	gga Gly	aac Asn	ctc Leu	acg Thr	gct Ala 500	cct Pro	tgc Cys	aat Asn	gaa Glu	aaa Lys 505	tgt Cys	aga Arg	tgc Cys	tca Ser	tct Ser 510	1657
tca Ser	att Ile	tat Tyr	tct Ser	tct Ser 515	ata Ile	tgt Cys	gga Gly	aga Arg	gat Asp 520	gat Asp	att Ile	gaa Glu	tat Tyr	ttt Phe 525	tct Ser	1705
gcc Ala	tgc Cys	ttt Phe	gca Ala 530	ggg Gly	tgt Cys	aca Thr	tat Tyr	tct Ser 535	aaa Lys	gca Ala	caa Gln	aac Asn	caa Gln 540	aaa Lys	aag Lys	1753
atg Met	tac Tyr	tac Tyr 545	aat Asn	tgt Cys	tct Ser	tgc Cys	att 1 1 e 550	aaa Lys	gaa Glu	gga Gly	tta Leu	ata Ile 555	act Thr	gca Ala	gat Asp	1801
gca Ala	gaa Glu 560	ggt Gly	gat Asp	ttt Phe	att Ile	gat Asp 565	gcc Ala	aga Arg	ccc Pro	Gly ggg	aaa Lys 570	tgt Cys	gat Asp	gca Ala	aag Lys	1849
tgc Cys 575	tat Tyr	aag Lys	tta Leu	cct Pro	ttg Leu 580	ttc Phe	att Ile	gct Ala	ttt Phe	atc Ile 585	ttt Phe	tct Ser	aca Thr	ctt Leu	ata Ile 590	1897
ttt Phe	tct Ser	ggt Gly	ttt Phe	tct Ser 595	ggt Gly	gta Val	cca Pro	atc Ile	gtc Val 600	ttg Leu	gcc Ala	atg Met	acg Thr	cgg Arg 605	gtt Val	1945
gta Val	cct Pro	gac Asp	aaa Lys 610	ctg Leu	cgt Arg	tct Ser	ctg Leu	gcc Ala 615	ttg Leu	ggt Gly	gta Val	agc Ser	tat Tyr 620	gtg Val	att Ile	1993
ttg Leu	aga Arg	ata Ile 625	ttt Phe	ggg Gly	act Thr	att Ile	cct Pro 630	gga Gly	cca Pro	tca Ser	atc Ile	ttt Phe 635	aaa Lys	atg Met	tca Ser	2041
gga Gly	gaa Glu 640	act Thr	tct Ser	tgt Cys	att Ile	tta Leu 645	cgg Arg	gat Asp	gtt Val	aat Asn	aaa Lys 650	tgt Cys	gga Gly	cac His	aca Thr	2089
gga G1 y 65 5	cgt Arg	tgt Cys	tgg Trp	ata Ile	tat Tyr 660	aac Asn	aag Lys	aca Thr	aaa Lys	atg Met 665	gct Ala	ttc Phe	tta Leu	ttg Leu	gta Val 670	2137
gga Gly	ata Ile	tgt Cys	ttt Phe	ctt Leu 675	tgc Cys	aaa Lys	cta Leu	tgc Cys	act Thr 680	atc Ile	atc Ile	ttc Phe	act Thr	act Thr 685	att Ile	2185
gca Ala	ttt Phe	ttc Phe	ata Ile 690	tac Tyr	aaa Lys	cgt Arg	cgt Arg	cta Leu 695	aat Asn	gag Glu	aac Asn	act Thr	gac Asp 700	ttc Phe	cca Pro	2233
gat Asp	gta Val	act Thr 705	gtg Val	aag Lys	aat Asn	cca Pro	aaa Lys 710	gtt Val	aag Lys	aaa Lys	Lys	gaa Glu 715	gaa Glu	act Thr	gac Asp	2281
ttg Leu	taac	tgga	tc a	tcat	tgtg	a tt	gcag	atca	ttt	gagg	atc	agag	tgtg	aa	٠	2334
aacg	agtt	tc t	cttt	taca	g at	tctc	caag	att	tgtt	tct	gtgc	ccaa	ct t	tcag	aagag	2394
													ga t	ctgc	attt	2454
cata	ataa	ag t	gtcc	tatt	g tg	aaac	aaaa	aaa	aaaa	aaa	aaaa,					2499
<210 <211		47 19												,		
<212 <213	> P	RT	sapi	ens										,;		
<400		47		_												
Met 1	Phe	Val (√al . 5	Ala i	Arg i	His		31y : 10	Ser	Gln i	Asp (Val 15	Ser	

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

Arg Gly Val Glu Pro Leu Glu Ala Ala Arg Ala Gln Pro Ala Lys Asp 20 25 30

Arg Arg Ala Lys Gly Thr Pro Lys Ser Ser Lys Pro Gly Lys Lys His $35 \hspace{1cm} 40 \hspace{1cm} 45$

Arg Tyr Leu Arg Leu Leu Pro Glu Ala Leu 11e Arg Phe Gly Gly Phe 50 55 60

Arg Lys Arg Lys Lys Ala Lys Ser Ser Val Ser Lys Lys Pro Gly Glu

Val Asp Asp Ser Leu Glu Gln Pro Cys Gly Leu Gly Cys Leu Val Ser 85 90 95

Thr Cys Cys Glu Cys Cys Asn Asn Ile Arg Cys Phe Met Ile Phe Tyr 100 105 110

Cys Ile Leu Leu Ile Cys Gln Gly Val Val Phe Gly Leu Ile Asp Val 115 120 125

Ser Ile Gly Asp Phe Gln Lys Glu Tyr Gln Leu Lys Thr Ile Glu Lys 130 135 140

Leu Ala Leu Glu Lys Ser Tyr Asp Ile Ser Ser Gly Leu Val Ala Ile 145 150 160

Phe Ile Ala Phe Tyr Gly Asp Arg Lys Lys Val Ile Trp Phe Val Ala 165 170 175

Ser Ser Phe Leu Ile Gly Leu Gly Ser Leu Leu Cys Ala Phe Pro Ser 180 185 190

Ile Asn Glu Glu Asn Lys Gln Ser Lys Val Gly Ile Glu Asp Ile Cys 195 200 205

Glu Glu Ile Lys Val Val Ser Gly Cys Gln Ser Ser Gly Ile Ser Phe 210 215 220

Gln Ser Lys Tyr Leu Ser Phe Phe Ile Leu Gly Gln Thr Val Gln Gly 225 230 240

Ile Ala Gly Met Pro Leu Tyr Ile Leu Gly Ile Thr Phe Ile Asp Glu 245 250 255

Asn Val Ala Thr His Ser Ala Gly Ile Tyr Leu Gly Ile Ala Glu Cys 265 270

Thr Ser Met Ile Gly Tyr Ala Leu Gly Tyr Val Leu Gly Ala Pro Leu 275 280 285

Val Lys Val Pro Glu Asn Thr Thr Ser Ala Thr Asn Thr Thr Val Asn 290 295 300

As Gly Ser Pro Glu Trp Leu Trp Thr Trp Trp Ile As n Phe Leu Phe 305 310 315 320

Ala Ala Val Val Ala Trp Cys Thr Leu Ile Pro Leu Ser Cys Phe Pro Page 160

		16U	200	PCT	FINAL. ST25
325	330				335

Asn Asn Met Pro Gly Ser Thr Arg Ile Lys Ala Arg Lys Arg Lys Gln 340 345 350

Leu His Phe Phe Asp Ser Arg Leu Lys Asp Leu Lys Leu Gly Thr Asn 355 360 365

Ile Lys Asp Leu Cys Ala Ala Leu Trp Ile Leu Met Arg Asn Pro Val $370 \ 375 \ 380$

Leu Ile Cys Leu Ala Leu Ser Lys Ala Thr Glu Tyr Leu Val Ile Ile 385 \$390\$

Gly Ala Ser Glu Phe Leu Pro Ile Tyr Leu Glu Asn Gln Phe Ile Leu 405 410 415

Thr Pro Thr Val Ala Thr Thr Leu Ala Gly Leu Val Leu Ile Pro Gly $420 \hspace{1cm} 425 \hspace{1cm} 430$

Met Ser Cys Lys Ala Leu Met Arg Phe Ile Met Val Thr Ser Val Ile 450 460

Ser Leu Ile Leu Leu Val Phe Ile Ile Phe Val Arg Cys Asn Pro Val 465 470 475 480

Gln Phe Ala Gly Ile Asn Glu Asp Tyr Asp Gly Thr Arg Lys Leu Gly 485 490 495

Asn Leu Thr Ala Pro Cys Asn Glu Lys Cys Arg Cys Ser Ser Ser Ile 500 505 510

Tyr Ser Ser Ile Cys Gly Arg Asp Asp Ile Glu Tyr Phe Ser Ala Cys 515 520 525

Phe Ala Gly Cys Thr Tyr Ser Lys Ala Gln Asn Gln Lys Lys Met Tyr 530 540

Tyr Asn Cys Ser Cys 11e Lys Glu Gly Leu Ile Thr Ala Asp Ala Glu 545 550 560

Gly Asp Phe Ile Asp Ala Arg Pro Gly Lys Cys Asp Ala Lys Cys Tyr 565 570 575

Lys Leu Pro Leu Phe Ile Ala Phe Ile Phe Ser Thr Leu Ile Phe Ser 580 585 590

Gly Phe Ser Gly Val Pro Ile Val Leu Ala Met Thr Arg Val Val Pro 595 600 605

Asp Lys Leu Arg Ser Leu Ala Leu Gly Val Ser Tyr Val Ile Leu Arg 610 620

Ile Phe Gly Thr Ile Pro Gly Pro Ser Ile Phe Lys Met Ser Gly Glu 625 630 635 640

WO 03/089583 PCT/US03/11497

Thr Ser Cys	Ile Leu Arç 645	Asp Val i		160 200 E Cys Gly I			
Cys Trp Ile	Tyr Asn Lys 660		Met Ala 665	Phe Leu i	Leu Val 670	Gly Ile	
Cys Phe Leu 675	Cys Lys Let	Cys Thr 1	Ile Ile		Thr Ile 685	Ala Phe	
Phe Ile Tyr 690	Lys Arg Arg	Leu Asn (695	Glu Asn	Thr Asp 1	Phe Pro	Asp Val	
Thr Val Lys 705	Asn Pro Lys			Glu Glu 1 715	Thr Asp	Leu	
<210> 248 <211> 851 <212> DNA <213> Homo	sapiens						
<220> <221> CDS <222> (37). <223>	(831)						
<400> 248 gcttctccca c	gctggagtag (ıtgggggagg		atg gag g Met Glu i 1			54
gtc aga tcc Val Arg Ser		Gly Ile I					102
gtg ctg ttg Val Leu Leu 25				Tyr Asn 1			150
aag atc aac Lys Ile Asn 40							198
cgg ggg tcc Arg Gly Ser 55							246
tgc ccc ctg Cys Pro Leu							294
ccc gta tgg Pro Val Trp		Gln Gly 1					342
tgg ggc agc Trp Gly Ser 105				Leu Leu S			390
cag ctg ggc Gln Leu Gly 120							438
ctg cac ggc Leu His Gly 135		Val Ala 1	Leu Leu				486
agg gtg tgt Arg Val Cys							534
aag gcc ttg Lys Ala Leu		Cys Val V					582

aga Arg	cgt Arq	t cto g Lev 185	ı Tyr	tgg Trp	tgg Trp	gtg Val	gaç Glu 190	ı Thr	ato Met	act	acc	cto	acc Thi	IAL.S c tco c Sei	T25 tgg Trp		630
cac His	Let 200	1 Ala	tat Tyr	ctc Leu	atc Ile	acc Thr 205	Trp	acc Thr	acc Thr	tgo Cys	cto Leu 210	Ala	tco Ser	cac His	ctg Leu		678
ctg Leu 215	GIr	g gct n Ala	gcc Ala	ttt Phe	gag Glu 220	His	acg Thr	acc Thr	cag Gln	cto Leu 225	Ala	gaç Glu	g gcc Ala	caç Glr	g gag Glu 230		726
gtt Val	gaa Glu	ccc Pro	cag Gln	gag Glu 235	Val	tca Ser	GJ A	tct Ser	Ser 240	Leu	ctg Leu	cco Pro	tea Ser	Lei 245	tct Ser		774
gcg Ala	tcc Ser	teg Ser	Asp 250	Ser	gag Glu	tct Ser	gga Gly	aca Thr 255	Val	t t g Le u	cca Pro	gag Glu	caa Gln 260	Glu	act Thr		822
		gaa Glu 265		atgt	atc ·	ccca	tctg	cc									851
<21 <21 <21 <21	1 > 2 >	249 265 PRT Homo	sap	iens													
< 40	0>	249															
Met 1	G1 u	Ala	Leu	Pro 5	Pro	Val	Arg	Ser	Ser 10	Leu	Leu	Gly	Ile	Leu 15	Leu		
Gln	Val	Thr	Arg 20	Leu	Ser	Val	Leu	Leu 25	Val	Gln	Asn	Arg	Asp 30	His	Leu		•
Tyr	Asn	Phe 35	Leu	Leu	Leu	Lys	Ile 40	Asn	Leu	Phe	Asn	His 45	Trp	Val	Ser		
Gly	Leu 50	Ala	Gln	Glu	Ala	Arg 55	Gly	Ser	Cys	Asn	Trp 60	Gln	Ala	His	Leu		
Pro 65	Leu	Gly	Ala	Ala	Ala 70	Cys	Pro	Leu	Gly	Gln 75	Ala	Leu	Trp	Ala	61 y		
Leu	Ala	Leu	Ile	Gln 85	Val	Pro	Val	Trp	Leu 90	Val	Leu	Gln	Gly	Pro 95	Arg		
Leu	Met	Trp	Ala 100	Gly	Met	Trp	Gly	Ser 105	Thr	Lys	Gly	Leu	Gly 110	Leu	Ala		
Leu	Leu	Ser 115	Ala	Trp	Glu	Gln	Leu 120	Gly	Leu	Ser	Val	Ala 125	Ile	Trp	Thr		
Asp	Leu 130	Phe	Leu	Ser	Cys	Leu 135	His	Gly	Leu	Met	Leu 140	Val	Ala	Leu	Leu		
Leu 145	Val	Val	Val	Thr	Trp 150	Arg	Val	Cys	Gln	Lys 155	Ser	His	Cys	Phe	Arg 160		
Leu	Gly	Arg	Gln	Leu 165	Ser	Lys	Ala	Leu	Gln 170	Val	Asn	Cys	Val	Val 175	Arg	,	
Lys	Leu	Leu	Val 180	Gln	Leu .	Arg .		Leu 185	Tyr	Trp	Trp	Val	Glu 190	Thr	Met		

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25 Thr Ala Leu Thr Ser Trp His Leu Ala Tyr Leu Ile Thr Trp Thr Thr Cys Leu Ala Ser His Leu Leu Gln Ala Ala Phe Glu His Thr Thr Gln Leu Ala Glu Ala Gln Glu Val Glu Pro Gln Glu Val Ser Gly Ser Ser Leu Leu Pro Ser Leu Ser Ala Ser Ser Asp Ser Glu Ser Gly Thr Val Leu Pro Glu Gln Glu Thr Pro Arg Glu 250 <210> <211> 784 <212> DNA <213> Homo sapiens <220> <221> CDS <222> (97) . . (579) <223> <400> 250 gctttcagtt gtaacggact tcatcacatc acaaattgta ctcgttctca tccttttaag aaagttcaga cccaggaaaa tttccatagt acctta atg aaa aag ata gaa atc Met Lys Lys Ile Glu Ile agt qgg acg tgt ctt tcc ttt cat ctc ctt ttc ggc ttg gaa atc aga Ser Gly Thr Cys Leu Ser Phe His Leu Leu Phe Gly Leu Glu He Arg

60 114 162 atg aga agg att gtt ttt gct ggt gtt atc tta ttc cgc ctc tta ggt Met Arg Arg Ile Val Phe Ala Gly Val Ile Leu Phe Arg Leu Leu Gly 210 gtt atc tta ttc cgc ctc tta ggt gtt atc tta ttc ggc cgc tta ggt Val Ile Leu Phe Arg Leu Leu Gly Val Ile Leu Phe Gly Arg Leu Gly 258 gac ctg gga acc tgc cag aca aaa cct ggt cag tac tgg aaa gaa gag 306 Asp Leu Gly Thr Cys Gln Thr Lys Pro Gly Gln Tyr Trp Lys Glu Glu gtc cac att caa gat gtt gga ggt ttg att tgc aga gca tgc aat ctt Val His Ile Gln Asp Val Gly Gly Leu Ile Cys Arg Ala Cys Asn Leu 354 80 tea etg ecc tte eat gga tgt ett tta gae etg gga acc tge eag gea 402 Ser Leu Pro Phe His Gly Cys Leu Leu Asp Leu Gly Thr Cys Gln Ala gaa cct ggt cag tac tgt aaa gaa gag gtc cac att caa ggt ggc att Glu Pro Gly Gln Tyr Cys Lys Glu Glu Val His Ile Gln Gly Gly Ile 450 caa tgg tat tca gtc aaa ggc tgc aca aag aac aca tca gag tgc ttc Gln Trp Tyr Ser Val Lys Gly Cys Thr Lys Asn Thr Ser Glu Cys Phe 498 aag agt act ctc gtc aag aga att ctg caa ctg cat gaa ctt gta'act Lys Ser Thr Leu Val Lys Arg Ile Leu Gln Leu His Glu Leu Val Thr 546 135 act cac tgc tgc aat cat tct ttg tgc aat ttc tgagtcagtg gcccatatct Thr His Cys Cys Asn His Ser Leu Cys Asn Phe 599

aaaatgcttg gcagatcaat cagtctcgaa gcctgacctg gctatcacaa aatgatggct 659

att	gtca	att	agcc	cact	tc a	gaaa	cete	a ga	ccct	tgta	ggt	agaa	igga	attt	tgatct	719
															atctga	
gaa																784
<21 <21 <21 <21	1> 2>	251 161 PRT Homo	sap	i ens												
< 40	0>	251														
Met 1	Lys	Lys	Ile	Glu 5	lle	Ser	Gly	Thr	Cys 10	Leu	Ser	Phe	His	Leu 15	Leu	
Phe	Gly	Leu	G1u 20	Ile	Arg	Met	Arg	Arg 25	Ile	Val	Phe	Ala	G1 y	Val	Ile	
Leu	Phe	Arg 35	Leu	Leu	G1 y	Val	Ile 40	Leu	Phe	Arg	Leu	Leu 45	Gly	Val	Ile	
Leu	Phe 50	Gly	Arg	Leu	Gly	Asp 55	Leu	Gly	Thr	Cys	Gln 60	Thr	Lys	Pro	G1 y	
Gln 65	Tyr	Trp	Lys	Glu	Glu 70	Val	His	Ile	Gln	Asp 75	Val	Gly	Gly	Leu	Ile 80	
Cys	Arg	Ala	Cys	Asn 85	Leu	Ser	Leu	Pro	Phe 90	His	Gly	Cys	Leu	Leu 95	Asp	
Leu	Gly	Thr	Cys 100	Gln	Ala	Glu	Pro	Gly 105	Gln	Туr	Cys	Lys	Glu 110	Glu	Val	
His	Ile	Gln 115	Gly	G1 y	Ile	Gln	Trp 120	Tyr	Ser	Val	Lys	Gly 125	Cys	Thr	Lys	
Asn	Thr 130	Ser	Glu	Cys	Phe	Lys 135	Ser	Thr	Leu	Val	Lys 140	Arg	Ile	Leu	Gln	
Leu 145	His	Glu	Leu	Val	Thr 150	Thr	His	Cys	Cys	Asn 155	His	Ser	Leu	Cys	Asn 160	
Phe																
<210 <211 <212 <213	> 2 !> I	252 2205 NA Iomo	sapi	ens												
<220 <221 <222 <223	> (DS (24).	. (18	50)												
<400 gggc		:52 ta g	tgca	ggtc	c gc	c at Me 1	g gc t Al	t ga a Gl	g gc u Al	g tc a Se 5	a cg r Ar	g tg g Tr	g ca p Hi	c cg s Ar	a ggc g Gly 10	53
ggg ggg	gct Ala	tcg Ser	Lys	cat His 15	aag Lys	ttg Leu	cat His	Tyr	aga Arg 20	aag Lys	gaa Glu	gta Val	Glu	att Ile 25	aca Thr	101
acc	aca	ctt	cag	gaa	ttg	tta	ctc	tac	ttt .	att		tta age :		aac	cta	149

Thr	Thr	Leu	Gln 30	Glu	Leu	Leu	Leu	Tyr 35	Phe		FINA 11e 40		
									cca Pro				197
									act Thr				245
									ata Ile				293
									tac Tyr 100				341
									agc Ser				389
					-		-		cgt Arg		-		437
									cag Gln				485
						-		_	gac Asp				533
									act Thr 180				581
									cga Arg				629
									aaa Lys				677
									act Thr				725
	_				-		-		cta Leu	-	_	-	773
									ata 11e 260				821
									agc Ser				869
									ttt Phe				917
									aag Lys				965
									ttg Leu				1013
									caa Gln 340				1061

Page 166

cag Gln	ctg Leu	ttg Leu	aaa Lys 350	agt Ser	act Thr	gaa Glu	aaa Lys	tat Tyr 355	tca Ser	gat	200 ttc Phe	tat	ttt	ctt Leu	aca	1109
tgc Cys	tgg Trp	cac His 365	att Ile	tat Tyr	tac Tyr	aat Asn	aat Asn 370	ata Ile	att Ile	gct Ala	att Ile	acc Thr 375	atc Ile	ttt Phe	ttt Phe	1157
gca Ala	tgg Trp 380	ata Ile	aag Lys	ata Ile	ttc Phe	aaa Lys 385	ttc Phe	ata Ile	agc Ser	ttt Phe	aac Asn 390	aag Lys	aca Thr	atg Met	tct Ser	1205
cag Gln 395	ctg Leu	tca Ser	tca Ser	acc Thr	ttg Leu 400	tcc Ser	cgt Arg	tgt Cys	gtt Val	aaa Lys 405	gac Asp	ata Ile	gta Val	gga Gly	ttt Phe 410	1253
gcc Ala	atc Ile	atg Met	ttt Phe	ttt Phe 415	ata Ile	ata Ile	ttc Phe	ttt Phe	gct Ala 420	tat Tyr	gcc Ala	cag Gln	tta Leu	gga Gly 425	ttt Phe	1301
ctt Leu	gtt Val	ttt Phe	gga Gly 430	tca Ser	caa Gln	gtt Val	gat Asp	gac Asp 435	ttt Phe	tcc Ser	act Thr	ttt Phe	cag Gln 440	aat Asn	tcc Ser	1349
ata Ile	ttt Phe	gca Ala 445	caa Gln	ttt Phe	cga Arg	att Ile	gtt Val 450	ctt Leu	gga Gly	gat Asp	ttt Phe	aat Asn 455	ttt Phe	gct Ala	ggt Gly	1397
att Ile	cag Gln 460	caa Gln	gcc Ala	aat Asn	cct Pro	atc Ile 465	ttg Leu	gga Gly	ccc Pro	att Ile	tac Tyr 470	ttc Phe	atc Ile	act Thr	ttc Phe	1445
atc Ile 475	ttt Phe	ttt Phe	gtg Val	ttc Phe	ttt Phe 480	gtc Val	ctg Leu	ctg Leu	aat Asn	atg Met 485	ttc Phe	ttg Leu	gca Ala	att Ile	att Ile 490	1493
aat Asn	gat Asp	acc Thr	tat Tyr	tct Ser 495	gaa Glu	gtg Val	aaa Lys	gct Ala	gac Asp 500	tat Tyr	tca Ser	ata Ile	ggc Gly	aga Arg 505	agg Arg	1541
cca Pro	gat Asp	ttt Phe	gaa Glu 510	ctt Leu	ggc Gly	aaa Lys	atg Met	att Ile 515	aaa Lys	cag Gln	agt Ser	tac Tyr	aaa Lys 520	aat Asn	gtt Val	1589
ctc Leu	gag Glu	aaa Lys 525	ttc Phe	aga Arg	ctg Leu	aag Lys	aaa Lys 530	gct Ala	caa Gln	aaa Lys	gat Asp	gaa Glu 535	gac Asp	aag Lys	aaa Lys	1637
acc Thr	aaa Lys 540	ggc Gly	agc Ser	gga Gly	gat Asp	ttg Leu 545	gct Ala	gaa Glu	caa Gln	gcc Ala	aga Arg 550	aga Arg	gaa Glu	ggc ggc	ttt Phe	1685
gac Asp 555	gaa Glu	aat Asn	gag Glu	att Ile	caa Gln 560	aac Asn	gca Ala	gag Glu	cag Gln	atg Met 565	aaa Lys	aaa Lys	tgg Trp	aaa Lys	gag Glu 570	1733
agg Arg	ctt Leu	gag Glu	Lys	aag Lys 575	tat Tyr	tat Tyr	tct Ser	atg Met	gaa Glu 580	att Ile	caa Gln	gat Asp	gac Asp	tac Tyr 585	cag Gln	1781
cct Pro	gtc Val	Thr	caa Gln 590	gaa Glu	gaa Glu	ttt Phe	Arg	gat Asp 595	ggc Gly	acc Thr	aca Thr	Thr	aag Lys 600	tac Tyr	aaa Lys	1829
atg Met	aga Arg	ttc Phe 605	tct Ser	ctg Leu	agt Ser	gcc Ala	tgac	aaaa	cg a	attt	aagt	a cc	agcc	aagt		1880
acac	acga	tg a	tagc	ttca	a gg	aata	caac	tga	cttt	atg .	atat	gaat	tt t	caag	gaacg	1940
tatc	ttat	at g	gatt	ttga	a ga	atct	tgtt	tgc	ttat	aag	aact	tcaa	ga a	gcct	aagct	2000
tggc	ttta	at t	ttct	tgta	c tc	tctg	tact	cct	caag	cac	tgga	acac	ga t	cctç	tttct	2060
gggc	attc	ct a	3 999	agaa	a at	aaaat	tttg	taa	tgtt	cta (gaga	tcat	tt g	gaaa	aaaag	2120
atcc	aaaa	gt to	gtct	taat	a tg	agaca	atac	tgt	tact	aaa (cata	agtt	ca a	ataa	aaagt	2180
tgtt	ctga	aa a	aaaa	aaaa	a aa	aaa										2205

Page 167

WO 03/089583 PCT/US03/11497

16U 200 PCT FINAL. ST25

<210> 253 <211> 609 PRT <213> Homo sapiens <400> 253 Met Ala Glu Ala Ser Arg Trp His Arg Gly Gly Ala Ser Lys His Lys 1 10 15 Leu His Tyr Arg Lys Glu Val Glu Ile Thr Thr Thr Leu Gln Glu Leu $20 \hspace{1cm} 25 \hspace{1cm} 30 \hspace{1cm}$ Leu Leu Tyr Phe lle Phe Leu Ile Asn Leu Cys Ile Leu Thr Phe Gly 35 40 45 Met Val Asn Pro His Met Tyr Tyr Leu Asn Lys Val Met Ser Ser Leu 50 60Phe Leu Asp Thr Ser Val Pro Gly Glu Glu Arg Thr Asn Phe Lys Ser 65 70 75 80Ile Arg Ser Ile Thr Asp Phe Trp Lys Phe Met Glu Gly Pro Leu Leu $85 \hspace{1.5cm} 90 \hspace{1.5cm} 95$ Glu Gly Leu Tyr Trp Asp Ser Trp Tyr Asn Asn Gln Gln Leu Tyr Asn 100 105 110 Leu Lys Asn Ser Ser Arg Ile Tyr Tyr Glu Asn Ile Leu Leu Gly Val 115 120 125Pro Arg Val Arg Gln Leu Lys Val Arg Asn Asn Thr Cys Lys Val Tyr 130 135 140 Ser Ser Phe Gln Ser Leu Met Ser Glu Cys Tyr Gly Lys Tyr Thr Ser 145 150 155 160 Ala Asn Glu Asp Leu Ser Asn Phe Gly Leu Gln Ile Asn Thr Glu Trp 165 170 175Arg Tyr Ser Thr Ser Asn Thr Asn Ser Pro Trp His Trp Gly Phe Leu 180 185 190 Gly Val Tyr Arg Asn Gly Gly Tyr Ile Phe Thr Leu Ser Lys 195 200 205Ser Glu Thr Lys Asn Lys Phe Ile Asp Leu Arg Leu Asn Ser Trp Ile 210 215 220 Thr Arg Gly Thr Arg Val Ile Phe Ile Asp Phe Ser Leu Tyr Asn Ala 225 230 235 240

Asn Val Asn Leu Phe Cys Ile Ile Arg Leu Val Ala Glu Phe Pro Ala 245 250 250 255 $^{\prime\prime}$

Thr Gly Gly Ile Leu Thr Ser Trp Gln Phe Tyr Ser Val Lys Leu Leu 260 265 270

Arg Tyr Val Ser Tyr Tyr Asp Tyr Phe Ile Ala Ser Cys Glu Ile Thr 275 280 285

- Phe Cys Ile Phe Leu Phe Val Phe Thr Thr Gln Glu Val Lys Lys Ile 290 295 300
- Lys Glu Phe Lys Ser Ala Tyr Phe Lys Ser Ile Trp Asn Trp Leu Glu 305 310 315 320
- Leu Leu Leu Leu Cys Phe Val Ala Val Ser Phe Asn Thr Tyr 325 330 335
- Tyr Asn Val Gln Ile Phe Leu Leu Leu Gly Gln Leu Leu Lys Ser Thr 340 345 350
- Asn Asn Ile Ile Ala Ile Thr Ile Phe Phe Ala Trp Ile Lys Ile Phe $370 \hspace{1.5cm} 375 \hspace{1.5cm} 380$
- Lys Phe Ile Ser Phe Asn Lys Thr Met Ser Gln Leu Ser Ser Thr Leu 385 390 395 400
- Ser Arg Cys Val Lys Asp Ile Val Gly Phe Ala Ile Met Phe Phe Ile 405 415
- Ile Phe Phe Ala Tyr Ala Gln Leu Gly Phe Leu Val Phe Gly Ser Gln $420 \hspace{1cm} 425 \hspace{1cm} 430 \hspace{1cm}$
- Val Asp Asp Phe Ser Thr Phe Gln Asn Ser Ile Phe Ala Gln Phe Arg 435 440 445
- lle Val Leu Gly Asp Phe Asn Phe Ala Gly Ile Gln Gln Ala Asn Pro 450 455 460
- Ile Leu Gly Pro Ile Tyr Phe Ile Thr Phe Ile Phe Phe Val Phe Phe 475 480
- Val Leu Leu Asn Met Phe Leu Ala Ile Ile Asn Asp Thr Tyr Ser Glu 485 490 495
- Val Lys Ala Asp Tyr Ser Ile Gly Arg Arg Pro Asp Phe Glu Leu Gly 500 500 510
- Lys Met Ile Lys Gln Ser Tyr Lys Asn Val Leu Glu Lys Phe Arg Leu 515 520 525
- Lys Lys Ala Gln Lys Asp Glu Asp Lys Lys Thr Lys Gly Ser Gly Asp 530 535 540
- Leu Ala Glu Gln Ala Arg Arg Glu Gly Phe Asp Glu Asn Glu Ile Gln 545 550 555 560
- Asn Ala Glu Gln Met Lys Lys Trp Lys Glu Arg Leu Glu Lys Lys Tyr 565 575 575
- Tyr Ser Met Glu Ile Gln Asp Asp Tyr Gln Pro Val Thr Gln Glu Glu 580 585 590
- Phe Arg Asp Gly Thr Thr Lys Tyr Lys Met Arg Phe Ser Leu Ser Page 169

16U 200 PCT FINAL.ST25 595 600 605

vra																	
<210 <211 <211 <211	l > 2 >	254 1615 DNA Homo	sap	iens													
<220 <221 <221 <221	1 > 2 >	CDS (1).	. (15	84)													
	gcc								atg Met 10								48
									t t c Phe								96
									cac His							:	144
									ttg Leu							1	192
									ata Ile							2	240
									cgc Arg 90							2	288
									gag Glu							3	336
									acg Thr							3	384
									aaa Lys							4	132
									ttg Leu							4	180
									tct Ser 170							5	28
									atg Met							5	76
									caa Gln							6	524
									ctc Leu							· 6	72
									ctg Leu							7	20
atg	acc	ttc	ctc	ttg	atg	ttg	ctg	ctc	atc	ttc	ttc	tac	att	ttt	gct	7	68

Page 170

Ala

Met	Thr	Phe	Leu	Let 245	Met	Leu	Leu	. Leu	11e	Phe	200 Phe	PCT Tyr	FIN Ile	AL.S Phe 255	e Ala	
gtg Val	act Thr	ggt Gly	gtc Val 260	tac Tyr	gtc Val	Phe	tca Ser	gag Glu 265	Tyr	acc Thi	cgt Arg	tca Ser	cct Pro 270	Arg	cag Gln	816
gac Asp	ctg Leu	gag Glu 275	tac Tyr	cat His	gtg Val	ttc Phe	ttc Phe 280	Ser	gac Asp	Leu	ccg Pro	aat Asn 285	Ser	ctg Leu	gta Val	864
aca Thr	gtg Val 290	Phe	att Ile	ctc Leu	ttc Phe	acc Thr 295	Leu	gat Asp	cat His	tgg Trp	tat Tyr 300	Ala	ctg Leu	ctt Leu	cag Gln	912
gac Asp 305	gtc Val	tgg Trp	aag Lys	gtg Val	cct Pro 310	Glu	gt <i>c</i> Val	agt Ser	cgc Arg	ato Ile 315	Phe	agc Ser	agc Ser	ato	tat Tyr 320	960
ttc Phe	atc Ile	ctt Leu	tgg Trp	ttg Leu 325	ttg Leu	ctt Leu	ggc Gly	tcc Ser	att 11e 330	atc Ile	ttt Phe	cga Arg	agt Ser	atc Ile 335	ata Ile	1008
gta Val	gcc Ala	atg Met	atg Met 340	gtt Val	act Thr	aac Asn	ttt Phe	cag Gln 345	aat Asn	atc Ile	agg Arg	aaa Lys	gag Glu 350	ctg Leu	aat Asn	1056
gag Glu	gag Glu	atg Met 355	gcg Ala	cgt Arg	cgg Arg	gag Glu	gtt Val 360	cag Gln	ctc Leu	aaa Lys	gct Ala	gac Asp 365	atg Met	ttc Phe	aag Lys	1104
cgg Arg	cag Gln 370	atc Ile	atc Ile	cag Gln	agg Arg	aga Arg 375	aaa Lys	aac Asn	atg Met	tca Ser	cat His 380	gaa Glu	gca Ala	ctg Leu	acg Thr	1152
tca Ser 385	agc Ser	cat His	agc Ser	aaa Lys	ata Ile 390	gag Glu	gac Asp	aga Arg	gga Gly	gct Ala 395	agt Ser	caa Gln	caa Gln	agg Arg	gaa Glu 400	1200
agt Ser	ttg Leu	gac Asp	tta Leu	tca Ser 405	gaa Glu	gtg Val	tct Ser	gaa Glu	gta Val 410	gag Glu	tct Ser	aat Asn	tat Tyr	ggt Gly 415	gcc Ala	1248
act Thr	gaa Glu	gag Glu	gat Asp 420	tta Leu	ata Ile	aca Thr	tct Ser	gca Ala 425	tca Ser	aaa Lys	aca Thr	gaa Glu	gag Glu 430	acc Thr	ttg Leu	1296
tca Ser	aaa Lys	aag Lys 435	aga Arg	gag Glu	tac Tyr	cag Gln	tct Ser 440	tcc Ser	tcc Ser	tgt Cys	gtc Val	tcc Ser 445	tcc Ser	aca Thr	tcc Ser	1344
tct Ser	tcc Ser 450	tat Tyr	tct Ser	tcc Ser	tct Ser	tct Ser 455	gaa Glu	tcc Ser	aga Arg	ttt Phe	tct Ser 460	gaa Glu	tct Ser	att Ile	ggt Gly	1392
cgt Arg 465	ttg Leu	gac Asp	tgg Trp	gag Glu	act Thr 470	ctt Leu	gtg Val	cac His	gaa Glu	aat Asn 475	ctg Leu	ccc Pro	G 1y	cta Leu	atg Met 480	1440
gaa Glu	atg Met	gat Asp	Gln	gat Asp 485	gac Asp	cgt Arg	gtt Val	tgg Trp	ccc Pro 490	aga Arg	gac Asp	tca Ser	ctc Leu	ttc Phe 495	cga Arg	1488
tat Tyr	ttt Phe	Glu	ttg Leu 500	cta Leu	gaa Glu	aag Lys	ctt Leu	cag Gln 505	tat Tyr	aac Asn	cta Leu	gag Glu	gaa Glu 510	cgt Arg	aag Lys	1536
aag Lys	Leu	caa Gln (515	gag Glu	ttt Phe	gca Ala	Val	cag Gln 520	gca Ala	ctg Leu	atg Met	aac Asn	ttg Leu 525	gaa Glu	gac Asp	aag Lys	1584
taaa	gcaa	tg g	atgg	cttc	a at	atcc	ttgg	g						, 1		1615
<210: <211: <212: <213:	> 5: > Pi	55 28 RT omo :	sapi	ens											-	·

Page 171

WO-03/089583 PCT/US03/11497

16U 200 PCT FINAL.ST25

<400> 255

Met Ala Ala Tyr Gln Gln Glu Glu Gln Met Gln Leu Pro Arg Ala Asp 1 5 10 15

Ala Ile Arg Ser Arg Leu Ile Asp Thr Phe Ser Leu Ile Glu His Leu 20 25 30

Gln Gly Leu Ser Gln Ala Val Pro Arg His Thr Ile Arg Glu Leu Leu 35 45

Asp Pro Ser Arg Gln Lys Lys Leu Val Leu Gly Asp Gln His Gln Leu 50 55 60

Val Arg Phe Ser Ile Lys Pro Gln Arg Ile Glu Gln Ile Ser His Ala 65 70 75 80

Gln Arg Leu Leu Ser Arg Leu His Val Arg Cys Ser Gln Arg Pro Pro 85 90 95

Leu Ser Leu Trp Ala Gly Trp Val Leu Glu Cys Pro Leu Phe Lys Asn 100 105 110

Phe Ile Ile Phe Leu Val Phe Leu Asn Thr Ile Ile Leu Met Val Glu 115 120 125

Ile Glu Leu Leu Glu Ser Thr Asn Thr Lys Leu Trp Pro Leu Lys Leu 130 135 140

Thr Leu Glu Val Ala Ala Trp Phe Ile Leu Leu Ile Phe Ile Leu Glu 145 150 155 160

lle Leu Leu Lys Trp Leu Ser Asn Phe Ser Val Phe Trp Lys Ser Ala 165 170 175

Trp Asn Val Phe Asp Phe Val Val Thr Met Leu Ser Leu Leu Pro Glu 180 185 190

Val Val Leu Val Gly Val Thr Gly Gln Ser Val Trp Leu Gln Leu 195 200 205

Leu Arg Ile Cys Arg Val Leu Arg Ser Leu Lys Leu Leu Ala Gln Phe 210 215 220

Arg Gln Ile Gln Ile Ile Ile Leu Val Leu Val Arg Ala Leu Lys Ser 225 230 235 240

Met Thr Phe Leu Leu Met Leu Leu Leu Ile Phe Phe Tyr Ile Phe Ala 245 250 255

Val Thr Gly Val Tyr Val Phe Ser Glu Tyr Thr Arg Ser Pro Arg Gln 260 265 270

Asp Leu Glu Tyr His Val Phe Phe Ser Asp Leu Pro Asn Ser Leu Val 275 280 285

Thr Val Phe Ile Leu Phe Thr Leu Asp His Trp Tyr Ala Leu Leu Gln 290 295 300

Asp Val Trp Lys Val Pro Glu Val Ser Arg Ile Phe Ser Ser Ile Tyr Page 172

16U 200 PCT FINAL.ST25 305 310 -Phe Ile Leu Trp Leu Leu Gly Ser Ile Ile Phe Arg Ser Ile Ile 325 330 335 Val Ala Met Met Val Thr Asn Phe Gln Asn Ile Arg Lys Glu Leu Asn Glu Glu Met Ala Arg Arg Glu Val Gln Leu Lys Ala Asp Met Phe Lys Arg Gln Ile Ile Gln Arg Arg Lys Asn Met Ser His Glu Ala Leu Thr 370 375 380Ser Ser His Ser Lys Ile Glu Asp Arg Gly Ala Ser Gln Gln Arg Glu 385 390 395 . 400 Ser Leu Asp Leu Ser Glu Val Ser Glu Val Glu Ser Asn Tyr Gly Ala 405 410 415Thr Glu Glu Asp Leu Ile Thr Ser Ala Ser Lys Thr Glu Glu Thr Leu $420 \hspace{1.5cm} 425 \hspace{1.5cm} 430$ Ser Lys Lys Arg Glu Tyr Gln Ser Ser Ser Cys Val Ser Ser Thr Ser 435 440 445Ser Ser Tyr Ser Ser Ser Ser Glu Ser Arg Phe Ser Glu Ser Ile Gly Arg Leu Asp Trp Glu Thr Leu Val His Glu Asn Leu Pro Gly Leu Met 465 470 475 480 Tyr Phe Glu Leu Glu Lys Leu Gln Tyr Asn Leu Glu Glu Arg Lys Lys Leu Gln Glu Phe Ala Val Gln Ala Leu Met Asn Leu Glu Asp Lys <210> 256 <211> 24 <212> DNA <213> Homo sapiens <400> 256 tcatggatca ccagctccac gctc 24 <210> 257 <211> 25 <212> DNA <213> Homo sapiens <400> 257 caccaagatc accaccatgg aagca <210> 258 <211> 50 <212> DNA <213> Homo sapiens <400> 258

WO 03/089583 PCT/US03/11497

				200 PCT FIN	IAL. ST25	
ggattc	aggc cttttaaacc	ccactcagtg	ggtgcatggc	agggctttga		50
-210 5	050					
<210> <211>	259 24					
<212>	DNA					
<213>						
<400>	259				•	
tgctga	cgaa tcttatgaac	cagg				24
<210>	260					
<211>	26					
<212> <213>	DNA Homo sapiens					
<400>	260					
tcacgt	cage cteteettee	tcagtg				26
<210>	261					
<211> <212>	50 DNA					
<213>	Homo sapiens					
<400>	261					
ccacaa	atca tataaattag	gggaaagaga	gaggcaggca	Caccecaaaa		50
	0.00					
<210> <211>	262 50					
<211>	DNA					
<213>	Homo sapiens					
<400>	262					
	ttat ttaaaagacc	tcagaaatgt	caccatactt	agttatttta		50
	,			agotto cocca		-
<210>	263					
<211>	23					
<212>	DNA					
<213>	Homo sapiens					
<400>	263					
ggccate	gac aatgtcacag	cag				23
<210>	264					
<211>	31					
<212>	DNA					
<213>	Homo sapiens					
<400>	264					
agcagac	aca tactgggcca	ttcataacca	С			31
<210>	265					
<211>	50					
<212> <213>	DNA Homo sapiens					
<400>	265					
ggtacta	ittc tatattttgg	gcacacagca	atgaagaaaa	cagaaaaacc	•	50
<210>	266					
<211> <212>	50 DNA					
<212>	Homo sapiens					
					•	
<400>	266	~~~~	***	*****		E 0
ceggget	tca taaatatgga	ycagaaagtt	tttacaaata	cagaacagca		50
-23.0 :	262					
<210> <211>	267 50					
<211>	DNA					
	Homo sapiens					

Page 174

<400> tagaa	267 tgtgt tataaaaaat gaagcagggc taggggaa	aag agatgggtga	50
<210> <211> <212> <213>	23		
<400> cctcat	268 ttggc ttcctcccac tcg		23
<210><211><211><212><213>	3'0 DNA		
<400> gccato	269 caaac tetgagetgg agatagtgae		30
<210><211><211><212><213>	50 DNA		
<400> ccaagg	270 gaact tttaaaactc ccattgcaca gttaccac	cc agaataatta	50
<210><211><211><212><213>	271 50 DNA Homo sapiens		
<400> catcct	271 Eggaa tatatttgcg tccaactctg caccttgo	tc tctattccct	50
<210><211><211><212><213>	272 50 DNA Homo sapiens		
<400> ctgggg	272 genee teaaaaaget eacetteeet eactteee	ac ttcaactgat	50
<210><211><211><212><213>	273 26 DNA Homo sapiens		
<400> tggcct	273 cgtt gaaagtgtca tcatcc		26
<210><211><211><212><213>	274 24 DNA Homo sapiens		
<400> ttggta	274 ccat ttacgaatgg ccgc		24
<210> <211> <212>	275 50 DNA		
<213> <400> aaacgg	Homo sapiens 275 catt ttaaaaatgc aggtttaaat tgttatcc	tc atctatggtt	50
<210> <211>	276 29		

Page 175

WO 03/089583 PCT/US03/11497

<212>	16U 200 PCT FINAL.ST	25
<213>	Homo sapiens	
<400>	276	
ctggaci	cttga gcagtaccac gtctggatc	29
<210>	277	
<211>		
<212> <213>	DNA Homo sapiens	
<400>	277	
	tccca cagcaatttt gacaatgg	28
/210 >	270	
<210> <211>	278 50	
<212>	DNA	
<213>	Homo sapiens	
<400>	278	
attttgg	ggita tatatagagg agictaggaa aagactcgig ggicigalic	50
<210>	279	
<211>		
<212> <213>	DNA Home canions	
(213)	Homo sapiens	
<400>	279	5.0
tactca	atatt tatatagcag caacttacat tgacccaggg agaactcagt	50
<210>	280	
<211>		
<212> <213>	DNA Homo sapiens	
<400>	200	
<400> gttacco	280 ccacc caaccgteac gacc	24
<210>	281	
<211> <212>	24	
<213>	DNA Homo sapiens	
<400>	281	
caggcga	gatgc cagagaagac gatg	24
<210>	202	
<210> <211>	282 50	
<212>	DNA	
<213>	Homo sapiens	
<400>	282	
ctagaat	attta cataaaaagg actggaggag cttttgcagc aactttgcat	50
<210>	203	
<210>	203 50	
<212>	DNA	
<213>	Homo sapiens	
<400>	283	
LLLCCC	ttctt ttaaaaacac gctttcactc tcaaaacagc agagaatgaa	50
<210>	284	•
<211>	50 .	
<212> <213>		
-C132	Homo sapiens	

<400> 284 aactggggtc tataagagag ccagggcact tattcatcca agggcagatg

Page 176

50

16U 200 PCT FINAL.ST25 <210> 285 <211> 26 <212> DNA <213> Homo sapiens <400> 285 26 ctatgacttc aacccacacc tgggca <210> 286 <211> 24 <212> DNA <213> Homo sapiens <400> 286 24 aaggtcgcca acttgtcctg gctc <210> 287 <211> 50 <212> DNA <213> Homo sapiens <400> 287 50 gggcgggagt aaaaggcaga gtccaattcc accggccccc agtgtgggtg <210> 288 <211> 24 <212> DNA <213> Homo sapiens <400> 288 24 tcaatgccat gcccaaactg agga <210> 289 <211> 24 <212> DNA <213> Homo sapiens <400> 289 24 caacaccgag atggacaccc tgct <210> 290 <211> 50 <212> DNA <213> Homo sapiens 50 cttttaaggt taaaaatgtg ggttttagat gattgtcctt tctaaacagc <210> 291 <211> 50 <212> DNA <213> Homo sapiens 50 tcaggatgtc taaaaaagat ctctctagtg tacacacgtg cacacacaca <210> 292 <211> 50 <212> DNA <213> Homo sapiens <400> 292 agtaactcta tttaaaagac ctaaaaaattt caaatcctaa aatgatctat . 50 <210> 293 <211> 5.0 <212> DNA <213> Homo sapiens <400> 293 aataaatgtt ttaaaagcac tcctttccga atggtggagc tggtggggc 50

<210>	294		
<211>	27		
<212>	DNA		
<213>	Homo sapiens		
<400>	294		
ctcagga	acga agatcatgat cggcatc		27
<210>	295		
<211>	28		
<212>	DNA		
<213>	Homo sapiens		
<400>	295		20
gaagatt	ttt gtgcccagct ttcccaag		28
<210>	296		
<211>	50		
<212>	DNA		
<213>	Homo sapiens		
	204		
<400>	296		50
tattete	cact tataagtggg agctaagcca tgagggcacc aaggcataag		30
4310 5	207		
<210>	297		
<211>	50		
<212>	DNA Name canions		
<213>	Homo sapiens		
<400>	297		
	atgt atacatgtgc catgctggtg tgctgcaccc attaactcgt		50
ccacac	acyc acacacycyc cacycogycy cyaryaraaa arranaaa		
<210>	298		
<211>	27		
<212>	DNA		
<213>	Homo sapiens		
	•		
< 400>	298		
tccatg	ctca gcttcatctc agctacc		27
<210>	299		
<211>	24		
<212>	DNA		
<213>	Homo sapiens		
<400>	299		24
tccatc	tcag accttggccc ttca		24
Z210×	200		
<210> <211>	300 50		
<212> <213>	DNA Homo sanjens		
\L13/	Homo sapiens		
<400>	300		
	cccc attaaaaagt gggcaaaggg catgaacact tttcaaaaga		50
<210>	301		
<211>	24		
<212>	DNA		
<213>	Homo sapiens		
<400>	301	i	
	gtaa ggagccatcg gaca		24
		,:	
<210>	302		
<211>	23		
<212>	DNA		
<213>	Homo sapiens		

Page 178

16U 200 PCT FINAL.ST25 <400> 302 cttgccaggt tctggtggct tgg 23 <210> <211> 303 50 <212> DNA <213> Homo sapiens <400> 303 tctttttgtc tataaatagg actttgattt tctggactag agaattgtat 50 <210> 304 <211> 24 <212> DNA <213> Homo sapiens <400> 304 acgaetecaa gaacageaag geeg 24 <210> 305 <211> 24 <212> DNA <213> Homo sapiens <400> 305 aaggtaacat cggcagaggc cagc 24 <210> 306 <211> 50 <212> DNA <213> Homo sapiens <400> 306 gctagcattt tttaaaagct gatgtettea etgggeaegg ggaeteaeae 50 <210> 307 <211> 24 <212> DNA <213> Homo sapiens <400> 307 cggccaggta ccaaagctca gctg 24 <210> 308 <211> 29 <212> DNA <213> Homo sapiens <400> 308 gccagattca ggagggaatg gaagagaac 29 <210> 309 <211> 50 <212> DNA <213> Homo sapiens <400> 309 tgatctactt tttaaaagga tcatgctggc tgctggtggg atttaggata 50 <210> 310 <211> 50 <212> DNA <213> Homo sapiens tgatagtgat aaaaaaaagt ggccagattt tggttatatt ttgaaataaa 50 <210> 311 <211> 50 <212> DNA


Page 179

	160	200 PCT FINAL.ST25	
<213> Homo sapiens			
<400> 311			
	ggggtctggg tgagataact	gatggaatga	50
grade	9999101999 1929211111	3333-	
<210> 312			
<211> 50			
<212> DNA			
<213> Homo sapiens			
<400> 312			
	ggagtcatta aaatggtgct	aagaagctga	50
	33-3	3 3 3	
<210> 313			
<211> 50			
<212> DNA			
<213> Homo sapiens			
<400> 313			
	gtgctaagaa gctgagctac	aagcagtggt	50
-5-9599-5	3-3	3 3 33	
<210> 314			
<211> 50			
<212> DNA			
<213> Homo sapiens			
<400> 314			
	ccactggatg aagtcccctc	cttccattaa	50
gacacceae ecadadacy	country adjusted		
<210> 315			
<211> 26			
<212> DNA			
<213> Homo sapiens			
<400> 315			
ttgggagaga ctagtgcacc	teagea		26
cegggagaga ceagegeace	ceagea		
<210> 316			
<211> 24			
<212> DNA			
<213> Homo sapiens			
<400> 316			
gagcaatece tettegtgge	aggt		24
gageaatece terregrage	. agge		
<210> 317			
<211> 50	•		
<212> DNA			
<213> Homo sapiens			
<400> 317			
	ı ggggtggagg ggcttatgag	aaggggacca	50
addagtgeet teadacaggg	, ggggeggagg ggeeeurgag	, 229999222	
<210> 318			
<211> 50	~		
<212> DNA			
<213> Homo sapiens			
<400> 318			
	gagatcagcc aggcgtggca	catacctata	50
continue taaaaatyca	. gagareagee aggegryges		
<210> 319			
<211> 50		,	
<212> DNA			
<213> Homo sapiens			
(400) 310			
<400> 319		tactagatac	50
aaaaaaaaa aaaaaaagcc	ctgtttatat cctacctcct	. Lyclyydyc	50
<210> 320			

16U 200 PCT FINAL.ST25 <211> 50 <212> DNA <213> Homo sapiens <400> 320 50 aaaataaaaa taaaaaatcc catctcctca catttccatt caacctcaat <210> 321 <211> 35 <212> DNA <213> Homo sapiens 35 acttccaaac atctacaact cetcagagte teatt <210> 322 <211> 25 <212> DNA <213> Homo sapiens <400> 322 tgcagcacca tcatgtaagg gacaa 25 <210> 323 <211> 50 <212> DNA <213> Homo sapiens <400> 323 50 tttttaaac tataaaaagt ggggatcaga aaacacagtc ataagggaaa <210> 324 <211> 50 <212> DNA <213> Homo sapiens <400> 324 50 gtatatgcta tatatatcag gattcacttt aatggcattg agttccagga <210> 325 <211> 50 <212> DNA <213> Homo sapiens <400> 325 ataaacaatt taaaaattag cccaccatgg tggtacacac ctgtcgttct 50 <210> 326 <211> 50 <212> DNA <213> Homo sapiens <400> 326 50 aaaaaagtgaa aaaaaaaggt gagggagact ttaactttct gaaatatatt <210> 327 <211> 24 <212> DNA <213> Homo sapiens <400> 327 24 ccaagaagcc gggagaagtg gatg <210> 328 <211> 28 <212> DNA <213> Homo sapiens <400> 328 28 tgacagagct aggcatatga gcactgga

```
<210> 329
<211> 50
<212> DNA
<213> Homo sapiens
<400> 329
                                                                        50
ctaaagagct tatatatcag cctaagaaaa gaaaaccaat aagaagttgc
<210> 330
<211> 26
<212> DNA
<213> Homo sapiens
<400> 330
                                                                        26
gcagttggtt cagaaccgag atcacc
<210> 331
<211> 29
<212> DNA
<213> Homo sapiens
<400> 331
ggcagatggg gatacattta ttctctggg
                                                                        29
<210> 332
<211>
       50
<212> DNA
<213> Homo sapiens
<400> 332
actaaaaata caaaaaagta gccgggtatg gtggtaggcg cctataatcc
                                                                        50
<210> 333
<211> 50
<212> DNA
<213> Homo sapiens
ggtaggcgcc tataatccca gctacttggg aggctgaggc aggagaattg
                                                                        50
<210> 334
<211> 26
<212> DNA
<213> Homo sapiens
<400> 334
tcggcttgga aatcagaatg agaagg
                                                                        26
<210> 335
<211> 30
<212> DNA
<213> Homo sapiens
<400> 335
                                                                        30
tgcacaaaga atgattgcag cagtgagtag
<210>
       336
<211> 50
<212> DNA
<213> Homo sapiens
                                                                   50
<210>
       337
<211>
       50
<212>
       DNA
<213> Homo sapiens
<400> 337
```

WO 03/089583



ggcca	16U 200 PCT FINAL.ST25 actta tataaaaggt ttatgttttt gttctgataa tttcgtttct	50
<210> <211> <212> <213>		
<400> aagtta	338 aagtt ttaaaaagaa caggctácaa agttatagct atggggtgat	50
<210> <211>	339 21	
<212> <213>	DNA	
<400>	339	
	tgta gtgcaggtcc g	21
<210>	340	
<211>	24	
<212> <213>	DNA Name Cambana	
	Homo sapiens	
<400>	340	
cccca	gttg cagggaattc tgcc	24
<210>	341	
<211> <212>	50 DNA	
<213>	Homo sapiens	
<400>	341	
aattca	aata tttaaaacgg actgtctcct cttcacaaaa gtctagatct	50
<210>	342	
<211>	24	
<212>	DNA	
<213>	Homo sapiens	
<400>	342	
ggctgt	tgag caggetteat gtge	24
<210>	343	
<211>	24	
<212> <213>	DNA	
	Homo sapiens	
<400>	343	
	tgga tgatetgeeg ettg	24
<210>	344	
<211>	50	
<212> <213>	DNA Homo sapiens	
<400> attqqqt	344 gca tatatattta ggatagttag ctcttcttgt tgaattgatc	50
. ,,,,		วบ